


2,4,6-TRICHLOROPHENOL

CAS No. 88-06-2

First Listed in the *Third Annual Report on Carcinogens*



CARCINOGENICITY

2,4,6-Trichlorophenol is *reasonably anticipated to be a human carcinogen* based on limited evidence for the carcinogenicity of combined exposures to polychlorophenols, including 2,4,6-trichlorophenol, in humans and sufficient evidence for the carcinogenicity of 2,4,6-trichlorophenol in experimental animals. Several cohort studies of workers involved in chemical manufacture of chlorophenols and in leather tanneries and sawmills where chlorophenols were used indicated a relationship between exposure to chlorophenols and the incidences of soft tissue sarcomas and non-Hodgkin's lymphomas. However, it was not possible to exclude the possible confounding effects of tetrachlorodibenzo-*p*-dioxin (TCDD) present in the chlorophenols or other chemicals in the work environment. Case-control studies have examined the association of non-Hodgkin's lymphoma, thyroid cancer, nasopharyngeal cancer, colon cancer, and liver cancer with chlorophenol exposure (IARC 1979, 1982, 1987, 1999).

When administered in the diet, 2,4,6-trichlorophenol increased the incidences of leukemias or lymphomas in male rats and hepatocellular carcinomas and adenomas in mice of both sexes (NCI 1979, IARC 1999).

PROPERTIES

2,4,6-Trichlorophenol occurs as yellow to pinkish orange flakes or colorless crystals, with a strong phenolic odor. It is slightly soluble in water and soluble in acetone, benzene, carbon tetrachloride, diacetone alcohol, diethyl ether, denatured ethanol, methanol, pine oil, Stoddard solvent, toluene, and turpentine (IARC 1979, 1999). In the United States, 2,4,6-trichlorophenol is available in aqueous formulations that may contain dioxins and dibenzofurans as impurities (IARC 1979, NTP 2001). This chemical is stable at temperatures up to its melting-point (69°C). It reacts with sodium carbonate to form the sodium salt; the hydroxyl group reacts with metals and amines to form ethers, esters and salts. Further, the aromatic portion undergoes substitution reactions such as nitration, alkylation, acetylation, and halogenation, and it is decomposed by strong oxidizing agents (IARC 1979). 2,4,6-Trichlorophenol does not readily ignite, but contact with metals may form flammable hydrogen gas. When heated to decomposition, 2,4,6-trichlorophenol emits toxic and corrosive fumes of hydrochloric acid or other toxic gases (HSDB 2001).

USE

2,4,6-Trichlorophenol has been used primarily in various pesticide formulations and as a wood preservative. These have included fungicides, glue preservatives, insecticides, bactericides, germicides, herbicides, defoliants, antiseptics, and as an antimildew agent for textiles (IARC 1979, NTP 2001). Most uses of 2,4,6-trichlorophenol have been cancelled in the U.S.; however, it continues to be used in the synthesis of some fungicides (HSDB 2001).

PRODUCTION

Commercial production of 2,4,6-trichlorophenol in the United States was first reported in 1950. An undisclosed quantity was produced in 1974, the last year that production was reported (IARC 1979). In 1975, production of the compound was discontinued by the only U.S. manufacturer because of the high cost of removing toxic impurities of chlorinated dibenzo-*p*-dioxins. 2,4,6-Trichlorophenol is no longer produced in the U.S. (HSDB 2001); however, Chem Sources (2001) identified 12 current U.S. suppliers of 2,4,6-trichlorophenol. Reported U.S. imports totaled 2,200 lb, 600 lb, and 550 lb in 1976, 1978, and 1980, respectively (IARC 1979, HSDB 2001). The International Trade Administration (ITA 2001) listed U.S. imports and exports of pentachlorophenol and 2,4,5-trichlorophenol but no values were listed for 2,4,6-trichlorophenol in 2000.

EXPOSURE

The primary routes of potential occupational exposure to 2,4,6-trichlorophenol are inhalation and dermal contact. The general population may be exposed to 2,4,6-trichlorophenol through ingestion of contaminated food or water and inhalation of contaminated air (IARC 1999, HSDB 2001). The risk of potential occupational exposure to 2,4,6-trichlorophenol is greatest for workers involved in wood preservation or the production of chlorophenols and chemicals made from chlorophenols (ATSDR 1999, HSDB 2001). The National Occupational Exposure Survey (1981-1983) indicated that 851 total workers, including 187 women, were potentially exposed to 2,4,6-trichlorophenol in the workplace (NIOSH 1984). The National Occupational Hazard Survey, conducted by NIOSH from 1972 to 1974, estimated that 110 workers were possibly exposed to 2,4,6-trichlorophenol in the workplace. Worker exposure was primarily in hospitals and in the leather tanning and finishing industry (NIOSH 1976), as well as in the pulp and wood industry (HSDB 2001). According to NCI (1979), substantial exposure of the general population is questionable; however, residues may be present throughout the environment since it was widely used as a pesticide.

2,4,6-Trichlorophenol can form when industrial wastewater containing phenol or certain aromatic acids is treated with hypochlorite or during the disinfection of drinking water sources. Investigators have detected trichlorophenol (unspecified isomers) in river water samples, landfill leachate samples, chemical plant effluent water samples, sewage treatment plant effluent samples, tap water samples, and in ambient air (ATSDR 1990, HSDB 2001). EPA's Toxic Chemical Release Inventory (TRI) listed five industrial facilities that reported environmental releases of 2,139 lb of 2,4,6-trichlorophenol in 1999 (TRI99 2001). Environmental releases reported to the TRI declined by almost 99% from 1988 to 1990, but releases showed little change from 1990 to 1999.

REGULATIONS

EPA regulates 2,4,6-trichlorophenol under the Clean Water Act (CWA), Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), and Resource Conservation and Recovery Act (RCRA). EPA published a water quality criteria document on 2,4,6-trichlorophenol for the protection of human health. Under CWA and CERCLA, a reportable quantity (RQ) of 10 lb has been established. Under RCRA, 2,4,6-trichlorophenol is subject to reporting and record-keeping requirements as a hazardous constituent of waste. 2,4,6-Trichlorophenol is also subject to permitting and pretreatment regulations for discharges of the compound.

OSHA regulates 2,4,6-trichlorophenol under the Hazard Communication Standard and as a chemical hazard in laboratories. Regulations are summarized in Volume II, Table 181.

REFERENCES

ATSDR. Agency for Toxic Substances and Disease Registry. Toxicological Profile for 2,4,6-Trichlorophenol. (Final Report). Atlanta, GA: ATSDR, Public Health Service, U.S. Department of Health and Human Services. 1990. 119 pp. NTIS Accession No. PB91-181545.

Chem Sources. Chemical Sources International, Inc. <http://www.chemsources.com>, 2001.

HSDB. Hazardous Substances Data Bank. Online database produced by the National Library of Medicine. 2,4,6-Trichlorophenol. Profile Last updated January 31, 2001. Last review date, May 7, 1998.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Some Halogenated Hydrocarbons. Vol. 20. 609 pp. Lyon, France: IARC, 1979.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Chemicals, Industrial Processes and Industries Associated with Cancer in Humans. Supplement 4. 292 pp. Lyon, France: IARC, 1982.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Overall Evaluations of Carcinogenicity. Supplement 7. 440 pp. Lyon, France: IARC, 1987.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Re-evaluation of Some Organic Chemicals, Hydrazine and Hydrogen Peroxide. Vol. 71. 1,589 pp. Lyon, France: IARC, 1999.

ITA. International Trade Administration. U.S. Department of Commerce. Subheading 290810: Derivatives of Phenols or Phenol-Alcohols Containing only Halogen Substituents and their Salts. <http://www.ita.doc.gov/td/industry/otea/Trade-Detail/Latest-December/>, 2001.

NCI. National Cancer Institute. Carcinogenesis, Technical Report Series No. 155. Bioassay of 2,4,6-Trichlorophenol for Possible Carcinogenicity (CAS No. 88-060-2). DHEW (NIH) Publication No. 79-1711. 115 pp. National Institutes of Health, Bethesda, MD, 1979.

2,4,6-Trichlorophenol (Continued)

NIOSH. National Institute for Occupational Safety and Health. National Occupational Hazard Survey (1972-74). Cincinnati, OH: Department of Health, Education, and Welfare, 1976.

NIOSH. National Institute for Occupational Safety and Health. National Occupational Exposure Survey (1981-83). Cincinnati, OH: Department of Health and Human Services, 1984.

NTP. National Toxicology Program. NTP Chemical Repository. 2,4,6-Trichlorophenol. Last updated August 13, 2001.

TRI99. Toxic Chemicals Release Inventory 1999. Data contained in the Toxic Chemical Release Inventory (TRI). Available from the U.S. Environmental Protection Agency Office of Environmental Information, <http://www.epa.gov/triexplorer/reports.htm>, 2001.