



ENVIRONMENTAL  
HEALTH  
PERSPECTIVES

<http://www.ehponline.org>

**Associations of Fine Particulate Matter Species with  
Mortality in the United States:  
A Multicity Time-Series Analysis**

**Lingzhen Dai, Antonella Zanobetti, Petros Koutrakis,  
and Joel D. Schwartz**

<http://dx.doi.org/10.1289/ehp.1307568>

**Received: 29 August 2013**

**Accepted: 2 May 2014**

**Advance Publication: 6 May 2014**

# **Associations of Fine Particulate Matter Species with Mortality in the United States: A Multicity Time-Series Analysis**

Lingzhen Dai, Antonella Zanobetti, Petros Koutrakis, and Joel D. Schwartz

Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts, USA

**Address correspondence to** Lingzhen Dai, Department of Environmental Health, Harvard School of Public Health, Landmark Center 415, 401 Park Drive, Boston, MA 02215 USA. Telephone: 617-384-8755. Fax: 617-384-8728. E-mail: [lid535@mail.harvard.edu](mailto:lid535@mail.harvard.edu)

**Short title:** Fine particulate matter species and mortality

**Acknowledgments:** This study was supported by the U.S. Environmental Protection Agency STAR grant R-834798-01.

**Competing financial interests:** The authors have no competing financial interests.

## **Abstract**

**Background:** Epidemiological studies have examined the association between PM<sub>2.5</sub> and mortality, but there remains uncertainty about the seasonal variations in PM<sub>2.5</sub>-related effects and the relative importance of species.

**Objectives:** to estimate the effects of PM<sub>2.5</sub> species on mortality and how infiltration rates may modify the association.

**Methods:** Using city-season specific Poisson regression, we estimated PM<sub>2.5</sub> effects on approximately 4.5 million deaths for all causes, CVD, MI, stroke, and respiratory diseases in 75 U.S. cities for 2000-2006. We added interaction terms between PM<sub>2.5</sub> and monthly average species-to-PM<sub>2.5</sub> proportions of individual species to determine the relative toxicity of each species. We combined results across cities using multivariate meta-regression, and controlled for infiltration.

**Results:** We estimated a 1.18% [95% confidence interval (CI): 0.93, 1.44%] increase in all-cause mortality, a 1.03% (95% CI: 0.65, 1.41%) increase in CVD, a 1.22% (95% CI: 0.62, 1.82%) increase in MI, a 1.76% (95% CI: 1.01, 2.52%) increase in stroke, and a 1.71% (95% CI: 1.06, 2.35%) increase in respiratory deaths in association with a 10- $\mu\text{g}/\text{m}^3$  increase in 2-day averaged PM<sub>2.5</sub> concentration. The associations were largest in the spring. Silicon, calcium, and sulfur were associated with more all-cause mortality, while sulfur was related to more respiratory deaths. County-level smoking and alcohol were associated with larger estimated PM<sub>2.5</sub> effects.

**Conclusions:** Our study showed an increased risk of mortality associated with PM<sub>2.5</sub>, which varied with seasons and species. The results suggest that mass alone might not be sufficient to evaluate the health effects of particles.

## Introduction

Over the past few decades, there has been much research on the adverse effects of ambient particulate matter (PM). A number of studies have used fine PM (PM<sub>2.5</sub>, particles < 2.5 μm in aerodynamic diameter) as an exposure metric and estimated the effects of PM<sub>2.5</sub> on human health (Laden et al. 2006; Ostro et al. 2006; Pope and Dockery 2006; Zanobetti and Schwartz 2009). Meanwhile, researchers have found some PM<sub>2.5</sub> species significantly modify PM<sub>2.5</sub>-related effects (Franklin et al. 2008; Lippmann et al. 2006; Zanobetti et al. 2009). PM<sub>2.5</sub> consists of many chemical components that originate from various sources, such as traffic, biomass burning and coal combustion. The U.S. National Research Council has emphasized the importance of examining the risk of PM species (NRC 2004). Determining the differential toxicity of PM<sub>2.5</sub> species and identifying species with greatest toxicity is of great importance to emission-control strategies and regulations.

The U.S. Environmental Protection Agency (EPA) established the PM<sub>2.5</sub> Speciation Trends Network (<http://www.epa.gov/ttnamti1/speciepg.html>) in 2000. Speciation sampling was conducted every third or sixth day, which limits statistical power for analysis of responses to acute exposure and also prevent the examination of e.g. two-day moving averages of exposure which most studies find more strongly associated with mortality and hospital admissions than single day exposures. As a result, there are a limited number of studies investigating the toxicity of PM<sub>2.5</sub> components. These investigations have reported numerous components that may be responsible for particle toxicity such as elemental and organic carbon, sulfate, nitrate, and metals, including zinc, nickel, iron, potassium, and chromium (Atkinson et al. 2010; Bell et al. 2009; Franklin et al. 2008; Ostro et al. 2006; Valdes et al. 2012; Zhou et al. 2011).

Recently, Krall et al. reported on the association of 1-day average concentrations of species from the speciation network and mortality in 72 cities for the years 2000-2005 (Krall et al. 2013). This paper addresses a similar question, but with the following differences. First, Krall et al. analyzed PM components without controlling for PM mass risks. As pointed out by Mostofsky et al. (Mostofsky et al. 2012), it is possible to find associations for components because they are highly correlated with mass, and not because they are themselves particularly toxic. Second, it focuses on single day exposures. PM<sub>2.5</sub> mortality studies have consistently reported that the associations are spread over more than one day. Thus when one uses separate time series for components which are measured only 1 day in 6 or 1 day in 3, this will bias downward estimates, possibly more for some components than others. In addition, the loss of two thirds to five sixths of the data substantially reduces power.

US adults, particularly the elderly who dominate mortality statistics, spend approximately 90% of their time indoors (U.S.EPA 1989). While particles penetrate indoors, the infiltration rates vary with the extent to which windows and doors are open, which in turn can vary with local temperature and may therefore modify the association. Previous studies have reported such modification (Franklin et al. 2008; Stafoggia et al. 2008; Zanobetti et al. 2009). In this paper we address these issues and in addition examine more species, add an additional year of observation, and look at specific causes of death.

## Materials and Methods

### *Study sites*

We included 75 U.S. cities in our study (see Supplemental Material, Table S1). Cities of interest were selected based on the availability of daily mortality, PM<sub>2.5</sub> mass, and speciation data for at least 400 days between 2000 and 2006.

### *Environmental data*

We conducted county-level analysis for most cities as the city lies within a single county, and used multiple counties for a city whose population extends beyond the boundary of one county (Zanobetti and Schwartz 2009). We obtained PM<sub>2.5</sub> mass and species concentration data from the U.S. EPA Air Quality System Technology Transfer Network (<http://www.epa.gov/ttn/airs/airsaqs/>). PM<sub>2.5</sub> mass samples were collected daily in most of the cities, while the speciation monitoring sites were operated on a 1-in-3 or 1-in-6 day schedule. Most of the cities had a single monitor. For cities with more than one sampling site concentration data were averaged. Our analysis focused on organic carbon (OC), elemental carbon (EC), sodium (Na), aluminum (Al), silicon (Si), sulfur (S), potassium (K), calcium (Ca), vanadium (V), iron (Fe), nickel (Ni), copper (Cu), and zinc (Zn), because these species have been shown to be representative of several sources (e.g., motor vehicles, oil combustion, coal combustion, wood burning, sea salt, and road dust) and their concentration levels are mostly above the method detection limits (Hopke et al. 2006). Furthermore, they have been studied by previous epidemiologic and toxicological studies (Bell et al. 2009; Franklin et al. 2008; Ostro et al. 2006; Zanobetti et al. 2009; Zhou et al. 2011). Monthly average proportions between each component and PM<sub>2.5</sub> mass were calculated for each city by dividing monthly concentrations of species by the respective mass mean.

Daily mean temperature in every city was obtained from the National Oceanic and Atmospheric Administration (<http://www.noaa.gov/>). We used 24-hour average temperature data from the closest weather station to the center of the city. Percent green space data were obtained from the National Land Cover Database, Multi-Resolution Land Characteristics Consortium (<http://www.mrlc.gov/>).

### ***Health data***

Daily mortality data were obtained from National Center for Health Statistics (<http://www.cdc.gov/nchs/>). We examined non-accidental deaths due to all causes and specific diseases, which were derived from the *International Statistical Classification of Disease*, 10<sup>th</sup> Revision (WHO 2007) codes as follows: all causes (ICD-10, A00-R99), cardiovascular diseases (ICD-10, I01–I59), respiratory diseases (ICD-10, J00–J99), myocardial infarction (ICD-10, I21-I22), and stroke (ICD-10, I60-I69).

We investigated several behavioral and other risk factors that have been reported to impact health (Baja et al. 2010; Dogra et al. 2007; Dwyer-Lindgren et al. 2013; Mora et al. 2007), including diabetes, being overweight or obese (i.e., BMI  $\geq$  25), smoking, quitting smoking, alcohol consumption (having > two drinks per day), asthma, and leisure time physical activity, from the Behavioral Risk Factor Surveillance System (BRFSS) (CDC 2006). We applied county-level weighting methodology to obtain county-level percentages of these variables in 2006. For counties that were not available, we used data from the closest metropolitan or micropolitan statistical area (MMSA) and applied MMSA-level weighting methodology.

### ***Statistical methods***

We applied a two-stage analysis in our study. In the first stage, a city-specific season-stratified time-series analysis using Poisson regression in a generalized additive model (GAM) was used to estimate the association between daily mortality and the mean of PM<sub>2.5</sub> mass on the day of death and the day before death in each city and each season (defined as Spring: March – May; Summer: June – August; Fall: September – November; Winter: December – February). We controlled for time trend with a natural cubic regression spline with 1.5 degrees of freedom (d.f.) per season per year, for day of the week with indicator variables, and for daily temperature on the same day (lag 0) and on the previous day (lag 1) with a natural cubic spline with 3 d.f. for each. For every species, we calculated the monthly average species-to-PM<sub>2.5</sub> proportions for each month as a solution to the missing speciation data problem due to the 1-in-6 or 1-in-3 day sampling frequency. We then added, one at the time, the interaction terms between PM<sub>2.5</sub> and the monthly average species-to-PM<sub>2.5</sub> proportions of each individual species (Valdes et al. 2012). The model is as below:

$$\text{Log}E(Y_t) = \text{Intercept} + ns(\text{time}, df) + ns(\text{temperature}_t, df) + ns(\text{temperature}_{t-1}, df) + \text{day of the week} + \alpha Z_{t-1,t} + \beta p^i + \gamma Z_{t-1,t} p^i \quad [1]$$

where,  $E(Y_t)$  is the expected death count at day  $t$ ,  $ns$  is the natural cubic splines,  $Z_{t-1,t}$  indicates 2-day averaged concentration of PM<sub>2.5</sub> at day  $t-1$  and  $t$ , and  $p^i$  is the mean monthly proportion of species  $i$  to mass.

By using an interaction with the monthly mean ratio we avoid losing most of the daily observations, since we are able to use more than one day's exposure, and control for PM mass.

While the use of the monthly ratio introduces some error in that variable, much of the variation

in species mass is across cities, and between months within cities. For example, organic carbon, sulfate and nitrate are products of photochemical reactions whose rates are temperature-dependent, and this varies substantially across the U.S. and differently by month in different locations (Baker and Scheff 2007; de Gouw et al. 2005). It is important to note that if a species ratio is not significant in this analysis that does not mean that the species has no effect, it means its effect is not different than the average PM effect. A species with low or no toxicity would be expected to have a significant negative interaction term.

In the second stage of the analysis, we conducted a multivariate random effects meta-analysis and combined the 300 (i.e., 75 cities \* 4 seasons) city-season specific effect estimates to obtain an overall association between PM<sub>2.5</sub> mass and its interaction with each species with mortality across all 75 cities:

$$\mathbf{Y}^i = \mathbf{X}\mathbf{B}^i \quad [2]$$

where,  $\mathbf{Y}^i$  is a  $(300 \times 2)$  matrix, whose first column contains 300 city-season specific coefficients for PM<sub>2.5</sub> and the second column contains 300 city-season specific coefficients for interaction with species  $i$ ,  $\mathbf{X}$  is a  $(300 \times 4)$  matrix for intercept, linear, quadratic and cubic temperature, and  $\mathbf{B}^i$  indicates a  $(4 \times 2)$  matrix of meta-regression coefficient for PM<sub>2.5</sub> and for interaction with species  $i$ .

It has been shown that high ventilation is seen at mild temperatures whereas low ventilation is seen at high and low temperatures (Koutrakis et al. 2005). Assuming that PM effect would not drop consistently as temperature increases, we added a cubic term in the model to allow for a plateau. We also examined whether the BRFSS factors modified PM<sub>2.5</sub> effects. The model is:

$$\hat{\beta}_{is} = \beta_0 + \beta_1 t_{is} + \beta_2 t_{is}^2 + \beta_3 t_{is}^3 + \beta_4 BRFSS_i \quad [3]$$

where  $\hat{\beta}_{is}$  is the estimated PM<sub>2.5</sub> coefficient for city  $i$  in season  $s$ ,  $t_{is}$  is the centered temperature (i.e., temperature – mean temperature) for city  $i$  in season  $s$ , and  $BRFSS_i$  is the BRFSS variable in city  $i$ . To estimate the effect of individual species, we performed the same meta-regression, but with the coefficient of the interaction term for species as the outcome being modeled. Here again we adjusted for city-season mean temperature as a surrogate for air exchange. We also investigated spatial variations between cities by focusing on a single outcome and exposure season to evaluate the effects in each city by mean exposure in that season for each city.

The effect estimates for PM<sub>2.5</sub> were expressed as the percent change in mortality associated with a 10 µg/m<sup>3</sup> increase in the 2-day averaged concentration of PM<sub>2.5</sub> mass, for comparability with most previous studies. We expressed the effect of species on mortality as the estimated percent increase in mortality at the 10<sup>th</sup> and 90<sup>th</sup> percentile of distribution of species-to-PM<sub>2.5</sub> proportion for each species, holding the PM<sub>2.5</sub> increase constant at 10 µg/m<sup>3</sup>.

Data management was performed with SAS version 9.1 (SAS 2006), and regression analysis with R version 3.0.0 (R 2013).

## Results

In this study, we examined 4,473,519 all-cause deaths, of which 1,429,968 were CVD, 308,235 MI, 255,430 stroke and 436,800 respiratory deaths.

Table 1 summarizes daily mortality, PM<sub>2.5</sub>, temperature in all cities. On average, there were 28 non-accidental deaths per day. Daily death count by season was higher in the winter (n=31) and

spring (n=28). Among the causes of interest, CVD killed the most people on average (9/day), followed by respiratory diseases (3/day). The overall mean concentration of PM<sub>2.5</sub> was 13.3 µg/m<sup>3</sup>. PM<sub>2.5</sub> mean concentration was highest in the summer (15.0 µg/m<sup>3</sup>) and lowest in the spring (11.6 µg/m<sup>3</sup>). Some of the species exhibited strong seasonal variability. For example, sulfur varied from 798 ng/m<sup>3</sup> in the winter to 1669 ng/m<sup>3</sup> in the summer, with larger variations in some cities.

The distributions of monthly average proportions of PM<sub>2.5</sub> species are shown in Table 2. OC had the largest mean proportion (37.9%), followed by sulfur (8.78%) and EC (6.31%). The mean proportions for all the metals were less than 1% of mass concentration.

Table 3 presents the estimated percent increase mortality for a 10 µg/m<sup>3</sup> increase in 2-day averaged PM<sub>2.5</sub> across the 75 cities. We found statistically significant associations between PM<sub>2.5</sub> and mortality. A 1.18% (95% CI: 0.93, 1.44%) increase in all-cause mortality was associated with a 10 µg/m<sup>3</sup> increase in the 2-day averaged concentration of PM<sub>2.5</sub>. The greatest effect estimate effect was observed for stroke mortality [1.76% (95% CI: 1.01, 2.52%)], followed by respiratory deaths [1.71% (95% CI: 1.06, 2.35%)]. We observed seasonal variations in PM<sub>2.5</sub> effects (see Supplemental Material, Figure S1). For a 10-µg/m<sup>3</sup> increase in 2-day averaged PM<sub>2.5</sub>, the percent increases in all mortality categories were greatest in the spring.

Figure 1 shows the effect estimates of PM<sub>2.5</sub> on all-cause mortality in each city by mean spring PM<sub>2.5</sub> in each city. We observed differential effects across cities.

Figure 2 shows the adjusted estimated percent increases in mortality for a 10-µg/m<sup>3</sup> increase in 2-day averaged PM<sub>2.5</sub> at the 10<sup>th</sup> or 90<sup>th</sup> percentile of distribution of the proportions of species.

For all-cause mortality, interaction terms between PM<sub>2.5</sub> and species silicon, calcium, and sulfur had a *p*-value less than or equal to 0.1. We found that a 10- $\mu\text{g}/\text{m}^3$  increase in 2-day averaged PM<sub>2.5</sub> was associated with an increase in all-cause mortality of 3.55% (95% CI: 1.35, 5.81%) at 90<sup>th</sup> percentile of distribution of the sulfur-to-PM<sub>2.5</sub> proportion versus 2.16% (95% CI: 1.27, 3.06%) at the 10<sup>th</sup> percentile of the sulfur-to-PM<sub>2.5</sub> ratio (see Supplemental Material, Table S2; same below). We also found silicon [3.25% (95% CI: 1.91, 4.62%) vs. 1.87% (95% CI: 1.42, 2.32%)] and calcium [3.42% (95% CI: 2.08, 4.77%) vs. 1.75% (95% CI: 1.34, 2.16%)] were associated with higher estimated effects of PM<sub>2.5</sub> on all-cause mortality. In addition, sulfur was associated with higher estimated PM<sub>2.5</sub> effect on respiratory deaths. The percent increase in respiratory mortality at the 90<sup>th</sup> percentile of the sulfur-to-PM<sub>2.5</sub> proportion was 8.96% (95% CI: 1.55, 16.90%), vs. 4.44% (95% CI: 1.46, 7.51%) at the 10<sup>th</sup> percentile.

Figure 3 indicates the relationship between effect estimates and city-season temperature, which serves as a surrogate for ventilation and thus particle penetration indoors. We observed an inverted U-shape relationship with a plateau at high temperatures. The *p*-value for cubic term is 0.06 in meta-regression without BRFSS factors and is 0.07 controlled for smoking and alcohol consumption. The all-cause mortality effect estimates first increase as temperature increases and peak around a seasonal average of 10 °C. After that, they decrease until they reach a plateau at around 28 °C.

County-level percent of green space, diabetes, obesity, asthma, or physical activity did not modify the effect of PM<sub>2.5</sub> on mortality. However, among the behavioral factors, we found the effects of PM<sub>2.5</sub> were higher in areas where people smoked more or had two drinks or more per day. Specifically, an IQR increase in the prevalence of smokers (8.8%) was associated with a

34% increase in estimated  $PM_{2.5}$  effects, while an IQR increase in the prevalence of heavy drinkers (7.0%) was associated with an increase of 40% in the estimated effects of fine particles.

## **Discussion**

In this nationwide time-series study, we estimated the effects of  $PM_{2.5}$  mass and species on daily mortality across 75 U.S. cities, covering over 4 million deaths. We found an increase in  $PM_{2.5}$  concentration at lag day 0-1 was statistically significantly associated with increased risk of all-cause mortality, CVD, MI, stroke, and respiratory mortality. We also found that  $PM_{2.5}$ -related effects were modified by certain species. Furthermore, analysis by season indicated that effect estimates were highest in the spring. To investigate this seasonal pattern we included city-season specific temperature in the meta-regression analysis. These seasonal variations may affect the characteristics of  $PM_{2.5}$  mixture and mediate its effects on health outcomes (Bell et al. 2007).

Controlling for this potential confounder and for  $PM_{2.5}$  mass, we found that a species related to coal combustion (i.e., sulfur) was associated with higher risks for all cause but particularly respiratory mortality. Sulfur is also a marker of regional pollution thus it may not only reflect exposures to power plant emissions. Changes in the proportion of OC mass in  $PM_{2.5}$  did not modify its effect on mortality for any cause, suggesting this species has average toxicity. We found higher silicon or calcium proportions were associated with increased estimated  $PM_{2.5}$  mortality risks. These crustal elements are often elevated near roads and can be a surrogate for increased road dust, which in addition to those elements contains various organic compounds, compounds from tire and brake wear, etc. (Rogge et al. 1993). Thus they may be a marker for pollution from traffic other than EC.

The BRFSS factors we examined were on the county level. The distributions of prevalence of smoking and heavy drinkers (i.e., > two drinks/day) in different cities were approximately normal distributed with a mean around 30% and 60%, respectively (see Supplemental Material, Figure S2). We found that cities with more smokers or heavy drinkers had larger estimated effects of PM<sub>2.5</sub>. These have not previously been identified as susceptibility factors for the effects of particles on health, and this requires greater attention.

The magnitudes of effects in our study are comparable to those reported by other studies. For example, a study that included 112 U.S. cities reported a 0.98% (95% CI: 0.75, 1.22%) increase, a 0.85% (95% CI: 0.46, 1.24%) increase, a 1.18% (95% CI: 0.48, 1.89%) increase, a 1.78% (95% CI: 0.96, 2.62%) increase, and a 1.68% (95% CI: 1.04, 2.33%) increase in all-cause, CVD, MI, stroke, and respiratory mortality, respectively, for a 10- $\mu\text{g}/\text{m}^3$  increase in 2-day averaged PM<sub>2.5</sub> (Zanobetti and Schwartz 2009). Our estimates are slightly higher than the above ones and are closer to those by a 27-city study, which found a 1.21% (95% CI: 0.29, 2.14%) increase in all-cause mortality, a 1.78% (95% CI: 0.20, 3.36%) increase in respiratory mortality and 1.03% (95% CI: 0.02, 2.04%) increase in stroke mortality for a 10  $\mu\text{g}/\text{m}^3$  increase in previous day's PM<sub>2.5</sub> (Franklin et al. 2007). Our study and these two studies used city-season specific models to allow for seasonal differences in the effects of temperature and day of the week.

The finding that effects were highest in the spring is consistent with previous studies (Zanobetti and Schwartz 2009; Zeka et al. 2006). Franklin et al. (2008) found similar pattern using linear and quadratic temperature in the meta-regression. Additionally, we included the cubic term, which was marginally significant and led to the small plateau at high temperatures. These results

indicated greater effects for moderate temperatures when windows are more likely to be open and particle penetration rates are higher.

EC is considered as a marker of traffic emissions (Viana et al. 2006). Previous research has reported EC was significantly associated with increased risk of mortality due to all causes or cardiovascular diseases (Bell et al. 2009; Metzger et al. 2004; Peng et al. 2009). In this study, we did observe that increase in the EC-to-PM<sub>2.5</sub> proportion increased the association between PM<sub>2.5</sub> and all-cause mortality and CVD mortality in crude meta-regression, but it was no longer significant when we controlled for city-season temperature. Similarly, there were two studies that also controlled for temperature in the meta-regression and did not find any effect modification by EC in the association between PM<sub>2.5</sub> and non-accidental mortality or hospital admissions for cardiovascular diseases (Franklin et al. 2008; Zanobetti et al. 2009).

Silicon and calcium, which may be associated with soil or road dust, were observed to modify the effects of PM<sub>2.5</sub> on all-cause mortality in our study. Crustal elements have been reported to have adverse effects on health. For example, Ostro et al. found strong association between silicon and mortality (Ostro et al. 2010); Franklin et al. (2008) observed silicon and aluminum were modifiers of the PM<sub>2.5</sub>-mortality effects. There were also studies that showed plausible biological mechanisms of inflammatory effects of road dust containing aluminum and/or silicon (Becker et al. 2005; Clarke et al. 2000). Additionally, road dust is often coated with organic compounds and metals from car exhaust, tire wear, etc. (Rogge et al. 1993), that may contribute to its toxicity.

Nickel, as a marker of oil combustion, was reported to have effect modification in the relationship between PM<sub>2.5</sub> and mortality or hospital admissions in previous studies (Franklin et al. 2008; Zanobetti et al. 2009), but we did not observe any. On average, nickel only accounted for 0.02% of the PM<sub>2.5</sub> concentration in this study. The concentrations of nickel are frequently lower than the method detection limit (Burnett et al. 2000), which may make us fail to detect its effects. Nevertheless, toxicological research has found evidence on its adverse effects (Gao et al. 2004; Lippmann et al. 2006). For example, Lippmann et al. (2006) found atherosclerotic prone mice that were exposed to concentrated air particles had a pronounced acute change in heart rate and heart rate variability when nickel was especially high. Lippmann and the New York studies which found nickel effects had high exposures due to the residual fuel burn in New York for heating. The levels for the entire country are lower.

We observed the effect of PM<sub>2.5</sub> mass on all-cause and respiratory mortality was modified by sulfur. This component is a marker of coal combustion emissions, which suggests species derived from coal combustion might have great toxicity on mortality, particularly due to respiratory diseases. Sulfate is the primary form of sulfur in particles. Sulfate has been implicated as a major toxic species in PM<sub>2.5</sub> (Amdur 1996) and reported to be associated with increased risk of various mortality outcomes in earlier epidemiological studies (Fairley 1999; Hoek et al. 2000; Laden et al. 2000; Mar et al. 2000). The importance of sulfates in the air may be due to the ability of acid sulfates to solubilize transition metals and thus making them bioavailable (Ghio et al. 1999). There were studies that found sulfate was associated with endothelial dysfunction (O' Neill et al. 2005), increased oxidative stress and coagulation (Chuang et al. 2007). These toxicology findings provide plausibility to sulfate health effects.

One disappointing aspect of this result is that despite the use of 75 cities and almost 4.5 million deaths, we were unable to distinguish much difference in toxicity for many of the species we examined. This may reflect only modest differences in toxicity, but may also reflect more fundamental difficulties in identifying differences between many correlated exposures with limited measurements over time. Evidence of the low power to detect differences can be seen in the difference between our results for all deaths and results for cardiovascular deaths. The pattern of higher estimated effects when PM mass has a larger fraction of silicon, sulfur, and calcium is present for cardiovascular deaths as well, but with a third as many deaths, it does not reach significance. One option to improve study power might be specifically selecting locations with high proportions of the species of interest.

There are several limitations in this study. First, our ability to capture spatial variability is constrained due to the location of U.S. EPA monitors. A previous study showed moderate to low monitor-to-monitor correlations between daily concentrations of several species (arsenic, EC, and nickel) in the New York City area, which suggest high spatial variability in some speciation concentrations (Ito et al. 2004). Differential measurement error between species that are better or worse represented by a single monitor may bias differential results. However, in a time-series study much of the geographic variability will result in Berkson error (Zeger et al. 2000), which will not produce bias. Meanwhile, failing to capture spatial variability might weaken study power and attenuate estimates. Second, we failed to capture day-to-day variation in the analysis. Although we used monthly average species-to-PM<sub>2.5</sub> proportions to gain more power, we still lost variation across days. Nevertheless, we believe that the day-to-day variation is random error in measurement, which induces little or no bias.

Third, as mentioned above, there are data limitations, such as the one-in-six and one-in-three sampling frequency for the species. Whether one takes the approach of Krall et al. (2013) and only analyzes those days, or our approach and gains power by analyzing every day but with more error prone monthly means of the species, there is a price that is paid for this lack of data. Together with moderate correlation among the species and with total particle mass, this makes the task difficult. We do not believe this is likely to produce false positives, and hence we believe our findings are well supported.

## References

- Amdur M. 1996. Animal toxicology. En: Wilson R, Spengler J, ed. Particles in our air. Concentrations and health effects. Boston:Harvard University Press, 84-121.
- Atkinson RW, Fuller GW, Anderson HR, Harrison RM, Armstrong B. 2010. Urban Ambient Particle Metrics and Health: A Time-series Analysis. *Epidemiology* 21(4):501-511.
- Baja ES, Schwartz JD, Wellenius GA, Coull BA, Zanobetti A, Vokonas PS, et al. 2010. Traffic-related air pollution and QT interval: modification by diabetes, obesity, and oxidative stress gene polymorphisms in the normative aging study. *Environ Health Perspect* 118(6):840-846.
- Baker K, Scheff P. 2007. Photochemical model performance for PM<sub>2.5</sub> sulfate, nitrate, ammonium, and precursor species SO<sub>2</sub>, HNO<sub>3</sub>, and NH<sub>3</sub> at background monitor locations in the central and eastern United States. *Atmos Environ* 41(29):6185-6195.
- Becker S, Dailey L, Soukup J, Grambow S, Devlin R, Huang Y. 2005. Seasonal variations in air pollution particle-induced inflammatory mediator release and oxidative stress. *Environ Health Perspect* 113:1032-1038.
- Bell M, Dominici F, Ebisu K, Zeger S, Samet J. 2007. Spatial and temporal variation in PM<sub>2.5</sub> chemical composition in the United States for health effects studies. *Environ Health Perspect* 115:989-995.
- Bell M, Ebisu K, Peng R, Samet J, Dominici F. 2009. Hospital Admissions and Chemical Composition of Fine Particle Air Pollution. *Am J Respir Crit Care Med* 179(12):1115-1120.
- Burnett R, Brook J, Dann T, Delocla C, Philips O, Cakmak S, et al. 2000. Association between particulate- and gas-phase components of urban air pollution and daily mortality in eight Canadian cities. *Inhal Toxicol* 12(Suppl 4):15-39.
- CDC. 2006. Behavioral Risk Factor Surveillance System (BRFSS). Available: <http://www.cdc.gov/brfss/> [accessed July 12 2013].
- Chuang KJ, Chan CC, Su TC, Lee CT, Tang CS. 2007. The effect of urban air pollution on inflammation, oxidative stress, coagulation, and autonomic dysfunction in young adults. *Am J Respir Crit Care Med* 176(4):370-376.
- Clarke R, Coull B, Reinisch U, Catalano P, Killingsworth C, Koutrakis P, et al. 2000. Inhaled concentrated ambient particles are associated with hematologic and bronchoalveolar lavage changes in canines. *Environ Health Perspect* 108:1179-1187.

- de Gouw JA, Middlebrook AM, Warneke C, Goldan PD, Kuster WC, Roberts JM, et al. 2005. Budget of organic carbon in a polluted atmosphere: Results from the New England Air Quality Study in 2002. *J Geophys Res*:110(D16):D16305.
- Dogra S, Arden CI, Baker J. 2007. The relationship between age of asthma onset and cardiovascular disease in Canadians. *J Asthma* 44(10):849-854.
- Dwyer-Lindgren L, Freedman G, Engell RE, Fleming TD, Lim SS, Murray CJ, et al. 2013. Prevalence of physical activity and obesity in US counties, 2001--2011: a road map for action. *Popul Health Metr* 11(1):7.
- Fairley D. 1999. Daily mortality and air pollution in Santa Clara County, California: 1989-1996. *Environ Health Perspect* 107(8):637-641.
- Franklin M, Koutrakis P, Schwartz J. 2008. The Role of Particle Composition on the Association Between PM<sub>2.5</sub> and Mortality. *Epidemiology* 19(5):680-689.
- Franklin M, Zeka A, Schwartz J. 2007. Association between PM<sub>2.5</sub> and all-cause and specific-cause mortality in 27 US communities. *J Expo Sci Environ Epidemiol* 17:279-287.
- Gao F, Barchowsky A, Nemec AA, Fabisiak JP. 2004. Microbial stimulation by *Mycoplasma fermentans* synergistically amplifies IL-6 release by human lung fibroblasts in response to residual oil fly ash (ROFA) and nickel. *Toxicol Sci* 81(2):467-479.
- Ghio AJ, Stoneheurner J, McGee JK, Kinsey JS. 1999. Sulfate content correlates with iron concentrations in ambient air pollution particles. *Inhal Toxicol* 11(4):293-307.
- Hoek G, Brunekreef B, Verhoeff A, van Wijnen J, Fischer P. 2000. Daily mortality and air pollution in The Netherlands. *J Air Waste Manag Assoc* 50(8):1380-1389.
- Hopke PK, Ito K, Mar T, Christensen WF, Eatough DJ, Henry RC, et al. 2006. PM source apportionment and health effects: 1. Intercomparison of source apportionment results. *J Expo Sci Environ Epidemiol* 16(3):275-286.
- Ito K, Xue N, Thurston G. 2004. Spatial variation of PM<sub>2.5</sub> chemical species and source-apportioned mass concentrations in New York City. *Atmos Environ* 38:5269-5282.
- Koutrakis P, Suh HH, Sarnat JA, Brown KW, Coull BA, Schwartz J. 2005. Characterization of particulate and gas exposures of sensitive subpopulations living in Baltimore and Boston. *Res Rep Health Eff Inst* 131:1-65.

- Krall JR, Anderson GB, Dominici F, Bell ML, Peng RD. 2013. Short-term Exposure to Particulate Matter Constituents and Mortality in a National Study of U.S. Urban Communities. *Environ Health Perspect* 2:2.
- Laden F, Neas LM, Dockery DW, Schwartz J. 2000. Association of fine particulate matter from different sources with daily mortality in six US cities. *Environ Health Perspect* 108(10):941-947.
- Laden F, Schwartz J, Speizer FE, Dockery DW. 2006. Reduction in fine particulate air pollution and mortality - Extended follow-up of the Harvard six cities study. *Am J Respir Crit Care Med* 173(6):667-672.
- Lippmann M, Ito K, Hwang J, Maciejczyk P, Chen L. 2006. Cardiovascular effects of nickel in ambient air. *Environ Health Perspect* 114:1662-1669.
- Mar TF, Norris GA, Koenig JQ, Larson TV. 2000. Associations between air pollution and mortality in Phoenix, 1995-1997. *Environ Health Perspect* 108(4):347-353.
- Metzger K, Tolbert P, Klein M, Peel J, Flanders W, Todd K, et al. 2004. Ambient air pollution and cardiovascular emergency department visits. *Epidemiology* 15:46-56.
- Mora S, Cook N, Buring JE, Ridker PM, Lee IM. 2007. Physical activity and reduced risk of cardiovascular events: potential mediating mechanisms. *Circulation* 116(19):2110-2118.
- Mostofsky E, Schwartz J, Coull BA, Koutrakis P, Wellenius GA, Suh HH, et al. 2012. Modeling the association between particle constituents of air pollution and health outcomes. *Am J Epidemiol* 176(4):317-326.
- NRC. 2004. Research Priorities for Airborne Particulate Matter: IV. Continuing Research Progress Washington, DC: National Research Council. Available: [http://epa.gov/ncer/science/pm/2008sab/exec\\_summary\\_2004\\_nrc\\_report.pdf](http://epa.gov/ncer/science/pm/2008sab/exec_summary_2004_nrc_report.pdf)
- O' Neill M, Veves A, Zanobetti A, Sarnat J, Gold D, Economides P, et al. 2005. Diabetes enhances vulnerability to particulate air pollution-associated impairment in vascular reactivity and endothelial function. *Circulation* 111:2913-2920.
- Ostro B, Broadwin R, Green S, Feng W, Lipsett M. 2006. Fine particulate air pollution and mortality in nine California counties: results from CALFINE. *Environ Health Perspect* 114:29-33.

- Ostro B, Lipsett M, Reynolds P, Goldberg D, Hertz A, Garcia C, et al. 2010. Long-Term Exposure to Constituents of Fine Particulate Air Pollution and Mortality: Results from the California Teachers Study. *Environ Health Perspect* 118(3):363-369.
- Peng R, Bell M, Geyh A, McDermott A, Zeger S, Samet J, et al. 2009. Emergency admissions for cardiovascular and respiratory diseases and the chemical composition of fine particle air pollution. *Environ Health Perspect* 117:957-963.
- Pope CA, 3rd, Dockery DW. 2006. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc* 56(6):709-742.
- R. 2013. R: A Language and Environment for Statistical Computing. 3.0.0 edition.
- Rogge WF, Hildemann LM, Mazurek MA, Cass GR, Simoneit BRT. 1993. Sources of fine organic aerosol. 2. Noncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks. *Environ Sci Technol* 27(4):636-651.
- SAS. 2006. SAS Software Release 9.1.
- Stafoggia M, Schwartz J, Forastiere F, Perucci CA, Group S. 2008. Does temperature modify the association between air pollution and mortality? A multicity case-crossover analysis in Italy. *Am J Epidemiol* 167(12):1476-1485.
- U.S.EPA. 1989. Report to Congress on indoor air quality: Volume 2. EPA/400/1-89/001C. Washington, DC.
- Valdes A, Zanobetti A, Halonen J, Cifuentes L, Morata D, Schwartz J. 2012. Elemental concentrations of ambient particles and cause specific mortality in Santiago, Chile: a time series study. *Environ Health* 11(1):82.
- Viana M, Querol X, Alastuey A, Gil JI, Menéndez M. 2006. Identification of PM sources by principal component analysis (PCA) coupled with wind direction data. *Chemosphere* 65(11):2411-2418.
- WHO, 2007. International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Version for 2007. Available: <http://apps.who.int/classifications/apps/icd/icd10online2007/>
- Zanobetti A, Franklin M, Koutrakis P, Schwartz J. 2009. Fine particulate air pollution and its components in association with cause-specific emergency admissions. *Environ Health* 8(1):58.

- Zanobetti A, Schwartz J. 2009. The Effect of Fine and Coarse Particulate Air Pollution on Mortality: A National Analysis. *Environ Health Perspect* 117(6):898-903.
- Zeger S, Thomas D, Dominici F, Samet J, Schwartz J, Dockery D, et al. 2000. Exposure measurement error in time-series studies of air pollution: concepts and consequences. *Environ Health Perspect* 108:419-426.
- Zeka A, Zanobetti A, Schwartz J. 2006. Individual-level modifiers of the effects of particulate matter on daily mortality. *Am J Epidemiol* 163(9):849-859.
- Zhou J, Ito K, Lall R, Lippmann M, Thurston G. 2011. Time-series analysis of mortality effects of fine particulate matter components in Detroit and Seattle. *Environ Health Perspect* 119(4):461-466.

**Table 1.** Summary of daily mortality counts, PM<sub>2.5</sub>, and temperature across all 75 cities in 2000–2006 (mean ± SD).

<b>Variable</b>	<b>Overall</b>	<b>Spring</b>	<b>Summer</b>	<b>Fall</b>	<b>Winter</b>
Mortality (no.)					
All causes	28.0 ± 33.9	28.2 ± 33.9	26.0 ± 31.4	27.1 ± 32.5	30.8 ± 37.2
CVD	9.0 ± 12.6	9.1 ± 12.6	8.2 ± 11.5	8.5 ± 11.8	10.0 ± 14.1
MI	1.9 ± 2.9	1.9 ± 2.9	1.8 ± 2.6	1.8 ± 2.7	2.2 ± 3.3
Stroke	1.6 ± 2.2	1.6 ± 2.2	1.5 ± 2.0	1.6 ± 2.1	1.8 ± 2.4
Respiratory diseases	2.7 ± 3.6	2.9 ± 3.7	2.3 ± 3.1	2.4 ± 3.2	3.3 ± 4.4
Temperature (°C)	14.1 ± 10.0	13.4 ± 7.5	24.0 ± 4.1	15.1 ± 7.3	3.6 ± 8.0
PM <sub>2.5</sub> (µg/m <sup>3</sup> )	13.3 ± 8.3	11.6 ± 6.5	15.0 ± 8.8	12.8 ± 8.4	13.9 ± 9.0
PM <sub>2.5</sub> species <sup>a</sup> (ng/m <sup>3</sup> )					
OC	4367 ± 2752	3688 ± 1806	4590 ± 2371	4491 ± 2716	4688 ± 3724
EC	724 ± 590	602 ± 438	628 ± 459	830 ± 647	842 ± 733
Na	80 ± 141	93 ± 165	89 ± 152	66 ± 117	71 ± 122
Al	31 ± 78	31 ± 55	51 ± 128	23 ± 45	15 ± 34
Si	117 ± 177	123 ± 134	171 ± 273	98 ± 125	69 ± 85
S	1174 ± 1019	1066 ± 731	1669 ± 1385	1107 ± 960	798 ± 512
K	79 ± 197	63 ± 49	103 ± 360	69 ± 62	79 ± 103
Ca	65 ± 77	65 ± 68	74 ± 77	68 ± 88	53 ± 72
V	2.5 ± 4.0	2.2 ± 3.4	2.7 ± 4.2	2.7 ± 4.4	2.5 ± 3.8
Fe	102 ± 124	93 ± 127	111 ± 111	108 ± 136	93 ± 121
Ni	2.5 ± 11.6	2.3 ± 6.0	2.2 ± 6.5	2.2 ± 5.7	3.2 ± 21.4
Cu	5.1 ± 8.9	4.2 ± 7.3	5.7 ± 11.5	5.0 ± 7.3	5.4 ± 8.6
Zn	18 ± 57	16 ± 39	16 ± 57	19 ± 53	22 ± 76

<sup>a</sup>Method detection limits for species can be found at <https://aqs.epa.gov/aqsweb/codes/data/Parameters-SPECIATION.csv>

**Table 2.** Distributions of monthly species-to-PM<sub>2.5</sub> proportions (%) across all 75 cities.<sup>a</sup>

<b>Species</b>	<b>Mean ± SD</b>	<b>10<sup>th</sup></b>	<b>25<sup>th</sup></b>	<b>50<sup>th</sup></b>	<b>75<sup>th</sup></b>	<b>90<sup>th</sup></b>
OC	37.9 ± 16.9	24.6	29.1	35.5	44.1	53.6
EC	6.31 ± 3.45	2.86	3.96	5.51	7.49	10.1
Na	0.82 ± 1.31	0.07	0.20	0.45	0.96	1.90
Al	0.28 ± 0.44	0.03	0.07	0.15	0.31	0.64
Si	1.07 ± 1.22	0.30	0.45	0.70	1.20	2.12
S	8.78 ± 3.80	4.54	6.83	8.96	11.1	12.7
K	0.64 ± 0.52	0.31	0.40	0.53	0.72	0.98
Ca	0.62 ± 0.67	0.17	0.26	0.44	0.73	1.24
V	0.02 ± 0.03	0.00	0.01	0.01	0.03	0.05
Fe	0.89 ± 0.72	0.33	0.47	0.70	1.07	1.57
Ni	0.02 ± 0.06	0.00	0.00	0.01	0.02	0.04
Cu	0.04 ± 0.05	0.01	0.02	0.03	0.05	0.08
Zn	0.15 ± 0.18	0.04	0.06	0.10	0.15	0.24

<sup>a</sup>The 3-7<sup>th</sup> columns indicate 10<sup>th</sup>, 25<sup>th</sup>, ..., 90<sup>th</sup> percentiles of the monthly species-to-PM<sub>2.5</sub> proportions.

**Table 3.** Estimated percent difference in mortality (95% CI) in association with a 10- $\mu\text{g}/\text{m}^3$  increase in  $\text{PM}_{2.5}$  at lag 0-1 by cause of death and season.

<b>Mortality</b>	<b>Overall</b>	<b>Spring</b>	<b>Summer</b>	<b>Fall</b>	<b>Winter</b>
All causes	1.18 (0.93, 1.44)	2.85 (2.23, 3.47)	0.85 (0.42, 1.28)	1.17 (0.72, 1.63)	0.46 (0.07, 0.85)
CVD	1.03 (0.65, 1.41)	2.47 (1.52, 3.43)	1.03 (0.38, 1.67)	0.87 (0.33, 1.42)	0.39 (-0.36, 1.14)
MI	1.22 (0.62, 1.82)	2.08 (0.72, 3.46)	1.23 (-0.19, 2.66)	0.81 (-0.32, 1.95)	0.41 (-1.12, 1.96)
Stroke	1.76 (1.01, 2.52)	3.31 (0.49, 6.22)	1.16 (-0.42, 2.76)	1.31 (0.05, 2.58)	1.59 (0.16, 3.03)
Respiratory diseases	1.71 (1.06, 2.35)	4.03 (2.85, 5.21)	1.09 (-0.58, 2.78)	0.58 (-0.39, 1.57)	0.86 (-0.11, 1.84)

## Figure legends

**Figure 1.** Spatial variations in estimated PM<sub>2.5</sub> effects between cities.

**Figure 2.** Estimated percent difference in mortality for a 10- $\mu\text{g}/\text{m}^3$  increase in PM<sub>2.5</sub> at lag 0-1 and an increase of 10<sup>th</sup> or 90<sup>th</sup> percentile of distribution of monthly species-to-PM<sub>2.5</sub> proportions, controlled for city-season specific temperature (\* indicates a  $p$ -value  $\leq 0.1$  for the interaction term, \*\* indicates a  $p$ -value  $\leq 0.05$  for the interaction term).

**Figure 3.** Relationship between estimated effects of PM<sub>2.5</sub> on all-cause mortality and temperature (controlled for smoking and alcohol consumption).

Figure 1.




Figure 2.




Figure 3.

