

Urinary Perchlorate and Thyroid Hormone Levels in Adolescent and Adult Men and Women Living in the United States

Benjamin C. Blount, James L. Pirkle, John D. Osterloh, Liza Valentin-Blasini, and Kathleen L. Caldwell

Division of Laboratory Sciences, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

BACKGROUND: Perchlorate is commonly found in the environment and known to inhibit thyroid function at high doses. Assessing the potential effect of low-level exposure to perchlorate on thyroid function is an area of ongoing research.

OBJECTIVES: We evaluated the potential relationship between urinary levels of perchlorate and serum levels of thyroid stimulating hormone (TSH) and total thyroxine (T_4) in 2,299 men and women, ≥ 12 years of age, participating in the National Health and Nutrition Examination Survey (NHANES) during 2001–2002.

METHODS: We used multiple regression models of T_4 and TSH that included perchlorate and covariates known to be or likely to be associated with T_4 or TSH levels: age, race/ethnicity, body mass index, estrogen use, menopausal status, pregnancy status, premenarche status, serum C-reactive protein, serum albumin, serum cotinine, hours of fasting, urinary thiocyanate, urinary nitrate, and selected medication groups.

RESULTS: Perchlorate was not a significant predictor of T_4 or TSH levels in men. For women overall, perchlorate was a significant predictor of both T_4 and TSH. For women with urinary iodine $< 100 \mu\text{g/L}$, perchlorate was a significant negative predictor of T_4 ($p < 0.0001$) and a positive predictor of TSH ($p = 0.001$). For women with urinary iodine $\geq 100 \mu\text{g/L}$, perchlorate was a significant positive predictor of TSH ($p = 0.025$) but not T_4 ($p = 0.550$).

CONCLUSIONS: These associations of perchlorate with T_4 and TSH are coherent in direction and independent of other variables known to affect thyroid function, but are present at perchlorate exposure levels that were unanticipated based on previous studies.

KEY WORDS: exposure, iodine, NHANES, perchlorate, thyroid, thyroxine, TSH. *Environ Health Perspect* 114:1865–1871 (2006). doi:10.1289/ehp.9466 available via <http://dx.doi.org/> [Online 5 October 2006]

Perchlorate is an inorganic anion used for a variety of products such as road flares, explosives, pyrotechnics, and solid rocket propellant (Mendiratta et al. 1996). Perchlorate can also form naturally in the atmosphere, leading to trace levels of perchlorate in precipitation (Dasgupta et al. 2005). Natural processes are considered to concentrate perchlorate in some locations such as regions of west Texas (Dasgupta et al. 2005) and northern Chile (Urbansky et al. 2001). A combination of human activities and natural sources has led to the widespread presence of perchlorate in the environment. As of November 2005, perchlorate was detected in drinking water samples from 4.1% of community water supplies in 26 different states, with levels ranging from the method detection limit of $4 \mu\text{g/L}$ to a maximum at $420 \mu\text{g/L}$ [U.S. Environmental Protection Agency (EPA) 2005]. Most of this drinking-water contamination is likely due to contaminated source waters, although in rare instances perchlorate formation has been reported to occur in water distribution systems (Jackson et al. 2004). Additionally, perchlorate exposure from the diet is probable because of the contamination of milk (Kirk et al. 2005), vegetables (Sanchez et al. 2005), fruit (Sanchez et al. 2006a), grain (Sanchez et al. 2006b), and forage crops (Jackson et al. 2005). Perchlorate contamination has also

been reported in dietary supplements and flavor enhancers (Snyder et al. 2006).

Trace levels of perchlorate in the environment leads to human exposure. Direct measurement of perchlorate in biological samples collected from people [National Research Council (NRC) 2005] is considered an excellent assessment of their exposure. We recently assessed perchlorate exposure in a nationally representative sample of 2,820 U.S. residents, ≥ 6 years of age, who participated in the National Health and Nutrition Examination Survey (NHANES) during 2001 and 2002 (Blount et al. 2006).

Environmental perchlorate exposure is of potential health concern because much larger doses of perchlorate have been shown to competitively inhibit iodide uptake (Greer et al. 2002; Wýngaarden et al. 1953). Populations with low intake of iodine or increased demand for iodine may be more vulnerable to inhibition of iodide uptake. Sustained inhibition of iodide uptake can lead to hypothyroidism, although perchlorate-induced changes to thyroid function have not been previously demonstrated in any human population exposed to perchlorate, even at doses as high as 0.5 mg/kg body weight per day (NRC 2005). The thyroid plays a crucial role in energy homeostasis and neurologic development. Hypothyroidism can lead to metabolic

problems in adults and abnormal development during gestation and infancy (Braverman and Utiger 2000). Severe hypothyroidism due to iodine deficiency during pregnancy is a preventable cause of cretinism, a permanent cognitive impairment of the developing fetus (Glinooer 2000). Mild hypothyroidism during pregnancy has been associated with subtle cognitive deficits in children (Haddow et al. 1999; Klein et al. 2001), leading the NRC to recommend that consideration be given to adding iodide to all prenatal vitamins (NRC 2005). Therefore, we examined relationships between urinary perchlorate and serum thyroid hormones in men and women, ≥ 12 years of age, who participated in NHANES 2001–2002.

Subjects and Methods

Study design. NHANES is conducted by the National Center for Health Statistics of the Centers for Disease Control and Prevention (CDC). This survey is designed to assess the health and nutrition status of the civilian, non-institutionalized U.S. population. NHANES uses a complex multistage probability sampling designed to be representative of the U.S. population based on age, sex, race/ethnicity, and income. Data reported in the present study were collected using an extensive household interview addressing health conditions and health-related behaviors and a standardized physical examination including medical blood and urine tests, which were conducted in mobile examination centers. NHANES 2001–2002 was conducted in 30 locations throughout the United States. Overall, the

Address correspondence to B.C. Blount, Division of Laboratory Sciences, National Center for Environmental Health, CDC, 4770 Buford Highway, NE, Mail Stop F47, Atlanta, GA 30341 USA. Telephone: (770) 488-7894. Fax: (770) 488-0181. E-mail: bkb3@cdc.gov

We thank the staff at the National Center for Health Statistics and Westat who were responsible for planning and conducting the National Health and Nutrition Examination Survey (NHANES), and E. Gunter and C. Pfeiffer for managing the National Center for Environmental Health's involvement with NHANES. We thank J. Morrow, J. Mauldin, S. Caudill, A. Delinsky, J. Phillips, and M. Smith for technical assistance.

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention.

The authors declare they have no competing financial interests.

Received 27 June 2006; accepted 4 October 2006.

survey interview response rate was 83.9% and the exam response rate was 79.6%. A full description of the NHANES survey is available on the NHANES website (CDC 2004). The study protocol was reviewed and approved by the CDC institutional review board; additionally, informed written consent was obtained from all subjects before they took part in the study.

Urinary perchlorate levels were measured by the Division of Laboratory Sciences, National Center for Environmental Health, on a representative random one-third subsample consisting of 2,820 study participants (males and females), ≥ 6 years of age (Blount et al. 2006). For ages ≥ 12 years, 2,517 persons were in the random subsample. Serum levels of thyroid stimulating hormone (TSH) and total thyroxine (T_4) were only available for 2,299 participants ≥ 12 years of age.

Demographic variables. Sociodemographic data were self-reported by study participants. Race/ethnicity was derived from self-reported questionnaire data and categorized as non-Hispanic white, non-Hispanic black, Mexican American, and "other." Each of these race/ethnicity categories was used in the regression modeling. Non-Hispanic whites were used as the referent group in regression analysis.

Laboratory methods. During the physical examinations, whole blood and spot urine specimens were collected from participants,

aliquoted, and stored cold (2–4°C) or frozen until shipment. Whole blood was collected into a red-top 15-mL Vacutainer tube, mixed, allowed to clot 30–45 min, and centrifuged; approximately 1 mL serum was stored frozen in a cryovial for future analysis for TSH and T_4 . Serum samples collected in 2001 were assayed for TSH and T_4 by the Coulston Foundation (Alamogordo, NM) using a microparticle enzyme immunoassay for the quantitative determination of TSH, and a Hitachi 704 chemistry analyzer (Hitachi Chemical Diagnostics, Mountain View, CA) for the quantitative determination of T_4 (CDC 2003). Serum samples collected in 2002 were assayed for TSH and T_4 by Collaborative Laboratory Services (Ottumwa, IA) using a chemiluminescent immunoassay (Access Immunoassay System; Beckman Instruments, Fullerton, CA) (CDC 2003). The National Center for Health Statistics evaluated the TSH and T_4 data sets from the two laboratories and determined that the values are comparable across the 2 years.

Surplus urine samples from NHANES 2001–2002 were shipped on dry ice to the Division of Laboratory Sciences and analyzed for perchlorate, thiocyanate, and nitrate using ion chromatography tandem mass spectrometry (Blount et al., 2006; Valentin-Blasini et al. 2005). These samples were stored frozen (–70°C) for up to 4 years before perchlorate

analysis. Experiments evaluating storage at –70°C for > 2 years indicated no changes in urinary levels of this analyte (Blount et al. 2006). Reported results for all assays met the division's quality control and quality assurance performance criteria for accuracy and precision [similar to specifications outlined by Westgard et al. (1981)]. Urine samples from the same study participants had previously been analyzed for iodine using inductively coupled plasma mass spectrometry (Caldwell et al. 2005).

Statistical analysis. Initial multiple regression analysis found perchlorate to be a significant predictor of both T_4 and log TSH in women, but perchlorate did not predict either T_4 or log TSH in men (data not shown). Therefore, we present subsequent analysis focused on women.

Of the 1,318 women ≥ 12 years of age, 92 had missing TSH and T_4 values, leaving 1,226. Of these 1,226 women, 91 were excluded from analysis because they reported a history of thyroid disease or current use of thyroid medications, leaving 1,135 women. Of these 1,135 women, 3 had extreme values of T_4 and/or TSH and were excluded. One of these women had a total T_4 of 27 $\mu\text{g}/\text{dL}$ and a TSH of 0.04 IU/L. This woman was clearly hyperthyroid and thus was excluded from the analysis. Two other women had very high TSH levels (43 and 68 IU/L) and were excluded. Of the remaining 1,132 women, 21 had missing perchlorate measurements, leaving a sample size of 1,111 women.

The major design variables for NHANES are age, sex, race/ethnicity, and income related to the poverty level. The values of these variables for the initial 1,318 women and the final 1,111 women, respectively, are as follows: mean age, 41.6 and 39.8 years; percent non-Hispanic whites, 70.8% and 69.4%; percent non-Hispanic blacks, 11.8% and 12.5%; percent Mexican Americans, 7.0% and 7.0%; and percent below the poverty level, 13.9% and 14.9%.

We chose covariates for the multiple regression analyses that are known to be or likely to be associated with T_4 or TSH. We selected a broad number of covariates to evaluate the independence of the perchlorate relationship. These covariates were age, race/ethnicity, body mass index (BMI), serum albumin, serum cotinine (a marker of tobacco smoke exposure), estimated total caloric intake, pregnancy status, postmenopausal status, premenarche status, serum C-reactive protein, hours fasting before sample collection, urinary thiocyanate, urinary nitrate, and use of selected medications.

For these covariates, Table 1 provides means (or geometric means if lognormally distributed) for continuous variables, percent in category for categorical variables, and number of missing results for each covariate. Thyroid

Table 1. Means and percent in category for covariates used in the multiple regression, women ≥ 12 years of age, NHANES 2001–2002.^a

Variable	No.	No. missing	Arithmetic mean (95% CI)	Geometric mean (95% CI)	Percent in category (95% CI)
Age (years)	1,111	0	39.8 (38.1–41.6)		
Fasting (hr)	1,111	0	10.4 (9.85–10.9)		
Serum albumin (g/dL)	1,111	0	4.20 (4.17–4.23)		
Serum T_4 ($\mu\text{g}/\text{dL}$)	1,111	0	8.27 (7.97–8.58)		
Total kilocalories (kcal/1,000)	1,072	39	1.93 (1.87–1.99)		
BMI	1,075	36		25.8 (25.2–26.5)	
Serum cotinine ($\mu\text{g}/\text{L}$)	1,104	7		0.33 (0.23–0.48)	
Serum C-reactive protein (mg/dL)	1,111	0		0.16 (0.14–0.18)	
Serum TSH (IU/L)	1,111	0		1.36 (1.31–1.42)	
Urine creatinine (mg/dL)	1,109	2		81.4 (76.7–86.5)	
Urine iodine ($\mu\text{g}/\text{L}$)	1,111	0		126 (115–138)	
Urine nitrate ($\mu\text{g}/\text{L} \times 1,000$)	1,106	5		38.0 (35.9–40.3)	
Urine perchlorate ($\mu\text{g}/\text{L}$)	1,111	0		2.84 (2.54–3.18)	
Urine thiocyanate ($\mu\text{g}/\text{L} \times 1,000$)	1,104	7		1.20 (1.08–1.33)	
Race/ethnicity					
Non-Hispanic white	1,111	0			69.4 (62.9–75.4)
Non-Hispanic black	1,111	0			12.5 (7.49–19.1)
Mexican American	1,111	0			7.02 (5.14–9.34)
Other race	1,111	0			11.1 (7.04–16.3)
Medication usage					
Furosemide	1,111	0			1.99 (1.25–3.01)
Glucocorticoids and androgens	1,111	0			2.23 (1.24–3.67)
Beta-blocker	1,111	0			4.48 (3.34–5.87)
Estrogen	1,111	0			17.1 (13.2–21.7)
Other drug	1,111	0			1.04 (0.52–1.88)
Menopausal or postmenopausal	1,028	83			35.9 (30.1–41.9)
Pregnant	1,111	0			3.84 (2.74–5.21)
Premenarchal	1,019	92			1.06 (0.48–2.02)

CI, confidence interval.

^aExcludes women with missing TSH, T_4 , or perchlorate, women with history of thyroid disease or taking thyroid drugs, and three women with outlier values of T_4 or TSH (see text).

function has been previously reported to vary with the constitutional variables of age, race, sex, pregnancy, and menopause (Braverman and Utiger 2000). Serum cotinine is a marker of tobacco smoke exposure, and smoking is associated with altered thyroid function (Bertelsen and Hegedus 1994). We included serum C-reactive protein as a marker for inflammatory conditions that have been associated with alterations in thyroid function. Both total caloric intake [based on a 24-hr dietary recall survey and a U.S. Department of Agriculture (USDA) database (Food and Nutrition Database for Dietary Studies; USDA 2004)] and BMI are related to thyroid function, but the interrelationship as to cause or effect is unclear.

Serum albumin was included in our analysis as a possible surrogate for T_4 serum protein binding. NHANES 2001–2002 included total T_4 measurements but not free T_4 measurements; total T_4 varies with the concentrations of specific binding proteins. Concentrations of these proteins can change with physiologic state and health conditions. Free T_4 varies less with such protein concentration changes than does total T_4 . Serum albumin accounts for 15–20% of T_4 binding, with thyroid binding protein and prealbumin (not measured in NHANES) accounting for the remaining percentage (Robbins 2000). Thyroid autoantibody measurements were not available for 2001–2002. For autoantibodies to affect the relationship between perchlorate and T_4 or TSH, presence of autoantibodies would have to correlate with perchlorate levels. We have found no such correlation in the literature and we are unaware of a rationale for such an association.

Medications known to affect thyroid function were also considered. As noted above, women taking medication containing thyroid hormone (e.g., levothyroxine) or antithyroid drugs (e.g., methimazole or propylthiouracil) were excluded. Use of beta-blockers, estrogen formulations, steroids, and furosemide were each modeled using an indicator variable in the regressions. An “other drug” category was also modeled by an indicator variable. This “other drug” category consisted of a heterogeneous group of other medications that have possible effects on thyroid function, protein binding, or measurements, including salicylates, dopaminergics, anticonvulsants and barbiturates, narcotic analgesics, androgenic agents, lithium, and several others (a total of 28 drug codes).

We included the log of urinary creatinine in the models to adjust for variable water excretion. A nonlinear relationship was evaluated by adding the square of the log of perchlorate to final models, but it was not significant. Models were also checked for significance of interaction terms involving main effects. We examined partial regression plots to identify any unduly influential data points; no unduly

influential points were found. Indicator variable coefficients in the models (e.g., for non-Hispanic blacks) were interpreted as follows: 1 = group member, and 0 = not a group member. Urine samples were collected in three sessions of the day from 0800 hours through 2200 hours. Mean perchlorate levels were not statistically different across sessions ($p = 0.49$).

We examined univariate statistics and distribution plots for each dependent and independent variable to look for outliers and to assess the distribution shape. TSH, perchlorate, cotinine, BMI, urinary thiocyanate, urinary nitrate, and C-reactive protein were \log_{10} -transformed to normalize their distributions.

Regression models, including log of perchlorate as one of the predictor variables, were constructed separately for thyroxine and log of TSH. For the initial phase of analysis, we used ordinary least-squares regression (OLS) (SAS Proc Reg, version 9.0; SAS Institute, Cary, NC) and purposefully did not adjust for the NHANES complex survey design in order to obtain a broad group of potentially significant predictor variables. Forward stepwise and backward elimination procedures were used on both population-weighted and unweighted data. The entry p -value for forward elimination models was 0.10 and the retaining p -value for backward elimination was 0.10 in order to identify significant and borderline significant predictors. The forward stepwise and backward elimination approaches produced models that were generally in good agreement.

This OLS analysis produced a generous list of significant and borderline-significant variables for regression analysis using SUDAAN (version 9.0.1; Research Triangle Institute, Research Triangle Park, NC), which provides an analysis that adjusts for the complex survey design. SUDAAN regression models were tested using a manual backward elimination approach starting with the variables obtained from the OLS regression modeling. Selected variables that were excluded in the SUDAAN backward elimination process were added to the final model to ensure they were not significant. The stability of the perchlorate coefficient was monitored during the SUDAAN backward elimination process.

In the main SUDAAN regression analysis, we used population weights to represent women ≥ 12 years of age in the U.S. population for the years 2001 and 2002. In addition, we performed separate regression analyses with SUDAAN using unweighted data and verified that regression coefficients were in good agreement with those obtained using population weights. Reported regression model results in the tables use the population-weighted analysis.

Women were categorized based on a urinary iodine cut point of 100 $\mu\text{g/L}$ and analyzed separately. The 100 $\mu\text{g/L}$ cut point was used based on the World Health Organization

(WHO) definition of sufficient iodine intake in populations (WHO 1994). The WHO noted that the prevalence of goiter begins to increase in populations with median urinary iodine $< 100 \mu\text{g/L}$. A urine iodine level of 100 $\mu\text{g/L}$ represents about the 36th percentile of urinary iodine concentrations in women living in the United States (Caldwell et al. 2005). Women with lower iodine intake could be more vulnerable to perchlorate's effects to impair iodine uptake. From this analysis, the significance of urinary perchlorate as a predictor of thyroid function in women was found to be largely determined by women with urinary iodine $< 100 \mu\text{g/L}$. Consequently, we report here results for women divided into groups based on urinary iodine levels.

Compared to the use of average multiple spot urine measurements or 24-hr urine specimens, the use of a single spot urine for perchlorate and iodine measurement has more imprecision in estimating true urine levels (Andersen et al. 2001). This imprecision is a source of random error (not bias) and therefore decreases statistical power to detect an association between perchlorate and either TSH or T_4 compared to these other urine collection approaches.

Results

For all women ≥ 12 years of age, multiple regression analysis found urinary perchlorate to be a significant predictor of serum TSH and a significant predictor of serum T_4 (data not shown). Because low iodine levels had potential to affect the relationship of perchlorate with T_4 and TSH, women with urinary iodine $< 100 \mu\text{g/L}$ were analyzed separately from women with urinary iodine $\geq 100 \mu\text{g/L}$. Results of this analysis are presented in Tables 2 and 3 for T_4 and in Tables 4 and 5 for TSH.

For women with urinary iodine $< 100 \mu\text{g/L}$, multiple regression analysis found perchlorate to be a significant predictor ($p < 0.0001$) of T_4 with a coefficient for log perchlorate of -0.8917 . The result of regression of T_4 on perchlorate and urinary creatinine without other covariates yielded a coefficient of -0.8604 ($p < 0.0001$). Perchlorate was also a significant predictor ($p = 0.0010$) of log TSH with a coefficient of 0.1230. The result of regression of log TSH on perchlorate and urinary creatinine without other covariates found a coefficient of 0.1117 ($p = 0.0031$). The signs of these coefficients are coherent, with increased perchlorate associated with less production of T_4 and an increase in TSH to stimulate additional T_4 production. For women with urinary iodine $\geq 100 \mu\text{g/L}$, perchlorate was not a significant predictor of T_4 ($p = 0.5503$) but remained a significant predictor of log TSH ($p = 0.0249$). The regression analysis results

in Tables 2–5 include variables that were borderline significant ($0.05 \leq p < 0.10$) to give ample opportunity for other variables to explain variance and better evaluate the independence of the perchlorate effect.

Regression results for men (not shown) indicated that perchlorate was not a significant predictor of either T_4 or log TSH. This finding also held when examining men with urinary iodine levels $< 100 \mu\text{g/L}$.

From the regression coefficients for women with urinary iodine $< 100 \mu\text{g/L}$, we calculated the predicted effect size (i.e., the change in T_4 and TSH) for different levels of perchlorate exposure. We chose perchlorate levels corresponding to the 5th, 10th, 25th, 50th, 75th, 90th, and 95th percentiles of urinary perchlorate in women ≥ 12 years of age. The minimum and maximum perchlorate values are observed results for this population sample;

they are not estimates of the 0th and 100th percentiles for the U.S. population. As such, they would be expected to change in another population sample. The effect size was calculated from the difference between the minimum level of perchlorate measured in women and the level of perchlorate corresponding to the specific percentile. For example, the 50th percentile of urinary perchlorate for women was $2.9 \mu\text{g/L}$ and the minimum level was $0.19 \mu\text{g/L}$. Increasing exposure from $0.19 \mu\text{g/L}$ to $2.9 \mu\text{g/L}$ would result in a predicted decrease in T_4 of $1.06 \mu\text{g/dL}$.

For TSH, one more step is needed in the calculation. Because TSH was modeled as log TSH, the change in TSH from a given change in perchlorate depends on the starting level of TSH. In our calculations we used the approximate 50th and 90th percentiles of TSH as starting points to estimate the predicted perchlorate effect size for TSH. Results of these calculations for T_4 and TSH are presented in Table 6. For comparison, the normal range is $5\text{--}12 \mu\text{g/dL}$ for T_4 and $0.3\text{--}4.5 \text{ IU/L}$ for TSH.

To search for a threshold for the perchlorate relationship with T_4 and TSH, piecewise regression models (Neter et al. 1985) were fit to the data. No inflection point was found for the perchlorate relationship with T_4 or TSH. However, statistical power is limited to detect such a threshold, if present.

Discussion

Increased urinary perchlorate was associated with increased TSH and decreased T_4 for women with urinary iodine levels $< 100 \mu\text{g/L}$, a group possibly more susceptible to competitive inhibition of thyroid iodine uptake by perchlorate. The statistically significant associations of urinary perchlorate with decreased serum T_4 and increased serum TSH were consistent with competitive inhibition of iodide uptake.

For women with urine iodine $\geq 100 \mu\text{g/L}$, perchlorate was also a statistically significant predictor for TSH but not for T_4 . Greater iodine intake may have diminished the effect of perchlorate on T_4 in these women. The significant association with TSH, but not with T_4 , in this group may be due to the greater sensitivity of TSH to impairment of thyroid function; that is, normal T_4 levels are maintained by increasing TSH to compensate for impaired thyroid function.

Predicted changes in serum TSH and T_4 with increasing perchlorate exposure (Table 6) can span a notable portion of the normal medical range of TSH and T_4 values. Compared with a urine level of $0.19 \mu\text{g/L}$, urinary perchlorate of $13 \mu\text{g/L}$ (95th percentile) yields a predicted decrease in T_4 of $1.64 \mu\text{g/dL}$. The normal range for T_4 is $5\text{--}12 \mu\text{g/dL}$. A similar exposure would increase TSH by 2.12 IU/L for a woman starting with a TSH level of 3.11 IU/L (90th percentile for TSH in

Table 2. Regression of serum T_4 on perchlorate and covariates for women ≥ 12 years of age with urine iodine $< 100 \mu\text{g/L}$, NHANES 2001–2002.

Independent variable	Coefficient	SE	p-Value
Intercept	8.6508	0.5428	< 0.0001
Log (urinary perchlorate)	-0.8917	0.1811	< 0.0001
Log (urinary creatinine)	0.6897	0.3338	0.0391
Estrogen use	1.5117	0.4421	0.0007
Log (C-reactive protein)	0.8249	0.1774	< 0.0001
Mexican American ^a	0.6296	0.3684	0.0878
Menopause	-0.5908	0.2578	0.0221
Pregnant (by test)	0.7389	0.3662	0.0439
Total kilocalorie intake (+ 1,000)	-0.3334	0.1173	0.0046
Premenarche	0.6401	0.2722	0.0189

Dependent variable: serum T_4 ($n = 348$; $R^2 = 0.240$).

^aReferent group for race is non-Hispanic white.

Table 3. Regression of serum T_4 on perchlorate and covariates for women ≥ 12 years of age with urine iodine $\geq 100 \mu\text{g/L}$, NHANES 2001–2002.

Independent variables	Coefficient	SE	p-Value
Intercept	10.6652	1.2345	< 0.0001
Log (urinary perchlorate)	0.2203	0.3687	0.5503
Log (urinary creatinine)	1.3138	0.7183	0.0677
Estrogen use	0.8278	0.2722	0.0024
Log (C-reactive protein)	0.5783	0.1247	< 0.0001
Mexican-American ^a	0.5763	0.2522	0.0225
Pregnant (by test)	1.6175	0.3334	< 0.0001
Log (urinary nitrate)	-1.1215	0.4994	0.0249
Hours of fasting	0.0290	0.0156	0.0630

Dependent variable: serum T_4 ($n = 724$; $R^2 = 0.149$).

^aReferent group for race is non-Hispanic white.

Table 4. Regression of serum TSH on perchlorate and covariates for women ≥ 12 years of age with urine iodine $< 100 \mu\text{g/L}$, NHANES 2001–2002.

Independent variables	Coefficient	SE	p-Value
Intercept	0.2654	0.1183	0.0403
Log (urinary perchlorate)	0.1230	0.0373	0.0010
Log (urinary creatinine)	-0.0954	0.0761	0.2103
Beta-blocker use	0.1881	0.0595	0.0016
Estrogen use	-0.0918	0.0404	0.0233
Premenarche	0.1288	0.0262	< 0.0001

Dependent variable: log of serum TSH ($n = 356$; $R^2 = 0.061$).

Table 5. Regression of serum TSH on perchlorate and covariates for women ≥ 12 years of age with urine iodine $\geq 100 \mu\text{g/L}$, NHANES 2001–2002.

Independent variables	Coefficient	SE	p-Value
Intercept	-0.6948	0.3415	0.0600
Log (urinary perchlorate)	0.1137	0.0506	0.0249
Log (urinary creatinine)	-0.1198	0.0910	0.1884
Age in years	0.0025	0.0006	< 0.0001
Log (BMI)	0.4812	0.1346	0.0004
Non-Hispanic black ^a	-0.1125	0.0335	0.0008
Log (urinary nitrate)	0.1087	0.0591	0.0660
Log (urinary thiocyanate)	-0.0816	0.0352	0.0206

Dependent variable: log of serum TSH ($n = 697$; $R^2 = 0.145$).

^aReferent group for race is non-Hispanic white.

women ≥ 12 years of age). The normal range for TSH is 0.3–4.5 IU/L. Effect size estimates that start with the 90th percentile of TSH have more uncertainty than estimates starting with the 50th percentile because the predicted TSH levels fall further from the central portions of the original data.

The mechanism of perchlorate's effect is competitive inhibition of iodide uptake by the thyroid (Clewell et al. 2004; Wolff 1998). Based on this mechanism, individuals with less iodide available to compete with perchlorate may be more vulnerable to impaired iodide uptake. Chronically impaired iodide uptake could lead to changes in serum thyroid hormones, consistent with the increased TSH and decreased T_4 we find associated with increased perchlorate exposure in women with urinary iodine $< 100 \mu\text{g/L}$. The WHO (2004) has identified median urinary iodine levels $\geq 100 \mu\text{g/L}$ as indicating sufficient iodine intake for a population. Based on concerns about adequate iodine intake, the NRC (2005) recently recommended that consideration be given to adding iodine to all prenatal vitamins.

In the present study, perchlorate was not found to be a significant predictor of T_4 or TSH in men. Previous studies report that women have a much higher risk of goiter than do men, especially in populations with marginal iodine intake (Laurberg et al. 2000). The increased vulnerability of women may partially be caused by increased susceptibility to autoimmune thyroid disease in women, the increased demands on the thyroid during pregnancy, or the effect of estrogens on thyroid function. Estradiol has been shown to block TSH-induced sodium/iodide symporter (NIS) expression in the FRTL5 rat follicular cell line (Furlanetto et al. 1999). Impaired NIS expression could lead to reduced ability of the thyroid follicular cells to import iodide, and thus an increased vulnerability to NIS-inhibitors such as perchlorate. Also, estrogens increase T_4 -binding globulin and thus increase the demand for T_4 so that free T_4 levels can remain constant.

Covariates in the regression models predicted T_4 and TSH levels in a manner generally consistent with previous studies. We found that estrogen use was a significant, independent, and positive predictor of T_4 in both low and sufficient iodine models of women ≥ 12 years of age, but was not a significant predictor in either of the TSH models. Similar to estrogen use, pregnancy was a significant or borderline significant predictor of T_4 but not TSH. Both estrogen use and pregnancy raise estrogen levels, increase thyroid binding proteins, and increase serum T_4 concentrations (Glinoe 1997). Menopause lowers estrogen levels and was a significant predictor of T_4 in the regression for women with urinary iodine levels $< 100 \mu\text{g/L}$.

In NHANES III (1988–1994), non-Hispanic blacks were reported to have lower TSH than other groups, and Mexican Americans had higher T_4 levels than non-Hispanic blacks and whites (Hollowell et al. 2002). The models for TSH and T_4 in the present study were consistent with these previous findings concerning race/ethnicity. Non-Hispanic blacks have also been shown to have lower urinary perchlorate levels than non-Hispanic whites, although the reason for this difference is not known (Blount et al. 2006). Age was positively associated with TSH in women with urinary iodine levels $\geq 100 \mu\text{g/L}$, but not significant for women with urinary iodine levels $< 100 \mu\text{g/L}$. A positive association of age and TSH was seen in NHANES III and other studies (Canaris et al. 2000; Hollowell et al. 2002).

BMI was significant in the TSH model for women with urinary iodine levels $\geq 100 \mu\text{g/L}$, and total caloric intake was significant in the T_4 model for women with urinary iodine levels $< 100 \mu\text{g/L}$. Thyroid function clearly has an effect on BMI, as seen clinically and documented in populations (Nyrnes et al. 2006). The reverse is also true, because BMI and total caloric intake can influence the hypothalamic–pituitary–thyroidal axis, although usually at the extremes of body weight and caloric intake (Acheson et al. 1984; Burger et al. 1987; Danforth et al. 1979; Loucks et al. 1992; Loucks and Heath 1994). Total caloric intake in NHANES is a 24-hr recall of food intake. Depending on how well recent intake reflects long-term intake, total caloric intake may parallel the effect of BMI, which was not seen in the present study. Increased caloric intake is known to increase thyroid hormone disposition through deiodination pathways (Burger et al. 1987; Danforth et al. 1979), increasing the conversion of T_4 to the active form, triiodothyronine (T_3), and increasing conversion of T_3 to inactive forms. The effect of changes in calories and carbohydrate composition of the diet on thyroid disposition may have different short- and long-term effects on T_3 and T_4 levels. In the present study, hours of fasting before sample collection was a

borderline significant predictor in one regression model: T_4 in women with sufficient iodine. Fasting for 60 hr can reduce TSH in humans, but fasting for shorter periods has unknown effects on thyroid function.

Beta-blocker drugs are commonly used to treat hypertension and other cardiovascular conditions. Beta-blockers inhibit the conversion of T_4 to the more active form, T_3 , and increase serum TSH (Kayser et al. 1991). Use of these drugs was positively associated with TSH in the regression for women with urinary iodine $< 100 \mu\text{g/L}$. Serum C-reactive protein was positively associated with T_4 in women in each of the iodine groups. C-reactive protein is an acute phase reactant protein increased in many inflammatory conditions in response to production of tissue-generated cytokines, particularly interleukin-6, and has been used as a marker for both specific and systemic low-level inflammation conditions. It is unclear if C-reactive protein is associated with thyroid function other than thyroiditis (Jublanc et al. 2004; Pearce et al. 2003; Tuzcu et al. 2005). However, the stimulus for C-reactive protein, interleukin-6, has a firm inverse relationship with serum T_3 in nonthyroidal illnesses. Also, C-reactive protein and serum T_4 binding proteins are synthesized by the liver; C-reactive protein may vary with an unrecognized health or physiologic condition that affects the synthesis of both proteins. The association of C-reactive protein and T_4 in our study is unclear.

Other variables that are known to possibly affect thyroid function or measurements were not significant predictors in the regression models, including the categories of medications (other than estrogen use and beta-blockers), serum albumin, and serum cotinine. Generally, other medication categories were small and unlikely to have significant effects. Serum albumin did not appear in the final models. Factors such as estrogen use that increase protein binding of thyroid hormones may have accounted for variance in T_4 due to protein binding that serum albumin may have otherwise explained. Serum cotinine is a marker of tobacco smoke exposure, and

Table 6. Predicted change in serum T_4 ^a and serum TSH^b levels based on changes in urinary perchlorate levels in women ≥ 12 years of age, with urine iodine $< 100 \mu\text{g/L}$, NHANES 2001–2002.

Change in urine perchlorate ^d	Change in T_4 ($\mu\text{g/dL}$)	Change in TSH (IU/L) ^c	
		Initial TSH of 1.40 IU/L (50th TSH percentile)	Initial TSH of 3.11 IU/L (90th TSH percentile)
0.19 to 0.65 $\mu\text{g/L}$ (5th percentile)	0.48	0.23	0.51
0.19 to 0.92 $\mu\text{g/L}$ (10th percentile)	0.61	0.30	0.67
0.19 to 1.6 $\mu\text{g/L}$ (25th percentile)	0.83	0.42	0.93
0.19 to 2.9 $\mu\text{g/L}$ (50th percentile)	1.06	0.56	1.24
0.19 to 5.2 $\mu\text{g/L}$ (75th percentile)	1.28	0.70	1.56
0.19 to 9.0 $\mu\text{g/L}$ (90th percentile)	1.49	0.85	1.89
0.19 to 13 $\mu\text{g/L}$ (95th percentile)	1.64	0.95	2.12
0.19 to 100 $\mu\text{g/L}$ (maximum)	2.43	1.63	3.61

^aNormal range for T_4 : 5–12 $\mu\text{g/dL}$. ^bNormal range for TSH: 0.3–4.5 IU/L. ^cDepends on initial TSH level. ^dMinimum level measured, 0.19 $\mu\text{g/L}$.

smoking is associated with altered thyroid function (Belin et al. 2004; Bertelsen and Hegedus 1994). However, tobacco smoke also contains other factors that can inhibit TSH secretion (Bartalena et al. 1995), and perhaps is an explanation for the absence of an association of serum cotinine with either TSH or T₄.

Cyanide in tobacco smoke is metabolized to thiocyanate, a competitive inhibitor of iodide uptake (Tonacchera et al. 2004). Also, nitrate from dietary sources and from formation by intestinal bacteria can compete with iodide. *In vitro* studies indicate that perchlorate is a more potent inhibitor of human NIS, with potencies 15, 30, and 240 times greater than thiocyanate, iodide, and nitrate, respectively (Tonacchera et al. 2004). Thus, the ability of NIS to transport adequate amounts of iodide depends on the relative concentrations of these competing anions. Based on the relative concentrations of perchlorate, nitrate, and thiocyanate likely to be found in human serum, several researchers have predicted that nitrate and thiocyanate are more likely than perchlorate to impair thyroid function (DeGroef et al. 2006; Gibbs 2006). Thiocyanate-induced NIS inhibition is a plausible explanation of the association of smoking with goiter in populations with low iodine intake (Knudsen et al. 2002) and is analogous to the association of perchlorate exposure with thyroid hormone levels observed in our study. However, in women with urinary iodine levels ≥ 100 $\mu\text{g/L}$, urinary thiocyanate was negatively associated with serum TSH, a direction unexpected based on a mechanism of NIS inhibition. The explanation for this is unclear. Urinary nitrate was negatively associated with serum T₄ in women with urinary iodine levels ≥ 100 $\mu\text{g/L}$, a direction consistent with inhibition of NIS. Goitrogenic effects of nitrate intake in animal studies have been observed (Wyngaarden et al. 1953), but there are few studies in humans.

Recently the NRC (2005) evaluated the potential health effects of perchlorate ingestion. Based on studies of long-term treatment of hyperthyroidism and clinical studies of healthy adults, the NRC panel estimated that

a perchlorate dose of > 0.40 mg/kg/day would be required to cause hypothyroidism in adults, although lower doses may lead to hypothyroidism in sensitive subpopulations (NRC 2005).

Comparison of our results to previous studies requires consideration of *a*) target population group studied, *b*) estimated dose of perchlorate, *c*) duration of exposure to perchlorate dose, and *d*) sample size (statistical power). First, for men, we found no relationship with perchlorate and T₄ or TSH. This finding is in general agreement with predicted effects of this level of perchlorate exposure based on reported studies of exposure in men. Lawrence et al. (2000) administered 10 mg perchlorate daily (~ 0.14 mg/kg) to iodine-sufficient adult males for 14 days and found a 10% decrease in radioactive iodine uptake (RAIU), but with no change in TSH or free T₄.

Greer et al. (2002) administered perchlorate to 16 male and 21 female volunteers for 14 days, and found increasing RAIU inhibition for doses between 0.02 and 0.5 mg/kg/day, with no perchlorate-related change in TSH or free T₄. An unknown number of women in that study may have had urinary iodine < 100 $\mu\text{g/L}$, but if the women were typical of the U.S. population (Caldwell et al. 2005), the predicted number of women with low urinary iodine would be 7–8. Braverman et al. (2006) administered perchlorate to 13 iodine-sufficient male and female volunteers at daily doses of 0.5 mg and 3 mg for 6 months, and found no change in RAIU, TSH, or free T₄. Two other studies have also found that workers exposed to perchlorate intermittently for long periods did not have significant changes to serum TSH or T₄ levels (Braverman et al. 2005; Lamm et al. 1999). These study populations were either exclusively (Braverman et al. 2005) or predominantly (Lamm et al. 1999) male.

For women, only two perchlorate studies have focused on women or included a large percentage of women. A recent study of 184 pregnant Chilean women, with mean urinary perchlorate levels near the 99th percentile for women in NHANES 2001–2002, found no perchlorate relationship with thyroid function (Tellez et al. 2005). Of these 184 women,

181 had mean urinary iodine levels ≥ 100 $\mu\text{g/L}$ and only 3 had mean levels < 100 $\mu\text{g/L}$. Therefore, the results of Tellez et al. (2005) would compare to the present results for women with urinary iodine levels ≥ 100 $\mu\text{g/L}$. Urinary iodine levels in the Chilean study population (median 269 $\mu\text{g/L}$) were higher than urinary iodine levels found in the NHANES 2001–2002 population [median 168 $\mu\text{g/L}$; 95% confidence interval, 159–178 $\mu\text{g/L}$]. The Chilean women (Tellez et al. 2005) were also pregnant, which increases the variability in T₄ and TSH. This increased variability would make an association between perchlorate and thyroid function harder to find. The second study with a large percentage of women was Greer et al. (2002) discussed above. These two studies are compared with the present study in Table 7.

Table 7 indicates that our study is the first to target and separately analyze results for women with lower levels of urinary iodine, a potentially susceptible population. A second special attribute of the present study is the much larger sample size of women, affording more statistical power to detect a potential effect. By averaging over many women, the current data likely represents a good approximation of a population steady-state exposure to perchlorate that women have had for a long period of time. If a mid- to long-term exposure is needed for perchlorate to affect thyroid function, this data would have a better opportunity to detect that effect than study designs using short-term exposures. The influence of duration of exposure merits further study.

Accurate assessment of exposure is critical to detect biochemical end points potentially related to exposure. Our laboratory recently developed an improved method for measuring urinary perchlorate, which enhances individual perchlorate exposure assessment (Valentin-Blasini et al. 2005). The use of this new urinary perchlorate measurement strengthens the ability of the present study to detect potential associations with T₄ and TSH.

The present study has the general limitations of a cross-sectional analysis. Therefore, the relationship between urinary perchlorate and thyroid function was examined with attention to the potential influences of chance, bias, or confounding. Perchlorate (as with any of the significant predictor variables) could be a surrogate for another unrecognized determinant of thyroid function. We also assumed in this analysis that urinary perchlorate correlates with levels in the thyroid stroma and tissue, a kinetically distinct compartment. This would be the case in a population with stable, chronic exposures, which is likely but not certain in this population. A large sample size helps to average such potential kinetic differences. Finally, a measurement of free T₄ would be an improvement to the study.

Table 7. Comparison of perchlorate studies targeting women or including a high percentage of women.

	Greer et al. (2002)	Tellez et al. (2005)	Present study
No. of females studied	21 (37 total subjects)	184	1,111
No. of females with urine iodine < 100 $\mu\text{g/L}$	Unknown (estimate 7–8)	3 ^a	348, T ₄ analysis 356, TSH analysis
Females with urine iodine < 100 $\mu\text{g/L}$ analyzed separately	No	No	Yes
Perchlorate dose and duration of exposure	Up to 0.5 mg/kg/day for 14 days	Long-term environmental exposure	Long-term environmental exposure
Comments		All women pregnant, increasing variability of T ₄ and TSH	

^aAverage of one to three spot urine samples.

Conclusions

Urinary perchlorate is associated with an increased TSH and decreased total T_4 in women ≥ 12 years of age with urine iodine levels $< 100 \mu\text{g/L}$ in the U.S. population during 2001–2002. For women with urine iodine levels $\geq 100 \mu\text{g/L}$, urine perchlorate is a significant predictor of TSH but not T_4 . These effects of perchlorate on T_4 and TSH are coherent in direction and independent of other variables known to affect thyroid function, but are found at perchlorate exposure levels that were unanticipated based on previous studies. Further research is recommended to affirm these findings.

REFERENCES

- Acheson K, Jequier E, Burger A, Danforth E. 1984. Thyroid hormones and thermogenesis: the metabolic cost of food and exercise. *Metabolism* 33:262–265.
- Andersen S, Pedersen KM, Pedersen IB, Laurberg P. 2001. Variations in urinary iodine excretion and thyroid function. A 1-year study in healthy men. *Eur J Endocrinol* 144:461–465.
- Bartalena L, Bogazzi F, Tanda ML, Manetti L, Dell'Unto E, Martino E. 1995. Cigarette smoking and the thyroid. *Eur J Endocrinol* 133:507–512.
- Belin RM, Astor BC, Powe NR, Ladenson PW. 2004. Smoke exposure is associated with a lower prevalence of serum thyroid autoantibodies and thyrotropin concentration elevation and a higher prevalence of mild thyrotropin concentration suppression in the third National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab* 89:6077–6086.
- Bertelsen JB, Hegedus L. 1994. Cigarette smoking and the thyroid. *Thyroid* 4:327–331.
- Blount BC, Valentin-Blasini L, Osterloh JD, Mauldin JP, Pirkle JL. 2006. Perchlorate exposure of the U.S. population, 2001–2002. *J Exp Sci Environ Epidemiol*; doi:10.1038/sj.jes.7500535 [Online 18 October 2006].
- Braverman LE, He X, Pino S, Cross M, Magnani B, Lamm SH, et al. 2005. The effect of perchlorate, thiocyanate, and nitrate on thyroid function in workers exposed to perchlorate long-term. *J Clin Endocrinol Metab* 90:700–706.
- Braverman LE, Pearce EN, He X, Pino S, Seely M, Beck B, et al. 2006. Effects of six months of daily low-dose perchlorate exposure on thyroid function in healthy volunteers. *J Clin Endocrinol Metab* 91:2721–2724; doi:10.1210/jc.2006-0184 [Online 24 April 2006].
- Braverman LE, Utiger RD. 2000. Introduction to hypothyroidism. In: *Werner & Ingbar's The Thyroid: A Fundamental and Clinical Text*. (Braverman LE, Utiger RD, eds). 8th ed. Philadelphia:Lippincott Williams & Wilkins, 719–720.
- Burger AG, O'Connell M, Scheidegger K, Woo R, Danforth E. 1987. Monodeiodination of triiodothyronine and reverse triiodothyronine during low and high calorie diets. *J Clin Endocrinol Metab* 65:829–835.
- Caldwell KL, Jones R, Hollowell JG. 2005. Urinary iodine concentration: United States National Health and Nutrition Examination Survey 2001–2002. *Thyroid* 15:692–699.
- Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. 2000. The Colorado thyroid disease prevalence study. *Arch Intern Med* 160:526–534.
- CDC (Centers for Disease Control and Prevention). 2003. Laboratory Procedure Manual. Available: http://www.cdc.gov/nchs/data/nhanes/nhanes_01_02/140t4_b_met_b_t4.pdf [accessed 20 March 2006].
- CDC (Centers for Disease Control and Prevention). 2004. National Health and Nutrition Examination Survey. Available: <http://www.cdc.gov/nchs/nhanes.htm> [accessed 20 March 2006].
- Clewell RA, Merrill EA, Narayanan L, Gearhart JM, Robinson PJ. 2004. Evidence for competitive inhibition of iodide uptake by perchlorate and translocation of perchlorate into the thyroid. *Int J Toxicol* 23:17–23.
- Danforth E Jr, Horton ES, O'Connell M, Sims EA, Burger AG, Ingbar SH, et al. 1979. Dietary-induced alterations in thyroid hormone metabolism during overnutrition. *J Clin Invest* 64:1336–1347.
- Dasgupta PK, Martinelango PK, Jackson WA, Anderson TA, Tian K, Tock RW, et al. 2005. The origin of naturally occurring perchlorate: the role of atmospheric processes. *Environ Sci Technol* 39:1569–1575.
- DeGroef B, Decallonne BR, van der Geyten S, Darras VM, Bouillon R. 2006. Perchlorate versus other environmental sodium/iodide symporter inhibitors: potential thyroid-related health effects. *Eur J Endocrinol* 155:17–25.
- Furlanetto TW, Nguyen LQ, Jameson JL. 1999. Estradiol increases proliferation and down-regulates the sodium/iodide symporter gene in FRTL-5 cells. *Endocrinology* 140:5705–5711.
- Gibbs JP. 2006. A comparative toxicological assessment of perchlorate and thiocyanate based on competitive inhibition of iodide uptake as the common mode of action. *Hum Ecol Risk Assess* 12:157–173.
- Gliozzi D. 1997. The regulation of thyroid function in pregnancy: pathways of endocrine adaptation from physiology to pathology. *Endocr Rev* 18:404–433.
- Gliozzi D. 2000. Thyroid disease during pregnancy. In: *Werner & Ingbar's The Thyroid: A Fundamental and Clinical Text* (Braverman LE, Utiger RD, eds). Philadelphia:Lippincott Williams & Wilkins, 1013–1027.
- Greer MA, Goodman G, Pleus RC, Greer SE. 2002. Health effects assessment for environmental perchlorate contamination: the dose response for inhibition of thyroidal radioiodine uptake in humans. *Environ Health Perspect* 110:927–937.
- Haddow JE, Palomaki GE, Allan WC, Williams JR, Knight GJ, Gagnon J, et al. 1999. Maternal thyroid deficiency during pregnancy and subsequent neuropsychological development of the child. *N Engl J Med* 341:549–555.
- Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, et al. 2002. Serum TSH, T_4 , and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab* 87:489–499.
- Jackson A, Arunagiri S, Tock R, Anderson TA, Rainwater K. 2004. Electrochemical generation of perchlorate in municipal drinking water systems. *J Am Water Works Assoc* 96:103–108.
- Jackson WA, Joseph P, Laxman P, Tan K, Smith PN, Yu L, et al. 2005. Perchlorate accumulation in forage and edible vegetation. *J Agric Food Chem* 53:369–373.
- Jubanc C, Bruckert E, Giral P, Chapman MJ, Leenhardt L, Carreau V, et al. 2004. Relationship of circulating C-reactive protein levels to thyroid status and cardiovascular risk in hyperlipidemic euthyroid subjects: low free thyroxine is associated with elevated hsCRP. *Atherosclerosis* 172:7–11.
- Kayser L, Perrild H, Feldt-Rasmussen U, Hegedus L, Skovsted L, Hansen JE. 1991. The thyroid function and size in healthy man during 3 weeks treatment with beta-adrenoceptor-antagonists. *Horm Metab Res* 23:35–37.
- Kirk AB, Martinelango PK, Tian K, Dutta A, Smith EE, Dasgupta PK. 2005. Perchlorate and iodide in dairy and breast milk. *Environ Sci Technol* 39:2011–2017.
- Klein RZ, Sargent JD, Larsen PR, Waitsbren SE, Haddow JE, Mitchell ML. 2001. Relation of severity of maternal hypothyroidism to cognitive development of offspring. *J Med Screen* 8:18–20.
- Knudsen N, Bulow I, Laurberg P, Ovesen L, Perrild H, Jorgensen T. 2002. Association of tobacco smoking with goiter in a low-iodine-intake area. *Arch Intern Med* 162:439–443.
- Lamm SH, Braverman LE, Li FX, Richman K, Pino S, Howarth G. 1999. Thyroid health status of ammonium perchlorate workers: a cross-sectional occupational health study. *J Occup Environ Med* 41:248–260.
- Laurberg P, Nohr SB, Pedersen KM, Hreidarsson AB, Andersen S, Bulow-Pedersen I, et al. 2000. Thyroid disorders in mild iodine deficiency. *Thyroid* 10:951–963.
- Lawrence JE, Lamm SH, Pino S, Richman K, Braverman LE. 2000. The effect of short-term low-dose perchlorate on various aspects of thyroid function. *Thyroid* 10:659–663.
- Loucks AB, Heath EM. 1994. Induction of low- T_3 syndrome in exercising women occurs at a threshold of energy availability. *Am J Physiol* 266:R817–R823.
- Loucks AB, Laughlin GA, Mortola JF, Girton L, Nelson JC, Yen SS. 1992. Hypothalamic-pituitary-thyroidal function in eumenorrheic and amenorrheic athletes. *J Clin Endocrinol Metab* 75:514–518.
- Mendiratta SK, Dotson RL, Brooker RT. 1996. Perchloric acid and perchlorates. In: *Kirk-Othmer Encyclopedia of Chemical Technology*, Vol 18 (Kroschwitz JJ, Howe-Grant M, eds). 4th ed. New York:John Wiley & Sons, Inc, 157–170.
- National Research Council. 2005. Health Implications of Perchlorate Ingestion. Washington, DC:National Academy Press.
- Neter J, Wasserman W, Kutner M. 1985. Applied Linear Statistical Models. 2nd ed. Homewood, IL:Richard D. Irwin, Inc.
- Nyrenes A, Jorde R, Sundsfjord J. 2006. Serum TSH is positively associated with BMI. *Int J Obes (Lond)* 30:100–105.
- Pearce EN, Bogazzi F, Martino E, Brogioni S, Pardini E, Pellegrini G, et al. 2003. The prevalence of elevated serum C-reactive protein levels in inflammatory and noninflammatory thyroid disease. *Thyroid* 13:643–648.
- Robbins J. 2000. Thyroid hormone transport proteins and the physiology of hormone binding. In: *Werner & Ingbar's The Thyroid: A Fundamental and Clinical Text* (Braverman LE, Utiger RD, eds). Philadelphia:Lippincott Williams & Wilkins, 105–120.
- Sanchez CA, Krieger RI, Khandaker N, Moore RC, Holts KC, Neidel LL. 2005. Accumulation and perchlorate exposure potential of lettuce produced in the Lower Colorado River region. *J Agric Food Chem* 53:5479–5486.
- Sanchez CA, Krieger RI, Khandaker N, Valentin-Blasini L, Blount BC. 2006a. Potential perchlorate exposure from citrus sp. irrigated with contaminated water. *Analytica Chimica Acta* 567:33–38.
- Sanchez CA, Krieger RI, Valentin-Blasini L, Blount BC, Khandaker N. 2006b. Perchlorate accumulation and potential exposure from durum wheat irrigated with Colorado River water. *J ASTM Int*; doi:10.1520/JAI100397 [Online 28 June 2006].
- Snyder SA, Pleus RC, Vanderford BJ, Holady JC. 2006. Perchlorate and chlorate in dietary supplements and flavor enhancing ingredients. *Analytica Chimica Acta* 567:26–32; doi:10.1016/j.aca.2006.03.029 [Online 16 March 2006].
- Tellez RT, Chacon PM, Abarca CR, Blount BC, Landingham CB, Crump KS, et al. 2005. Long-term environmental exposure to perchlorate through drinking water and thyroid function during pregnancy and the neonatal period. *Thyroid* 15:963–975.
- Tonacchera M, Pinchera A, Dimida A, Ferrarini E, Agretti P, Vitti P, et al. 2004. Relative potencies and additivity of perchlorate, thiocyanate, nitrate, and iodide on the inhibition of radioactive iodide uptake by the human sodium iodide symporter. *Thyroid* 14:1012–1019.
- Tuzcu A, Bahceci M, Gokalp D, Tuzun Y, Gunes K. 2005. Subclinical hypothyroidism may be associated with elevated high-sensitive C-reactive protein (low grade inflammation) and fasting hyperinsulinemia. *Endocr J* 52:89–94.
- Urbansky ET, Brown SK, Magnuson ML, Keltly CA. 2001. Perchlorate levels in samples of sodium nitrate fertilizer derived from Chilean caliche. *Environ Pollut* 112:299–302.
- USDA (U.S. Department of Agriculture). 2004. USDA Food and Nutrient Database for Dietary Studies, 1.0. Beltsville, MD:Agricultural Research Service, Food Surveys Research Group. Available: <http://www.ars.usda.gov/Services/docs.htm?docid=7673> [accessed 20 March 2006].
- U.S. EPA (U.S. Environmental Protection Agency). 2005. Occurrence Data: Accessing Unregulated Contaminant Monitoring Data. Available: <http://www.epa.gov/safewater/ucmr/data.html> [accessed 20 March 2006].
- Valentin-Blasini L, Mauldin JP, Maple D, Blount BC. 2005. Analysis of perchlorate in human urine using ion chromatography and electrospray tandem mass spectrometry. *Anal Chem* 77:2475–2481.
- Westgard JO, Barry PL, Hunt MR, Groth T. 1981. A multi-rule Shewhart chart for quality control in clinical chemistry. *Clin Chem* 27:493–501.
- WHO (World Health Organization). 1994. Indicators for Assessing Iodine Deficiency Disorders and Their Control through Salt Iodization. WHO/NUT/94.6. Geneva:World Health Organization/International Council for the Control of Iodine Deficiency Disorders.
- Wolff J. 1998. Perchlorate and the thyroid gland. *Pharmacol Rev* 50:89–105.
- Wyngaarden JB, Stanbury JB, Rapp B. 1953. The effects of iodide, perchlorate, thiocyanate and nitrate administration upon the iodide concentrating mechanism of the rat thyroid. *Endocrinology* 52:568–574.