

Spatial Modeling of PM₁₀ and NO₂ 1985-2000 in the Continental US Online Supplement

Jaime E. Hart ^{1,2,3}, Jeff D. Yanosky ², Robin C. Puett ^{1,4,5}, Louise Ryan ⁶, Douglas W. Dockery ^{1,3}, Thomas J. Smith ¹, Eric Garshick ^{2,7}, Francine Laden ^{1,2,3}

¹Exposure, Epidemiology and Risk Program, Department of Environmental Health, Harvard School of Public Health, Boston, MA

²Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

³Department of Epidemiology, Harvard School of Public Health, Harvard School of Public Health, Boston, MA


⁴South Carolina Cancer Prevention and Control Program, University of South Carolina, Columbia, SC

⁵Departments of Environmental Health Sciences and Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC

⁶Department of Biostatistics, Harvard School of Public Health, Boston, MA

⁷Pulmonary and Critical Care Medicine Section, Medical Service, VA Boston Healthcare System, Boston, MA

Supplement Figure 1: Measured versus model predicted PM₁₀ (μg/m³) from the model cross-validation



Supplement Figure 2: Measured versus model predicted NO₂ (ppb) from the model cross-validation

