



ENVIRONMENTAL
HEALTH
PERSPECTIVES

<http://www.ehponline.org>

**Air Pollution Exposures During Adulthood and Risk of
Endometriosis in the Nurses' Health Study II**

**Shruthi Mahalingaiah, Jaime E. Hart, Francine Laden,
Ann Aschengrau, and Stacey A. Missmer**

<http://dx.doi.org/10.1289/ehp.1306627>

Received: 7 February 2013

Accepted: 8 November 2013

Advance Publication: 13 November 2013

Air Pollution Exposures During Adulthood and Risk of Endometriosis in the Nurses' Health Study II

Shruthi Mahalingaiah,¹ Jaime E. Hart,^{2,3} Francine Laden,^{2,3,4} Ann Aschengrau,⁵ and Stacey A. Missmer^{2,4,6}

¹Department of Obstetrics and Gynecology, Boston University School of Medicine, Boston, Massachusetts, USA;

²Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA;

³ Exposure, Epidemiology, and Risk Program, Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts, USA;

⁴ Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts, USA;

⁵ Department of Epidemiology, Boston University School of Public Health, Boston, Massachusetts, USA;

⁶ Department of Obstetrics, Gynecology, and Reproductive Biology, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA.

Address correspondence to:

Shruthi Mahalingaiah, MD

Department of Obstetrics and Gynecology

Boston University School of Medicine/ Boston Medical Center

85 East Concord St. 6th Floor

Boston, MA 02118

Phone: 1-617-414-7305

Fax: 1-617-414-7300

Email: shruthi@bu.edu

Running Title: Air pollution and Endometriosis Risk

Competing Financial Interests: The authors have no actual or potential competing financial interests to disclose.

Acknowledgments: The authors would like to thank Tam Nguyen for programming assistance and the participants of the Nurses' Health Study II for their continued enthusiastic participation. Supported by grants 5K12HD043444-10 and R01HD57210 from the National Institute of Child Health and Human Development, R01CA50385 from the National Cancer Institute, 5R01ES017017 from the National Institute for Environmental Health Sciences, a research grant from the Boston University Department of Obstetrics and Gynecology, and the Massachusetts Institute of Technology Center for Environmental Health Sciences (MIT CEHS) Translational Pilot Project Program. The Nurses' Health Study II is additionally supported by the infrastructure grant UM1 CA176726 from the National Cancer Institute.

ABSTRACT

Background: Particulate matter and proximity to large roadways may promote disease mechanisms, including systemic inflammation, hormonal alteration, and vascular proliferation, that may contribute to the development and severity of endometriosis.

Objective: To determine the association of air pollution exposures during adulthood, including distance to road, particulate matter less than 10 microns, less than 2.5 microns, and between 2.5 and 10 microns ($PM_{2.5}$, $PM_{10-2.5}$, PM_{10}), and timing of exposure with risk of endometriosis in the Nurses' Health Study II.

Methods: Proximity to major roadways and outdoor levels of $PM_{2.5}$, $PM_{10-2.5}$, PM_{10} were determined for all residential addresses from 1993 to 2007. Multivariable-adjusted time-varying Cox proportional hazard models were used to estimate the relation between these air pollution exposures and endometriosis risk.

Results: Among 84,060 women 2,486 incident cases of surgically confirmed endometriosis were identified over 710,230 person-years of follow-up. There was no evidence of an association, between distance to road, $PM_{2.5}$, $PM_{10-2.5}$, PM_{10} averaged over follow-up or during the previous 2- or 4- year period and endometriosis risk.

Conclusions: Traffic and air pollution exposures during adulthood were not associated with incident endometriosis in this cohort of women.

Introduction

Endometriosis is defined as the presence of endometrial-like tissue outside of the lining of the uterus and may be asymptomatic, or accompanied by menstrual cycle pain and pelvic pain of varying severity. Surgical visualization is the gold-standard for establishing a diagnosis of endometriosis. However, patients may be given a diagnosis based on clinically suspected disease based on history, physical exam, and imaging findings (Hsu et al. 2010). Endometriosis is staged based on the extent of surface area and locations involved from mild to severe disease. The reported prevalence of endometriosis varies by clinical population and method of ascertainment from 4-50% (Abbas et al. 2012). The prevalence of undiagnosed endometriosis of all stages in the general population has been estimated at 11%, (Buck Louis et al. 2011). However, the likely prevalence of severe endometriosis is less than 2% (Zondervan et al. 2002). In a study of white females, 15-49 years of age in Rochester, Minnesota from 1970-1979 presenting to area hospitals and diagnosed with endometriosis, the authors describe the incidence of endometriosis by level of diagnostic certainty. There were 252 cases /100,000 person years of endometriosis defined by the gold standard (histologically confirmed or surgically visualized) (Houston et al. 1987). In a previous study evaluating *in utero* exposures and incidence of endometriosis of the first 10 years of follow-up in the Nurses' Health Study II (NHSII) cohort, the incidence of self-reported laparoscopically confirmed endometriosis was 298 cases/100,000 person-years among women aged between 25-42. (Missmer et al. 2004b).

Endometriosis is observed rarely in non-human primates, though animal models have been developed using transplantation of autologous endometrial tissue or human endometrial tissue into the peritoneum (Grummer 2006). In monkeys (*macaca mulatta* and rhesus), endometriosis has been induced through long-term exposure to radiation (Fanton and Golden 1991; Splitter et

al. 1972) and chronic exposure to dioxin-compounds (Rier et al. 1993), with a minimum latency period of 7 years (Fanton and Golden 1991; Rier et al. 1993; Wood et al. 1983). There is limited data in humans regarding the etiologically relevant time window of exposure.

Several factors may contribute to endometriosis development and disease severity, including anatomic (Breech and Laufer 1999), anthropometric (Missmer et al. 2004a), hormonal, immunologic (Siristatidis et al. 2006), inflammatory, and genetic factors (Augoulea et al. 2012; Nyholt et al. 2012). Inflammatory markers are elevated in women with endometriosis (Gentilini et al. 2011); a state of inflammation is considered to be both a result of the disease and may promote disease progression (Agic et al. 2006).

Air pollution exposures include traffic related exhaust (diesel and non-diesel) as well as particulate matter (PM) characterized by its size fraction. Smaller sizes of particulate matter, such as 2.5 microns in diameter (PM_{2.5}) or smaller, can cross into the blood stream, deposit at distant tissues, and promote local and systemic inflammation (Brook et al. 2010; Calderón-Garcidueñas et al. 2011). Laboratory studies have demonstrated that diesel exhaust particles have hormonal activity in estrogenic and androgenic activity assays (Misaki et al. 2008; Oh et al. 2008; Sidlova et al. 2009; Wang et al. 2005). In a rat model of the *in utero* and postnatal effects of diesel exhaust on endometriosis, exposure to diesel exhaust enhanced persistence of endometriosis lesions, defined by persistence of surrounding collagen fibers (Umezawa et al. 2011). In the present study, we evaluated the association of exposure to traffic related exhaust and PM during adulthood with the incidence of endometriosis in the NHSII cohort.

Methods

Study Population

Women in this study were participants in the Nurses' Health Study II (NHSII), an ongoing prospective cohort study with a one-time enrollment in 1989 and ten completed biennial follow-up surveys through May 2007. A total of 116,687 female registered nurses completed the baseline questionnaire. Nurses ranged in age from 25 to 43 years and initially resided in 14 of the United States, including California, Connecticut, Indiana, Iowa, Kentucky, Massachusetts, Michigan, Missouri, New York, North Carolina, Ohio, Pennsylvania, South Carolina, and Texas. However, as of the mid-1990s, they have resided in all 50 states and the District of Columbia (Figure 1). Questionnaires were used to collect information regarding the incidence of disease outcomes and a variety of biologic, environmental, dietary, and life-style risk factors. Surveys used to conduct this analysis were updated and mailed biennially with a >90% response in each questionnaire cycle. Of the initial enrolled cohort, 93% were Caucasian – representative of the ethnic diversity of nurses in the US at enrollment – and most resided in urban areas in the Northeastern and Midwestern United States. Women were included in the current study if they were alive, continued to respond to questionnaires throughout the course of follow-up, and did not have a diagnosis of endometriosis prior to 1993. In order to assess exposure, the women also had to have at least one home address within the continental United States that could be geocoded to the street segment level. The study was approved by the Institutional Review Boards (IRB) of Boston University School of Medicine/Boston Medical Center and Brigham and Women's Hospital. Informed consent was implied through the return of questionnaires.

Case ascertainment

In 1993, the participants were asked if they “ever had physician-diagnosed endometriosis.” If they responded “yes,” they were asked to indicate whether the diagnosis was laparoscopically confirmed, as surgical diagnosis is the current diagnostic gold standard (Hsu et al. 2010). These questions were asked again on all subsequent biennial questionnaires. As described previously (Missmer et al. 2004a), a validation study was performed in 1994 based on a medical record review of a random sample of the 1,766 nurses who reported physician-diagnosed endometriosis after enrollment into the NHSII. The 131 women whose records were reviewed included 105 who reported a laparoscopic diagnosis of endometriosis. Among these women, medical record review confirmed 93 cases, and the other 12 received clinical care for endometriosis, despite having no visual confirmation of disease during laparoscopy. Of the 26 women who did not report laparoscopic diagnosis, records indicated a clinical diagnosis of endometriosis in 14, and no evidence of a diagnosis in the remaining 12 (Missmer et al. 2004a).

Women who reported endometriosis on the 1993 questionnaire were excluded from the present study. Women who reported a laparoscopic diagnosis of endometriosis after 1993 through 2007 were included as cases. Women who reported a clinical diagnosis of endometriosis without laparoscopy did not contribute any additional person-time but were allowed to re-enter the analysis in the future as a case if endometriosis was later confirmed by laparoscopy. The midpoint between the receipts of the questionnaire before and after diagnosis was assigned as the date of diagnosis.

Exposure Assessment

Residential address information was updated in the NHSII cohort every two years as part of the questionnaire mailing process, and was geocoded to obtain latitude and longitude for each mailing address from 1989-2007. We used geographic information system (GIS) software (ArcGIS, version 9.2; ESRI, Redlands, CA) to calculate the distance to the nearest road (in meters) at each residential address as a proxy measure of all exposures related to traffic. Road segments in the 2000 U.S. Census Topologically Integrated Geographic Encoding and Referencing system (TIGER) (U.S.CensusBureau 1992) files were selected by U.S. Census feature class code to include: A1 (primary roads, typically interstate highways, with limited access, division between the opposing directions of traffic, and defined exits), A2 (primary major, non-interstate highways and major roads without access restrictions), or A3 (smaller, secondary roads, usually with more than two lanes) road segments. The shortest distance between each address and the closest road segment was calculated. Analyses were conducted using the distance to the closest of all three road types and distance to the two largest road types (A1, A2). Based on the distribution of distance to road in this cohort and previous exposure studies showing exponential decay in exposures with increasing distance from a road, for our primary analyses we categorized distance to road as 0–50, 50–200, or ≥ 200 m (Adar and Kaufman 2007; Karner et al. 2010; Lipfert et al. 2006; Lipfert and Wyzga 2008; Sahlodin et al. 2007; Zhu et al. 2002). We also considered cut-points up to 500 m in sensitivity analyses.

Predicted ambient exposure to particulate matter less than 10 microns (PM_{10}) and less than 2.5 microns ($PM_{2.5}$) in aerodynamic diameter was available for each month since January 1988 through 2007 at the home address(es) of each cohort member. These values were generated from nationwide expansions of previously validated spatiotemporal models (Yanosky et al. 2008a;

Yanosky et al. 2009). The models used monthly average PM_{10} and/or $PM_{2.5}$ data from USEPA's Air Quality System, a nationwide network of continuous and filter-based monitors, as well as monitoring data from various other sources. The models also used GIS to incorporate information on several geospatial predictors. All PM data and GIS data were used in generalized additive statistical models (Yanosky et al. 2008b) with smooth terms of space and time to create separate PM prediction surfaces for each month. Since monitoring data on $PM_{2.5}$ are limited prior to 1999, $PM_{2.5}$, before 1999, was modeled using data on PM_{10} and airport visibility data (Yanosky et al. 2009). $PM_{2.5-10}$ was estimated by subtracting monthly average values for $PM_{2.5}$ from those for PM_{10} . We calculated three different time-varying exposure measures: the average air pollution in the prior 2 calendar years, the average air pollution in the prior 4 calendar years, and the cumulative average exposure.

Additional Covariates

We examined possible confounding by age (in months), race/ethnicity (African American, Asian, Caucasian, Hispanic, Other), age at menarche, and, updated biennially – smoking status (current/former/never), body mass index (BMI), parity, oral contraception use, infertility (ever and current), ever performed rotating shift work (Schernhammer et al. 2011), region (Northeast, Midwest, West, South), and area level socioeconomic status (Census tract level, median home value and median family income). Variables selected for potential inclusion in the model included those identified in prior studies to correlate with endometriosis (parity, age at menarche) (Missmer et al. 2004a) or with exposure (elevation, land use) (Hart et al. 2009). Each potential confounder (or set of indicator variables) was added separately to a model that included age and race, *a priori*. We defined confounders as variables that changed the coefficient of the

main effect of traffic exposure or PM by at least 10% when added to this basic model (Greenland 1989).

Statistical Analysis

Time-varying Cox proportional hazards models were used to assess the relation of laparoscopically confirmed endometriosis with exposure to traffic or particulate matter. Person-time accrued from September 1, 1993 until first surgical diagnosis of endometriosis, hysterectomy, menopause, loss-to-follow-up, cancer diagnosis (other than non-melanoma skin cancer), date of death, or the end of follow-up, whichever occurred first. Person-time was calculated starting in 1993. This allowed for the calculation of up to 4 years of previous particulate exposure from baseline enrollment in 1989 and for the inclusion of prospective cases only. Person-time was excluded from follow-up for any period in which the home address was outside of the continental United States or was unable to be geocoded to the street segment level. Hazard ratios (HRs) and 95% CIs were calculated for categories of distance to road or for each 10- $\mu\text{g}/\text{m}^3$ increase in PM. All Cox models were stratified by age in months and calendar year.

To determine if associations of traffic exposure and air pollution with endometriosis differed by personal characteristics, we examined effect modification by parity (nulliparous or parous), overweight/obesity (BMI >25 or ≤ 25 kg/m²), smoking status (ever or never smoker), age at menarche (<12 vs. 12 or older), infertility (current or ever versus never), and rotating shift work (ever vs. never). For each characteristic we performed time-varying stratified models (with the exception of age at menarche) to obtain strata-specific effect estimates and 95% CIs. To determine if there was statistically significant ($p < 0.05$) effect modification, we also ran models

that included multiplicative interaction terms. To account for potential regional differences in traffic volume or PM composition we also performed models stratified by region of the country.

The relation between endometriosis and infertility is complex. Of the women who have both endometriosis and infertility, endometriosis may be either causative or incidental (Missmer et al. 2004a). Furthermore, women with infertility may undergo more diagnostic testing including laparoscopic evaluation that may identify endometriosis. Because there may be a diagnostic bias in women with infertility, to reduce this bias, we performed a sensitivity analysis restricted to those without infertility in two ways 1) reported at anytime during follow-up and 2) ever vs never. Statistical analyses were performed in SAS version 9.2 (SAS Institute, Cary, NC). An alpha level of 0.05 was used to define statistical significance.

Results

A total of 84,060 women were in this study population for analyses of residential proximity to road, PM₁₀, PM_{10-2.5}, and PM_{2.5}. The mean (\pm standard deviation) age over the full course of follow-up was 42.0 years (\pm 5.1), the cohort was mostly parous, and over two-thirds were never smokers. Age-standardized characteristics of the cohort for the full duration of follow-up are presented in Table 1. There was little difference in covariates among the distance to road categories, with the exception of decreasing parity levels nearer to roadways and more women living nearer to roadways in the Northeast. The mean and median levels of each of the particulate matter metrics were similar for each averaging period, and there were wide ranges in all pollutants (Table 2).

Over 710,230 person-years of follow-up, there were a total of 2,486 incident cases of laparoscopically confirmed endometriosis. In the basic models adjusted only for age and calendar time, living closer to a roadway than the referent, regardless of specific road type or cut-point categorization used, was generally associated with small nonstatistically significant elevations in the risk of endometriosis, compared to living farther from a roadway (Table 3). Age, parity, oral contraception use, age at menarche, region, and area-level socioeconomic status were the variables that met our definition of confounding most consistently. In fully adjusted models, associations were slightly attenuated. In region specific analyses, non-significant positive associations were primarily evident in the Northeast. Particulate matter exposures were not positively associated with endometriosis in the population as a whole (Table 4). Models excluding women reporting infertility (including 1,578 incident cases over 565,243 person-years of follow-up) also demonstrated similar results (data not shown).

There was no evidence of effect modification in the distance to road models (data not shown). As shown in Figure 2, risks for a $10 \mu\text{g}/\text{m}^3$ increase in PM were elevated among the nulliparous, ever smokers, women with an age of menarche 12 or older, women with a BMI less than 25, and women diagnosed with infertility. However, only a few interactions were statistically significant.

Discussion

In this cohort of US women, there was no statistically significant increased risk of endometriosis with exposures to traffic or ambient particulate matter ($\text{PM}_{2.5}$, PM_{10} , $\text{PM}_{10-2.5}$) exposure during adulthood. To the best of our knowledge, this analysis represents the first human study to assess the relation between PM exposure and endometriosis.

Though many theories have been proposed, retrograde menstruation is the most widely accepted etiology for the initial establishment of peritoneal endometriosis (Sampson 1927). As up to 85% of women have retrograde menstruation, it is further hypothesized that endometriosis occurs in susceptible women in the presence of abnormal endometrial cells (Flores et al. 2007; Halme et al. 1984; Kruitwagen et al. 1991). As described by Flores et al, viable endometrial cells deposited in the peritoneum adhere to peritoneal surfaces and invade under the stimulation of inflammatory cells and cytokines (Flores et al. 2007). Lesions then continue to infiltrate, develop a blood supply, and synthesize and secrete hormones (exact hormonal milieu depends on each lesion's secretory profile) (Takeda et al. 2004). Systemic and local inflammation and oxidative stress (Andrade et al. 2010; Maybin et al. 2011) and hormonal alterations (including altered response to hormones and increased lesion hormone synthesis) are proposed mechanisms for endometriosis disease progression (Burney et al. 2007; Pan et al. 2007).

Air pollution has been shown to promote both local and systemic inflammation (*in-vivo*) and increase oxidative stress (Brook et al. 2004; Brook et al. 2010), as well as possess hormonal activity through hormone receptor binding (*in-vitro*) (Misaki et al. 2008; Oh et al. 2008; Sidlova et al. 2009; Wang et al. 2005). As such, air pollution may engage several proposed pathways in endometriosis disease progression. In human studies and non-human primate animal models of endometriosis, there are elevations of proinflammatory cytokines in the peritoneal fluid in diseased subjects (Chen et al. 2010; D'Hooghe et al. 2001; D'Hooghe and Debrock 2002), as well as increased oxidative stress (Van Langendonck et al. 2002). Additionally, dioxins (through the aryl hydrocarbon receptor), and endocrine disruptors found in diesel exhaust (including through the polycyclic aromatic hydrocarbon receptor) (Takeda et al. 2004) may influence the local hormonal milieu surrounding the lesion of endometriosis.

There are a few limitations of this study. We used ambient exposures at the residential address to estimate personal exposures. This could potentially lead to exposure misclassification in that we have no information on workplace exposures or the proportion of each day the woman spent at home, or on characteristics of the home (age, ventilation rate, air purification systems, etc.) that may influence the actual levels of ambient pollution exposure inside the home.

Restricting cases to those with laparoscopic confirmation may capture more “severe” cases only (Kennedy et al. 2005). However, among those with medical record review during our validation study (Missmer et al. 2004a), more than 60% of these women were stage I/II, i.e. minimal/mild disease at diagnosis. As mentioned previously, it is estimated that undiagnosed severe cases are <2% of the general population (Zondervan et al. 2002). Although it is plausible that there are undiagnosed cases in the cohort, their prevalence among the more than 70,000 comparison women is unlikely to explain our null results.

Although representative of the racial/ethnic and socioeconomic distribution of nurses at the time of cohort enrollment in 1989, the NHSII cohort is not representative of the general US population. The NHSII cohort comprises predominantly Caucasian women living in neighborhoods of medium to high socioeconomic status (SES). Another limitation is that we are only able to assess associations with exposures during adulthood. Specifically, the women were 25 to 43 years of age at enrollment in 1989, and were 43 to 61 by the end of follow-up in 2007. This window of exposure may not be at the most etiologically relevant time window in relation to endometriosis disease pathogenesis. Hence, this investigation does not confer any insight into earlier life exposure and endometriosis risk. As previously published from this cohort (Missmer et al. 2004b; Vitonis et al. 2010) and others (Kvaskoff et al. 2013; Treloar et al. 2010), exposures in earlier time windows, including intrauterine (diethylstilbestrol exposure, low birthweight) and

childhood exposures (lean childhood body size, earlier age at menarche), have been associated with endometriosis. As has been suggested by Woodruff and colleagues (Woodruff et al. 2008), it may be that exposures at these earlier and potentially more susceptible windows are the critical point of disease influence. Or it is possible that lifetime exposure information, including information on exposures during childhood, may be needed to elucidate the true relation between traffic and air pollution exposures and endometriosis.

This large nationwide prospective cohort of women also provided several important strengths. For this analysis, we had detailed biennially updated residential address history and only included residential addresses with a street segment level geocoding match. Our exposure model provided spatially and temporally resolved predictions over the course of follow-up, which likely limits exposure misclassification compared to a single annual estimate or predictions from a central monitoring location. The geographic distribution represented by the participants of this study provided information on regions throughout the continental US. Another strength was the prospective information on many potentially confounding covariates and effect modifiers.

Although our study did not support an association of adult exposures to PM with endometriosis risk, more research is needed on air pollution exposures and endometriosis.

References

- Abbas S, Ihle P, Koster I, Schubert I. 2012. Prevalence and incidence of diagnosed endometriosis and risk of endometriosis in patients with endometriosis-related symptoms: Findings from a statutory health insurance-based cohort in Germany. *European journal of obstetrics, gynecology, and reproductive biology* 160:79-83.
- Adar SD, Kaufman JD. 2007. Cardiovascular disease and air pollutants: Evaluating and improving epidemiological data implicating traffic exposure. *Inhal Toxicol* 19 Suppl 1:135-149.
- Agic A, Xu H, Finas D, Banz C, Diedrich K, Hornung D. 2006. Is endometriosis associated with systemic subclinical inflammation? *Gynecologic and obstetric investigation* 62:139-147.
- Andrade AZ, Rodrigues JK, Dib LA, Romao GS, Ferriani RA, Jordao Junior AA, et al. 2010. [serum markers of oxidative stress in infertile women with endometriosis]. *Rev Bras Ginecol Obstet* 32:279-285.
- Augoulea A, Alexandrou A, Creatsa M, Vrachnis N, Lambrinouadaki I. 2012. Pathogenesis of endometriosis: The role of genetics, inflammation and oxidative stress. *Arch Gynecol Obstet* 286:99-103.
- Breech LL, Laufer MR. 1999. Obstructive anomalies of the female reproductive tract. *J Reprod Med* 44:233-240.
- Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, et al. 2004. Air pollution and cardiovascular disease: A statement for healthcare professionals from the expert panel on population and prevention science of the American Heart Association. *Circulation* 109:2655-2671.
- Brook RD, Rajagopalan S, Pope CA, 3rd, Brook JR, Bhatnagar A, Diez-Roux AV, et al. 2010. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation* 121:2331-2378.
- Buck Louis GM, Hediger ML, Peterson CM, Croughan M, Sundaram R, Stanford J, et al. 2011. Incidence of endometriosis by study population and diagnostic method: The Endo Study. *Fertility and sterility* 96:360-365.

- Burney RO, Talbi S, Hamilton AE, Vo KC, Nyegaard M, Nezhat CR, et al. 2007. Gene expression analysis of endometrium reveals progesterone resistance and candidate susceptibility genes in women with endometriosis. *Endocrinology* 148:3814-3826.
- Calderón-Garcidueñas L, Engle R, Mora-Tiscareño A, Styner M, Gómez-Garza G, Zhu H, et al. 2011. Exposure to severe urban air pollution influences cognitive outcomes, brain volume and systemic inflammation in clinically healthy children. *Brain and Cognition* 77:345-355.
- Chen QH, Zhou WD, Su ZY, Huang QS, Jiang JN, Chen QX. 2010. Change of proinflammatory cytokines follows certain patterns after induction of endometriosis in a mouse model. *Fertility and sterility* 93:1448-1454.
- D'Hooghe TM, Bamba CS, Xiao L, Peixe K, Hill JA. 2001. Effect of menstruation and intrapelvic injection of endometrium on inflammatory parameters of peritoneal fluid in the baboon (*papio anubis* and *papio cynocephalus*). *American journal of obstetrics and gynecology* 184:917-925.
- D'Hooghe TM, Debrock S. 2002. Endometriosis, retrograde menstruation and peritoneal inflammation in women and in baboons. *Hum Reprod Update* 8:84-88.
- Draper LB, Chong H, Wang E, Woodruff TK. 1997. The uterine myometrium is a target for increased levels of activin a during pregnancy. *Endocrinology* 138:3042-3046.
- Fanton JW, Golden JG. 1991. Radiation-induced endometriosis in macaca mulatta. *Radiat Res* 126:141-146.
- Flores I, Rivera E, Ruiz LA, Santiago OI, Vernon MW, Appleyard CB. 2007. Molecular profiling of experimental endometriosis identified gene expression patterns in common with human disease. *Fertility and sterility* 87:1180-1199.
- Gentilini D, Perino A, Vigano P, Chiodo I, Cucinella G, Vignali M, et al. 2011. Gene expression profiling of peripheral blood mononuclear cells in endometriosis identifies genes altered in non-gynaecologic chronic inflammatory diseases. *Hum Reprod* 26:3109-3117.
- Greenland S. 1989. Modeling and variable selection in epidemiologic analysis. *Am J Public Health* 79:340-349.
- Grummer R. 2006. Animal models in endometriosis research. *Hum Reprod Update* 12:641-649.
- Halme J, Hammond MG, Hulka JF, Raj SG, Talbert LM. 1984. Retrograde menstruation in healthy women and in patients with endometriosis. *Obstetrics and gynecology* 64:151-154.

- Hart JE, Yanosky JD, Puett RC, Ryan L, Dockery DW, Smith TJ, et al. 2009. Spatial modeling of pm10 and no2 in the continental united states, 1985-2000. *Environmental health perspectives* 117:1690-1696.
- Houston DE, Noller KL, Melton LJ, 3rd, Selwyn BJ, Hardy RJ. 1987. Incidence of pelvic endometriosis in rochester, minnesota, 1970-1979. *American journal of epidemiology* 125:959-969.
- Hsu AL, Khachikyan I, Stratton P. 2010. Invasive and noninvasive methods for the diagnosis of endometriosis. *Clin Obstet Gynecol* 53:413-419.
- Karner AA, Eisinger DS, Niemeier DA. 2010. Near-roadway air quality: Synthesizing the findings from real-world data. *Environmental science & technology* 44:5334-5344.
- Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R, et al. 2005. Eshre guideline for the diagnosis and treatment of endometriosis. *Human reproduction* 20:2698-2704.
- Kruitwagen RF, Poels LG, Willemsen WN, de Ronde IJ, Jap PH, Rolland R. 1991. Endometrial epithelial cells in peritoneal fluid during the early follicular phase. *Fertility and sterility* 55:297-303.
- Kvaskoff M, Bijon A, Clavel-Chapelon F, Mesrine S, Boutron-Ruault MC. 2013. Childhood and adolescent exposures and the risk of endometriosis. *Epidemiology* 24:261-269.
- Lipfert FW, Baty JD, Miller JP, Wyzga RE. 2006. Pm2.5 constituents and related air quality variables as predictors of survival in a cohort of u.S. Military veterans. *Inhal Toxicol* 18:645-657.
- Lipfert FW, Wyzga RE. 2008. On exposure and response relationships for health effects associated with exposure to vehicular traffic. *J Expo Sci Environ Epidemiol* 18:588-599.
- Maybin JA, Critchley HO, Jabbour HN. 2011. Inflammatory pathways in endometrial disorders. *Mol Cell Endocrinol* 335:42-51.
- Misaki K, Suzuki M, Nakamura M, Handa H, Iida M, Kato T, et al. 2008. Aryl hydrocarbon receptor and estrogen receptor ligand activity of organic extracts from road dust and diesel exhaust particulates. *Arch Environ Contam Toxicol* 55:199-209.
- Missmer SA, Hankinson SE, Spiegelman D, Barbieri RL, Marshall LM, Hunter DJ. 2004a. Incidence of laparoscopically confirmed endometriosis by demographic, anthropometric, and lifestyle factors. *American journal of epidemiology* 160:784-796.

- Missmer SA, Hankinson SE, Spiegelman D, Barbieri RL, Michels KB, Hunter DJ. 2004b. In utero exposures and the incidence of endometriosis. *Fertility and sterility* 82:1501-1508.
- Nyholt DR, Low SK, Anderson CA, Painter JN, Uno S, Morris AP, et al. 2012. Genome-wide association meta-analysis identifies new endometriosis risk loci. *Nat Genet* 44:1355-1359.
- Oh SM, Ryu BT, Chung KH. 2008. Identification of estrogenic and antiestrogenic activities of respirable diesel exhaust particles by bioassay-directed fractionation. *Arch Pharm Res* 31:75-82.
- Pan Q, Luo X, Toloubeydokhti T, Chegini N. 2007. The expression profile of micro-rna in endometrium and endometriosis and the influence of ovarian steroids on their expression. *Molecular human reproduction* 13:797-806.
- Rier SE, Martin DC, Bowman RE, Dmowski WP, Becker JL. 1993. Endometriosis in rhesus monkeys (*macaca mulatta*) following chronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Fundam Appl Toxicol* 21:433-441.
- Sahlodin AM, Sotudeh-Gharebagh R, Zhu YF. 2007. Modeling of dispersion near roadways based on the vehicle-induced turbulence concept. *Atmos Environ* 41:92-102.
- Sampson J. 1927. Peritoneal endometriosis due to menstrual dissemination of endometrial tissue into the peritoneal cavity. *Am J Obstet Gynecol* 14:422-469.
- Schernhammer ES, Vitonis AF, Rich-Edwards J, Missmer SA. 2011. Rotating nightshift work and the risk of endometriosis in premenopausal women. *Am J Obstet Gynecol* 205:476 e471-478.
- Sidlova T, Novak J, Janosek J, Andel P, Giesy JP, Hilscherova K. 2009. Dioxin-like and endocrine disruptive activity of traffic-contaminated soil samples. *Arch Environ Contam Toxicol* 57:639-650.
- Siristatidis C, Nissotakis C, Chrelas C, Iacovidou H, Salamalekis E. 2006. Immunological factors and their role in the genesis and development of endometriosis. *J Obstet Gynaecol Res* 32:162-170.
- Splitter GA, Kirk JH, Mac Kenzie WF, Rawlings CA. 1972. Endometriosis in four irradiated rhesus monkeys. *Vet Pathol* 9:249-262.
- Takeda K, Tsukue N, Yoshida S. 2004. Endocrine-disrupting activity of chemicals in diesel exhaust and diesel exhaust particles. *Environ Sci* 11:33-45.

- Treloar SA, Bell TA, Nagle CM, Purdie DM, Green AC. 2010. Early menstrual characteristics associated with subsequent diagnosis of endometriosis. *American journal of obstetrics and gynecology* 202:534 e531-536.
- U.S.CensusBureau. 1992. Tiger products. Available: <http://www.census.gov/geo/maps-data/data/tiger.html> [accessed June 12 2005].
- Umezawa M, Sakata C, Tanaka N, Tabata M, Takeda K, Ihara T, et al. 2011. Pathological study for the effects of in utero and postnatal exposure to diesel exhaust on a rat endometriosis model. *J Toxicol Sci* 36:493-498.
- Van Langendonck A, Casanas-Roux F, Donnez J. 2002. Oxidative stress and peritoneal endometriosis. *Fertil Steril* 77:861-870.
- Vitonis AF, Baer HJ, Hankinson SE, Laufer MR, Missmer SA. 2010. A prospective study of body size during childhood and early adulthood and the incidence of endometriosis. *Human reproduction* 25:1325-1334.
- Wang J, Xie P, Kettrup A, Schramm KW. 2005. Inhibition of progesterone receptor activity in recombinant yeast by soot from fossil fuel combustion emissions and air particulate materials. *Sci Total Environ* 349:120-128.
- Wood DH, Yochmowitz MG, Salmon YL, Eason RL, Boster RA. 1983. Proton irradiation and endometriosis. *Aviat Space Environ Med* 54:718-724.
- Woodruff TJ, Carlson A, Schwartz JM, Giudice LC. 2008. Proceedings of the summit on environmental challenges to reproductive health and fertility: Executive summary. *Fertility and sterility* 89:281-300.
- Yanosky JD, Paciorek C, Schwartz J, Laden F, Puett R, Suh H. 2008a. Spatio-temporal modeling of chronic pm10 exposure for the nurses' health study. *Atmospheric Environment* doi:10.1016/j.atmosenv.2008.01.044.
- Yanosky JD, Schwartz J, Suh HH. 2008b. Associations between measures of socioeconomic position and chronic nitrogen dioxide exposure in worcester, massachusetts. *Journal of toxicology and environmental health Part A* 71:1593-1602.
- Yanosky JD, Paciorek CJ, Suh HH. 2009. Predicting chronic fine and coarse particulate exposures using spatiotemporal models for the northeastern and midwestern united states. *Environmental health perspectives* 117:522-529.

Zhu Y, Hinds WC, Kim S, Sioutas C. 2002. Concentration and size distribution of ultrafine particles near a major highway. *J Air Waste Manag Assoc* 52:1032-1042.

Zondervan KT, Cardon LR, Kennedy SH. 2002. What makes a good case-control study? Design issues for complex traits such as endometriosis. *Human reproduction* 17:1415-1423.

Table 1: Age-standardized characteristics^a of the cohort (N= 84,060) over the entire period of follow-up (1993-2007)

| Characteristics | Entire Cohort | By Distance (m) to Nearest A1-A3 Roadway ^b | | |
|--------------------------------------------|---------------|-------------------------------------------------------|-------------|-------------|
| | | 200+ m | 51-199 m | 0-50 m |
| Mean ± SD | | | | |
| Age (in years) ^c | 42.0 ± 5.1 | 42.1 ± 5.1 | 41.9 ± 5.2 | 42.0 ± 5.3 |
| Body Mass Index (kg/m ²) | 25.9 ± 6.0 | 25.8 ± 5.9 | 26.0 ± 6.1 | 26.4 ± 6.4 |
| Census tract median income (\$10,000) | 6.68 ± 2.38 | 6.75 ± 2.29 | 6.66 ± 2.47 | 6.33 ± 2.56 |
| Census tract median home value (\$100,000) | 1.72 ± 1.24 | 1.67 ± 1.08 | 1.84 ± 1.41 | 1.76 ± 1.57 |
| Caucasian race | 94 | 94 | 93 | 93 |
| Less than 12 | 23 | 23 | 23 | 23 |
| 12 | 30 | 30 | 30 | 31 |
| Greater than 12 | 46 | 47 | 46 | 46 |
| Nulliparous | 18 | 16 | 21 | 25 |
| Parous | 82 | 84 | 79 | 75 |
| Never | 14 | 13 | 16 | 15 |
| Past | 75 | 75 | 73 | 74 |
| Current | 10 | 10 | 10 | 10 |
| Never | 67 | 68 | 66 | 65 |
| Current | 9 | 8 | 10 | 10 |
| Former | 24 | 24 | 24 | 24 |
| Yes | 4 | 4 | 4 | 4 |
| No | 94 | 94 | 94 | 94 |
| Yes | 69 | 69 | 69 | 70 |
| No | 31 | 31 | 31 | 30 |
| Northeast | 34 | 32 | 37 | 42 |
| Midwest | 34 | 35 | 31 | 30 |
| West | 15 | 14 | 20 | 15 |
| South | 17 | 19 | 13 | 13 |

^a Values are standardized to the age distribution of the study population

^b Each cohort member may be in multiple distance categories over follow-up

^c Values are not age-adjusted

Table 2: Distributions of the particulate matter pollution metrics among 84,060 women in the Nurses' Health Study II

| Metric | Mean \pm SD | Median (IQR) | Min | Max |
|---------------------|----------------|--------------|-----|------|
| 2-year average | | | | |
| PM ₁₀ | 23.2 \pm 6.2 | 22.4 (6.9) | 3.7 | 72.3 |
| PM ₁₀₋₂₅ | 9.2 \pm 4.6 | 8.3 (5.2) | 0.0 | 54.0 |
| PM _{2.5} | 14.0 \pm 3.0 | 14.0 (4.1) | 2.0 | 28.2 |
| 4-year average | | | | |
| PM ₁₀ | 21.7 \pm 5.5 | 21.0 (6.2) | 4.0 | 60.1 |
| PM ₁₀₋₂₅ | 8.4 \pm 4.2 | 7.4 (4.9) | 0.1 | 46.3 |
| PM _{2.5} | 13.3 \pm 2.7 | 13.3 (3.7) | 2.2 | 24.1 |
| Cumulative average | | | | |
| PM ₁₀ | 26.2 \pm 6.5 | 25.4 (6.9) | 5.1 | 80.7 |
| PM ₁₀₋₂₅ | 10.9 \pm 4.9 | 9.8 (5.4) | 1.7 | 60.7 |
| PM _{2.5} | 15.4 \pm 3.1 | 15.5 (4.2) | 2.7 | 29.3 |

Table 3: Basic and fully adjusted hazard ratios and 95% confidence intervals (CIs) of endometriosis risk by residential proximity to roadway, among 84,060 women in the Nurses' Health Study II in the whole country and by region

| Exposure | Person-years | Cases | Whole Country | Adjusted HR 95% CI ^b | Northeast | Midwest | West | South |
|-----------------------------|-----------------------|-------|---------------------------------|------------------------------------|------------------------------------|------------------------------------|------------------------------------|------------------------------------|
| | | | (2,486 cases) | | (761 cases) | (898 cases) | (334 cases) | (493 cases) |
| | | | Basic HR 95% CI ^a | | Adjusted HR 95% CI ^b |
| Distance to A1-A3 Roads (m) | | | | | | | | |
| 0-50 | 82,982 | 308 | 1.08 (0.95, 1.22) | 1.04 (0.91, 1.17) | 1.10 (0.89, 1.35) | 0.98 (0.78, 1.22) | 0.96 (0.66, 1.39) | 1.06 (0.77, 1.46) |
| 51-199 | 182,567 | 700 | 1.11 (1.01, 1.22) | 1.09 (0.99, 1.19) | 1.07 (0.90, 1.27) | 1.07 (0.91, 1.25) | 0.99 (0.78, 1.28) | 1.28 (1.03, 1.59) |
| 200+ | 440,204 | 1,478 | reference | reference | reference | reference | reference | reference |
| Distance to A1-A3 Roads (m) | | | | | | | | |
| 0-50 | 82,982 | 308 | 1.11 (0.97, 1.27) | 1.07 (0.93, 1.23) | 1.22 (0.96, 1.55) | 1.00 (0.78, 1.27) | 0.98 (0.63, 1.51) | 0.98 (0.70, 1.37) |
| 51-499 | 410,164 | 1,488 | 1.10 (1.00, 1.20) | 1.09 (0.99, 1.19) | 1.20 (1.00, 1.43) | 1.07 (0.92, 1.24) | 1.03 (0.77, 1.38) | 0.99 (0.81, 1.20) |
| 500+ | 212,607 | 690 | reference | reference | reference | reference | reference | reference |
| Distance to A1-A2 Roads (m) | | | | | | | | |
| 0-50 | 8,889 | 36 | 1.11 (0.79, 1.56) | 1.03 (0.73, 1.44) | 1.08 (0.63, 1.84) | 0.74 (0.40, 1.40) | 0.69 (0.16, 2.91) | 1.68 (0.81, 3.48) |
| 51-199 | 36,796 | 119 | 0.91 (0.75, 1.10) | 0.86 (0.71, 1.04) | 0.73 (0.53, 1.00) | 0.79 (0.55, 1.13) | 1.22 (0.77, 1.93) | 1.23 (0.79, 1.91) |
| 200+ | 660,068 ^{bv} | 2,331 | reference | reference | reference | reference | reference | reference |

^a Adjusted for age and calendar time

^b Additionally adjusted for race, current BMI, smoking status, parity, OC use, age at menarche, infertility, ever performed rotating shift work, and Census tract level median income and median home value and region (in un-stratified models)

Table 4: Basic and fully adjusted hazard ratios and 95% confidence intervals (CIs) of endometriosis risk for each 10- $\mu\text{g}/\text{m}^3$ increase in particulate matter, among 84,060 women in the Nurses' Health Study II in the whole country and by region of residence

| Exposure Averaging Time | Whole country (2,486 cases) | | Northeast (761 cases) | Midwest (898 cases) | West (334 cases) | South (493 cases) |
|-------------------------------|---------------------------------|-----------------------------------------------------|-----------------------------------------------------|-----------------------------------------------------|-----------------------------------------------------|-----------------------------------------------------|
| | Basic HR 95% CI ^a | Multivariable Adjusted HR 95% CI ^b |
| 2-year | | | | | | |
| PM ₁₀ | 0.98 (0.92, 1.05) | 0.94 (0.87, 1.02) | 1.01 (0.84, 1.21) | 0.96 (0.82, 1.11) | 0.91 (0.80, 1.03) | 0.82 (0.61, 1.09) |
| PM _{10-2.5} | 0.98 (0.89, 1.07) | 0.91 (0.81, 1.02) | 0.88 (0.62, 1.24) | 0.99 (0.81, 1.20) | 0.86 (0.69, 1.05) | 0.76 (0.54, 1.07) |
| PM _{2.5} | 0.98 (0.85, 1.12) | 0.95 (0.83, 1.10) | 1.18 (0.84, 1.65) | 0.88 (0.67, 1.16) | 0.85 (0.66, 1.11) | 1.00 (0.72, 1.37) |
| 4-year | | | | | | |
| PM ₁₀ | 0.97 (0.90, 1.05) | 0.92 (0.84, 1.00) | 0.98 (0.80, 1.20) | 0.90 (0.76, 1.06) | 0.92 (0.80, 1.06) | 0.78 (0.55, 1.11) |
| PM _{10-2.5} | 0.97 (0.88, 1.07) | 0.88 (0.77, 1.00) | 0.88 (0.62, 1.24) | 0.99 (0.81, 1.20) | 0.86 (0.69, 1.05) | 0.76 (0.54, 1.07) |
| PM _{2.5} | 0.97 (0.83, 1.13) | 0.92 (0.79, 1.07) | 1.14 (0.80, 1.63) | 0.80 (0.59, 1.09) | 0.86 (0.65, 1.13) | 1.08 (0.75, 1.55) |
| Cumulative | | | | | | |
| PM ₁₀ | 0.97 (0.91, 1.04) | 0.93 (0.87, 1.01) | 1.02 (0.86, 1.21) | 0.95 (0.82, 1.10) | 0.91 (0.80, 1.02) | 0.78 (0.61, 1.00) |
| PM _{10-2.5} | 0.98 (0.91, 1.07) | 0.92 (0.82, 1.02) | 0.90 (0.65, 1.24) | 0.99 (0.82, 1.19) | 0.89 (0.73, 1.08) | 0.72 (0.52, 1.00) |
| PM _{2.5} | 0.93 (0.81, 1.05) | 0.91 (0.80, 1.04) | 1.19 (0.87, 1.62) | 0.87 (0.68, 1.12) | 0.79 (0.61, 1.02) | 0.92 (0.69, 1.23) |

^a Adjusted for age and calendar time

^b Additionally adjusted for race, current BMI, smoking status, parity, OC use, age at menarche, infertility, ever performed rotating shift work, and Census tract level median income and median home value and region (in un-stratified models)

Figure Legends

Figure 1: Geographic distribution of nurses' addresses from 1989 – 2007.

Figure 2: Examination of possible effect modification of a 10-ug/m³ increase in particulate matter and risk of incident endometriosis by (top left to bottom right): parity, smoking status, age at menarche, BMI, infertility, and rotating shift work. All models are adjusted for age and calendar time, race, current BMI, smoking status, parity, oral contraceptive use, age at menarche, infertility, ever performed rotating shift work, region, and Census tract level median income and median home value as appropriate.




Figure 1.




Figure 2.