



ENVIRONMENTAL
HEALTH
PERSPECTIVES

<http://www.ehponline.org>

**Prenatal Exposure to Environmental Phenols:
Concentrations in Amniotic Fluid and Variability in
Urinary Concentrations during Pregnancy**

**Claire Philippat, Mary S. Wolff, Antonia M. Calafat, Xiaoyun Ye,
Rebecca Bausell, Molly Meadows, Joanne Stone, Rémy Slama
and Stephanie M. Engel**

<http://dx.doi.org/10.1289/ehp.1206335>

Received: 30 November 2012

Accepted: 9 August 2013

Advance Publication: 13 August 2013



Prenatal Exposure to Environmental Phenols: Concentrations in Amniotic Fluid and Variability in Urinary Concentrations during Pregnancy

Claire Philippat,^{1,2} Mary S. Wolff,³ Antonia M. Calafat,⁴ Xiaoyun Ye,⁴ Rebecca Bausell,⁵ Molly Meadows,⁵ Joanne Stone,⁶ Rémy Slama,^{1,2} and Stephanie M. Engel⁷

1: Inserm, Institut Albert Bonniot (U823), Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, Grenoble, France.

2: Grenoble University, Institut Albert Bonniot, Grenoble, France.

3: Department of Preventive Medicine, Mount Sinai School of Medicine, New York, New York, USA.

4: Centers for Disease Control and Prevention, Atlanta, Georgia, USA.

5: Columbia University College of Physicians and Surgeons, New York, New York, USA.

6: Department of Obstetrics, Gynecology and Reproductive Sciences, Mount Sinai School of Medicine, New York, New York, USA.

7: Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA.

Address correspondance to:

Claire Philippat

Institut Albert Bonniot - Centre de Recherche INSERM-UJF U823, UJF Site Santé

BP 170 - La Tronche - Grenoble 38042 Cedex 9 France

claire.philippat@e.ujf-grenoble.fr

Phone: +33 4 76 54 94 66

Fax: +33 4 76 54 94 14

Running Title: Phenols in amniotic fluid and maternal urine -

Acknowledgments: We acknowledge Larry Engel for his helpful comments, and Xiaoliu Zhou, -
Joshua Kramer, and Tao Jia for measuring the concentrations of the phenols. -

Funding: This research was supported by funding from NIEHS (R21ES014034). Ms. Philippat -
was supported by a grant from Region Rhône-Alpes. -

Competing financial interest: None -

Disclaimer: The findings and conclusions in this report are those of the authors and do not -
necessarily represent the official position of the Centers for Disease Control and Prevention. -

Abstract

Background: Maternal urinary biomarkers are often used to assess fetal exposure to phenols and their precursors. Their effectiveness as a measure of exposure in epidemiological studies depends on their variability during pregnancy and their ability to accurately predict fetal exposure.

Objectives: We assessed the relationship between urinary and amniotic fluid concentrations of nine environmental phenols, and the reproducibility of urinary concentrations, among pregnant women.

Methods: Seventy-one women presenting for amniocentesis were included. Maternal urine was collected at the time of the amniocentesis appointment and on two subsequent occasions. Urine and amniotic fluid were analyzed for 2,4- and 2,5-dichlorophenols, bisphenol A, benzophenone-3, triclosan, and methyl-, ethyl-, propyl-, and butyl-parabens using online solid phase extraction–high performance liquid chromatography–isotope dilution tandem mass spectrometry.

Results: Only benzophenone-3 and propylparaben were detectable in more than half of the amniotic fluid samples; for these phenols, concentrations in amniotic fluid and maternal urine collected on the same day were positively correlated ($\rho = 0.53$ and 0.32 , respectively). Other phenols were detected infrequently in amniotic fluid (e.g., bisphenol A was detected in only two samples). The intraclass correlation coefficients (ICC) of urinary concentrations in samples from individual women ranged from 0.48 and 0.62 for all phenols except bisphenol A (ICC = 0.11).

Discussion: Amniotic fluid detection frequencies for most phenols were low. The reproducibility of urine measures was poor for bisphenol A, but good for the other phenols. Although a single sample may provide a reasonable estimate of exposure for some phenols, collecting multiple urine samples during pregnancy is an option to reduce exposure measurement error in studies regarding the effects of phenol prenatal exposure on health.

Background

Phenolic compounds are used in numerous consumer products such as cosmetics (parabens), sunscreens (benzophenone-3), antibacterial soaps (triclosan), food packaging and polycarbonate plastics (bisphenol A). 2,5-dichlorophenol is a metabolite of p-dichlorobenzene, a chemical used in moth balls and bathroom deodorizers. 2,4-dichlorophenol is primarily used in the production of phenoxy acid herbicides (e.g., 2,4-diphenoxyacetic acid) and for the synthesis of pharmaceuticals and antiseptics. Exposure to these chemicals or their precursors is ubiquitous in the general population, including pregnant women (Casas et al. 2013; Wolff et al. 2008). Metabolites of phenols have also been detected in amniotic fluid (Edlow et al. 2012; Engel et al. 2006; Yamada et al. 2002) and cord blood (Chou et al. 2011; Fenichel et al. 2012), suggesting fetal exposure.

Certain environmental chemicals (e.g., phenols) are rapidly metabolized and excreted in urine after exposure. Thus, urinary concentrations are commonly used in epidemiological studies to estimate exposure to these chemicals. The extent to which concentrations in maternal urine reflect fetal exposure is uncertain. Alternative matrices, such as amniotic fluid, might provide a more accurate measure of direct fetal exposure. During pregnancy, the fetus swallows amniotic fluid and fetal urine is one of the main sources of amniotic fluid in the second half of pregnancy (Modena and Fieni 2004; Underwood et al. 2005). Thus, concentrations measured in this matrix might be a relevant dosimeter of fetal exposure. In humans, bisphenol A (Edlow et al. 2012; Engel et al. 2006; Ikezuki et al. 2002; Yamada et al. 2002) and 2,5-dichlorophenol (Bradman et al. 2003) have been detected in amniotic fluid collected from pregnant women receiving amniocentesis as part of their routine prenatal care. However, to our knowledge, no published study has evaluated human amniotic fluid concentrations of other prevalent environmental

phenols, nor has there been a comparison of phenol biomarker concentrations in amniotic fluid with corresponding concentrations in maternal urine.

There have been few studies of the health effects of prenatal exposures to phenols or their precursors, and all have relied on a relatively small number of maternal urine samples collected during pregnancy (1 to 3) to assess exposures (Philippat et al. 2012; Wolff et al. 2008). However, many phenols have short half-lives (less than 1 day) in humans (Janjua et al. 2008; Teeguarden et al. 2011; Volkel et al. 2002), and exposures to these compounds are closely related to dietary intakes and episodic personal behaviors such as cosmetic use. Therefore, urinary concentrations are likely to vary throughout pregnancy. Variation in sample collection and processing may be an additional source of variability in phenol biomarker concentrations, and a statistical method to standardize measured concentrations for such factors (Mortamais et al. 2012) has been applied in epidemiological studies of associations between urinary biomarkers and health endpoints (Chevrier et al. 2012; Philippat et al. 2012). However, to our knowledge, the impact of standardizing biomarker concentrations on reproducibility measures has not been studied.

Our objectives were to assess the relationship between phenol biomarker concentrations measured in maternal urine and amniotic fluid collected on the same day, and to study the temporal variability in urinary phenol concentrations over the course of pregnancy. A secondary aim was to assess the impact of collection conditions on the measured urine phenol concentrations, and determine whether standardization for these conditions improves reliability measures in phenol concentrations during pregnancy.

Methods

Study population

The Study of Advanced Reproductive Age and Environmental Health (SARAEH) recruited 97 women who were referred for amniocentesis screening at the Mount Sinai Medical Center in New York, NY. Recruitment occurred between 2005 and 2008. Study eligibility requirements were fluency in English or Spanish, maternal age between 18 and 40 years, singleton pregnancy, and intent to deliver at Mount Sinai Hospital. Women were approached for participation in the waiting area prior to their amniocentesis. Those who consented to participate in the study provided a spot urine sample immediately following their procedure. Two additional spot urine samples were collected from each participant later in their pregnancy, with collections planned at least two weeks apart, and the final sample collected after the 27th week of gestation. The present study comprises 71 participants who provided all three urine samples, including 69 for whom an amniotic fluid sample also was available. Following delivery, medical records were abstracted for information pertaining to fetal health.

This study was approved by the Mount Sinai Program for the Protection of Human Subjects and all women gave informed consent to be part of the study. The involvement of the Centers for Disease Control and Prevention (CDC) laboratory was determined not to constitute engagement in human subjects research.

Exposure assessment

Following collection, amniotic fluid was delivered to the Mount Sinai Medical Center clinical cytogenetics laboratory for standard clinical care processing, which includes centrifugation and cell separation. Before being released for research purposes, amniotic fluid supernatant was

stored in polypropylene containers at -20°C (time range: 0 to 38 weeks). Once released, SARAHEH research personnel thawed the amniotic fluid overnight at 4°C, and then aliquoted the supernatant into 2 mL cryovials for storage at -80°C. Following collection, urine was immediately aliquoted into 2 mL cryovials and stored at -80°C. Urine and amniotic fluid samples were shipped overnight on dry ice to the CDC for measurements of total (free plus conjugated) concentrations of 2,4- and 2,5-dichlorophenols, bisphenol A, benzophenone-3, triclosan, and methyl-, ethyl-, propyl-, and butyl-parabens by using online solid phase extraction–high performance liquid chromatography–isotope dilution tandem mass spectrometry (Ye et al. 2005). Specific gravity of urine was measured using a handheld refractometer in each thawed aliquot prior to shipment to the CDC, with the exception of 11 urine samples that had specific gravity measured using an Atago PAL-10S refractometer (Bellevue, WA, USA) at CDC. Urinary creatinine concentration was measured at CDC by an enzymatic reaction using a Roche Hitachi 912 Chemistry Analyzer (Hitachi, Pleasanton, CA, USA).

Statistical analyses

Analyses were conducted on natural-log-transformed concentrations (ln). Amniotic fluid concentrations below the limit of detection (LOD) were replaced with the instrument reading values for phenols detected in at least 50% of the samples. Instrument reading values were not available for urine assays; therefore, urine phenol concentrations below the LOD were replaced with the $LOD/\sqrt{2}$. The following formula was used to correct urinary concentrations for specific gravity: $C_{SG} = C \times [(SG_{mean} - 1)/(SG - 1)]$, where C_{SG} is the specific gravity corrected biomarker concentration, SG_{mean} is the specific gravity arithmetic mean in our population, and C is the measured biomarker concentration. Creatinine corrected concentrations were calculated by dividing the phenol concentrations by the creatinine concentration.

Relation between maternal urine and amniotic fluid biomarker concentrations

We compared amniotic fluid and urinary biomarker concentrations collected on the same day, and computed ratios of uncorrected urine to amniotic fluid concentrations. For benzophenone-3 and propylparaben, the two phenols detected in more than 50% of the amniotic fluid samples, we computed Spearman correlation coefficients comparing concentrations in amniotic fluid and urinary samples (uncorrected and specific gravity corrected) collected on the same day. To explore the possible predictors of benzophenone-3 and propylparaben concentrations in amniotic fluid, we performed Tobit regression models for a left-censored dependent variable (Lubin et al. 2004). We regressed amniotic fluid biomarker concentrations on maternal urinary concentrations in samples collected on the same day, fetal sex, gestational age at amniocentesis (< 17, 17 to 18.9, ≥ 19 gestational weeks), maternal age (< 31, 31 to 35.9, ≥ 36), maternal pre-pregnancy body mass index (BMI, < 25, ≥ 25), race/ethnicity (white non-Hispanic or other), pregnancy complications related to placental function (including preeclampsia, placenta previa, small placenta, oligohydroamnios and chorioamnionitis), and the time between amniotic fluid collection and processing (≤ 4 , 4 to 16, 16 to 20, or > 20 weeks). Models were simultaneously adjusted for all of these factors. We also performed sensitivity analyses excluding four women with abnormal amniotic fluid conditions (oligohydramnios or polyhydramnios).

Variability in urinary concentrations

We evaluated variability in measures of dilution in spot urine samples across pregnancy by plotting urinary creatinine concentrations and urine specific gravity as a function of gestational age. To assess variability in phenol concentrations across pregnancy, we computed Intraclass Correlation Coefficients (ICCs) between concentrations measured in the three spot urine samples using random intercept linear mixed models (Rabe-Hesketh 2008). The ICC is the ratio of the

between-woman variability to the total variability (between- plus within-woman variability). We also computed Spearman correlations between pairs of phenol concentrations measured in the three spot urine samples from each woman. We classified the comparability of samples based on ICCs and Spearman correlation coefficients according to the following general guidelines: < 0.4 : weak, 0.4 to 0.6 : moderate, > 0.6 : good. Analyses of urinary concentration variability were performed using uncorrected, specific gravity-corrected, and creatinine-corrected concentrations. Creatinine concentrations were missing for 11 of the first spot urine samples. Thus, analyses based on creatinine-corrected concentrations were restricted to 60 women with complete creatinine and environmental phenol biomarker data.

In a secondary analysis, we computed Spearman correlations among pairs of urine samples collected within specific time intervals: fewer than 4 weeks apart, 4-6 weeks apart, 6-10 weeks apart, 10-12 weeks apart, 12-14 weeks apart, 14-16 weeks apart, and greater than 16 weeks apart. Samples were not independent within a given category because multiple samples from an individual woman could be included in a single category if they were collected at equally spaced intervals.

Finally, we estimated associations between urinary phenol biomarker concentrations and collection conditions (gestational age at collection; hour, day and season of sampling; urine specific gravity) using a random intercept linear mixed model simultaneously adjusted for each collection condition along with BMI, maternal age, year of collection, maternal education, and maternal race/ethnicity (modeled as indicated previously) (Mortamais et al. 2012). We used the measured urinary biomarker concentrations and the estimated effects of collection conditions on the measured urine concentrations (for conditions that predicted urine concentrations with $p < 0.02$) to derive standardized concentrations, i.e., concentrations that would have been observed if all

samples had been collected under the same conditions (Mortamais et al. 2012). We estimated ICCs based on concentrations standardized for collection conditions to determine whether standardization improved reliability across repeated spot urine samples.

All analyses were performed using STATA/SE, version 12 (StataCorp, College Station, TX, USA).

Results

Study population and exposure

Women were 35.5 years old on average, mostly white (72%), and 55% attended graduate school (Table 1). The three urine samples were collected on average (\pm SD), at 17.6 ± 1.6 , 22.8 ± 3.5 , and 32.8 ± 3.1 gestational weeks, respectively. On average, the first two urine samples were collected 5 weeks apart (± 2.8 weeks) and the second and third were collected 10 weeks apart (± 3.9 weeks). Overall, the first urine sample was collected later in the day than the other samples, and the number of women with urine samples collected before 10 a.m. was 1, 15, and 16 for the first, second, and third samples, respectively. With the exception of ethylparaben, median values of measured phenol concentrations (uncorrected for urine dilution), creatinine concentrations and specific gravity- were lowest in the first urine samples (Table 2). Detection frequencies of environmental phenols in urine were high: 2,5-dichlorophenol, benzophenone-3, methyl-paraben, and propyl-paraben were detected in at least 99% of the samples, and detection frequencies for the other phenols ranged between 59 and 94% (Table 2). Detection frequencies in amniotic fluid were much lower: only benzophenone-3 and propylparaben were detectable in at least half of the samples (61% and 58%, respectively). Bisphenol A was detected in two amniotic fluid samples

only (at a concentration equal to the LOD of 0.4 µg/L), but was detected in 62% of the urine samples collected on the same day (Table 2).

Relation between maternal urine and amniotic fluid biomarker concentrations

Among women with no abnormal amniotic fluid condition (i.e. excluding four participants with oligohydramnios or polyhydramnios), ratios of concentrations in the first urine sample (not corrected for urine dilution) to concentrations in amniotic fluid were greater than 1 (Table 2). However, benzophenone-3 and butylparaben concentrations were higher in amniotic fluid than in the corresponding urine sample for one woman with oligohydramnios (urine to amniotic fluid ratios of 0.2 for benzophenone-3 and 0.7 for butylparaben).

With the exception of butylparaben, which was detected in only four amniotic fluid samples, median urinary concentrations of phenol biomarkers (with or without correction for dilution) were higher in women with detectable amniotic fluid concentrations (Figure 1). The correlation between amniotic fluid and urinary concentrations (uncorrected for urine dilution) was moderate for benzophenone-3 ($\rho = 0.53$) and weak for propylparaben ($\rho = 0.32$). Correction for urinary dilution had little effect on these correlations (data not shown).

After adjustment (for storage time, gestational age at sampling, maternal age, BMI, race/ethnicity, pregnancy complications, and child's sex) ln-amniotic fluid concentrations increased by 0.52 (95% CI: 0.3, 0.7) and 0.36 (95% CI: 0.2, 0.5) with a one-unit increase in the ln-transformed specific-gravity-corrected urinary concentration of benzophenone-3 and propylparaben, respectively (Table 3). Including urine specific gravity as a covariate instead of using phenol concentrations corrected for specific gravity did not change the results (data not shown). After exclusion of the four women with abnormal amniotic fluid conditions

(oligohydramnios and polyhydramnios) urine and amniotic fluid biomarker concentrations were still positively associated (data not shown). There was a limited range of gestational ages, and no clear associations between gestational age at urine collection and amniotic fluid concentrations. Maternal age (p for trend = 0.09) and being non-white (p -value = 0.16) were negatively associated with amniotic concentrations of benzophenone-3. Propylparaben and benzophenone-3 concentrations in amniotic fluid were lower for girl fetuses than boy fetuses (β = -0.56; 95% CI: -1.2, 0.1 and β = -0.37; 95% CI: -1.2, 0.4, respectively) (Table 3).

Variability in urinary concentrations

We observed similar variations in urinary specific gravity and creatinine concentrations throughout pregnancy (Supplemental Material, Figure S1), and both measures of urinary dilution were highly correlated (spearman correlation coefficient = 0.93). ICCs (Table 4) and Spearman correlations (Supplemental Material, Table S1) were similar for urine concentrations corrected for creatinine or specific-gravity and, with the exception of ethylparaben and bisphenol A, were always higher than corresponding estimates for concentrations that were not corrected for dilution. Given that, and because creatinine concentrations are missing for 11 urine samples, we discuss only specific gravity-corrected estimates going forward.

ICCs were moderate (0.4–0.6) to good (> 0.6), and of comparable magnitude, for 2,4-, 2,5-dichlorophenol, benzophenone-3, triclosan, and methyl-, ethyl- propyl-, and butyl-parabens (Table 4), with similar patterns for Spearman correlation coefficients, which ranged from 0.41 to 0.77 (Supplemental Material, Table S1). In contrast, the ICC for bisphenol A was only 0.11 (Table 4). Although collections of spot urines were planned to occur at least two weeks apart,

four women had one interval of less than two weeks between successive collections. However, excluding these women did not change our results (data not shown).

Correlations between urinary concentrations measured within different time intervals are shown in Supplemental Material, Table S2. Our expectation was that correlations would be higher for samples collected with shorter intervening time intervals, and longer for samples collected with longer intervening time intervals. As expected, for 2,4-, 2,5-dichlorophenol, ethylparaben and benzophenone-3, but not for the other phenols, we overall observed lower coefficients (i.e., less reliability) for samples with longer time intervals between collections. Correlations for bisphenol A were weak, regardless of the time between sample collections.

Specific gravity was positively associated with all urinary phenol biomarker concentrations (p -values ≤ 0.1) (Supplemental Material, Table S3). Similar associations were observed with creatinine concentrations (data not shown). We observed statistically significant ($p < 0.05$) heterogeneity in 2,4-dichlorophenol, benzophenone-3, triclosan and butylparaben concentrations according to hour of sampling and in bisphenol A concentration according to gestational age. Benzophenone-3 concentrations were significantly higher in samples collected in spring and summer (adjusted geometric mean: 110 $\mu\text{g/L}$) than in samples collected in fall and winter (adjusted geometric mean: 69 $\mu\text{g/L}$). Race/ethnicity was associated with benzophenone-3 concentration in urine, which was higher among white non-Hispanic compared to non-white women ($\beta = -1.2$, 95%CI, -2.1; -0.3). Urinary concentrations standardized for collection conditions (Mortamais et al. 2012) and urinary concentrations corrected for specific gravity were strongly correlated (Spearman correlation coefficients > 0.96). The use of concentrations standardized for collection conditions did not substantially alter ICCs for spot urine collections from ICCs for concentrations that were uncorrected or corrected for dilution (Table 4).

Discussion

In our population of pregnant women referred for amniocentesis, the detection frequencies of phenol biomarkers in amniotic fluid were low, with only two phenols detected in 50% or more of the samples. With the exception of bisphenol A (ICC = 0.11), the reproducibility of urine phenol biomarker concentrations in urine collected at 3 times during pregnancy was fairly good (ICCs of 0.48 – 0.62 for specific gravity corrected concentrations).

Phenol concentrations in maternal urine

Urine collected on the day of the amniocentesis tended to be more dilute than urine collected later in pregnancy; however no specific instructions (with respect to hydration) or procedures (e.g., use of intravenous therapy) that would impact urinary dilution were given before or during the amniocentesis. Methyl- and propyl- parabens are the most commonly used parabens in food and cosmetics (Soni et al. 2005), and as expected, urine concentrations of these phenols were higher than those observed for ethyl- and butyl- parabens. Triclosan and bisphenol A concentrations were similar, and 2,5-dichlorophenol concentrations were lower than those reported in a previous mother-child cohort study conducted at the Mount Sinai Medical Center between 1998 and 2002, which included predominantly Hispanic and African American participants (medians were 11 µg/L for triclosan, 1.3 µg/L for bisphenol A, and 53 µg/L for 2,5-dichlorophenol (Wolff et al. 2008), compared with medians of 6.5–16.2, 0.6–1.8, and 7.5–14.2 µg/L, respectively, in our study population). Urinary concentrations of parabens were not reported in the study published by Wolff et al. Median urinary concentrations of benzophenone-3 in our study population (54–77 µg/L) were considerably higher than those reported in the aforementioned study (median of 7.5 µg/L). It is likely that this partly reflects the different

racial/ethnic composition of these two Mount Sinai studies, because exposure to benzophenone-3 in sunscreens (Summers et al. 2011) would be more common in the predominantly white participants. In our study being white non-Hispanic has been associated with higher benzophenone-3 concentrations in urine and in amniotic fluid.

Phenol concentrations in amniotic fluid

While the presence of phenol biomarkers in amniotic fluid highlights the potential for direct fetal exposure, the low detection frequency observed for the majority of these compounds may be suggestive of low transfer and/or rapid clearance of the phenols in amniotic fluid. That being said, we cannot exclude the possibility that the low detection of phenol biomarkers in amniotic fluid was the result of processing and/or storage of samples in the clinical laboratory prior to being released for research purposes. Ye et al. reported good stability of total phenol concentrations in urine stored at -70°C (Ye et al. 2007), but, to our knowledge, there are no published data regarding the impact of storage at -20°C or the stability of biomarker concentrations in amniotic fluid.

We detected bisphenol A in only one amniotic fluid sample. Low concentrations of bisphenol A in amniotic fluid collected during pregnancy and at birth have been previously reported (medians ranged from 0.26 to 1.1 $\mu\text{g/L}$) (Edlow et al. 2012; Engel et al. 2006; Ikezuki et al. 2002; Yamada et al. 2002). In an ex-vivo experimental system, where 7 human placentas were in contact with a perfusate enriched in bisphenol A (10 $\mu\text{g/L}$), authors observed rapid transfer of this phenol across the placenta (Balakrishnan et al. 2010); however, the concentration used in the experimental study was higher than those typically observed in the urine of pregnant women from the general population (Woodruff et al. 2011). Bradman et al. (2003) reported higher

geometric mean of 2,5-dichlorophenol concentrations (0.39 $\mu\text{g/L}$) in second trimester amniotic fluid samples from 20 women referred for amniocentesis at a Children's Hospital Central California in Madera (California) than we observed in our population (geometric mean = 0.29 $\mu\text{g/L}$). To our knowledge, no previous study has examined amniotic fluid concentrations of benzophenone-3, triclosan, or parabens in humans.

Predictors of amniotic fluid concentrations

Women with the highest concentrations of phenols in urine were more likely to have detectable concentrations of environmental phenols in amniotic fluid. Because of the high percentage of undetectable concentrations, we were only able to evaluate predictors of amniotic fluid concentrations for benzophenone-3 and propylparaben, and for these substances, maternal urine concentration was found to be a strong predictor of amniotic fluid concentration. There have been few comparisons of maternal and fetal exposure for phenolic compounds (Ikezuki et al. 2002; Yamada et al. 2002). Ikezuki et al. (2002) compared bisphenol A concentration in maternal serum (n = 37 samples) and amniotic fluid (n = 32 samples collected between 15–18 weeks), and reported amniotic fluid biomarker concentrations (arithmetic mean 8.3 $\mu\text{g/L}$) about 5 times higher than in maternal serum (arithmetic mean 1.5 $\mu\text{g/L}$). However mean concentration of 8.3 $\mu\text{g/L}$ is substantially higher than amniotic fluid concentrations reported by others (median values from 0.26 to 1.1 $\mu\text{g/L}$) (Edlow et al. 2012; Engel et al. 2006; Yamada et al. 2002), raising concerns regarding potential external contamination from environmental sources of free bisphenol A.

We found no evidence of an association between gestational age at collection and benzophenone-3 or propylparaben amniotic fluid concentrations. However, the range of

gestational ages at amniocentesis (15 to 23 gestational weeks) was limited in our study population. Two previous studies reported higher bisphenol A concentrations in amniotic fluid collected in the second trimester than in samples collected later pregnancy. However, in these studies, the samples collected in the second and third trimesters were from different women who may have had different levels of exposure (Edlow et al. 2012; Ikezuki et al. 2002). In our study population, the concentration of propylparaben in amniotic fluid was lower in female fetuses compared with male fetuses. This difference may have been due to chance, or could indicate sex differences in fetal metabolism or excretion.

Variability in urinary concentrations

The poor reproducibility of bisphenol A concentrations measured in serial spot urine samples collected during pregnancy is consistent with previous reports (Braun et al. 2011; Braun et al. 2012) and taken together, these findings suggest that substantial exposure misclassification might occur in epidemiological studies when prenatal bisphenol A exposure is classified using a single spot urine sample. Exposures to the eight other phenols or their precursors, which are found in sunscreens, cosmetics and indoor deodorizers were more consistent throughout pregnancy in our study population and had a higher degree of reproducibility in urinary concentrations, with ICCs between 0.5 and 0.6. Although a correlation of this magnitude is greater than reported for many other rapidly metabolized compounds, it may still result in bias due to exposure misclassification if the putative window of susceptibility is distant from the sample collection period. In a study relying on repeated measurements in indoor dust, authors reported a 40% attenuation of the effect estimates when only one measurement per household, rather than the mean of the repeated measurements per household, was used to assess exposure to chemicals with ICC of about 0.6 (Whitehead et al. 2012). In order to limit the effects of exposure misclassification, studies of the

effects of phenol prenatal exposure on health should try to collect multiple urine samples during pregnancy.

Two previous studies among pregnant women reported ICCs for methyl-, propyl- and butylparabens (ICCs < 0.5) that were low relative to ICCs in our population; the authors did not measure ethylparaben concentration (Meeker et al. 2013; Smith et al. 2012). Meeker et al. (2013) also reported lower ICCs for triclosan, 2,4- and 2,5-dichlorophenol (0.47, 0.38 and 0.49, respectively), but their ICC for benzophenone-3 (0.62) was the same as in our study population. Differences may be attributable to differences in exposure patterns between study populations, or to differences in the time between consecutive urinary collections. For example, urine samples were collected at gestational weeks 5.8, 20.6 and 33.5 on average in the Smith et al. study (2012), compared with 17.6, 22.8, and 32.8 weeks in our study.

Correcting for urine dilution improved the temporal reproducibility in urinary concentration for most of the phenols that we evaluated. Urinary creatinine and specific gravity followed similar patterns throughout pregnancy, and the estimates of urinary concentration reliability (correlations, ICCs) were similar for both methods. These findings suggest that either could be used to correct for urine dilution in pregnant women.

Because benzophenone-3 is commonly used in sunscreen for its sun-blocking properties (Calafat et al. 2008), exposure is more likely to be high during the summer months than the rest of the year, consistent with the seasonal variability in benzophenone-3 concentrations in our population (adjusted geometric means of 110 µg/L in April–September compared with 69 µg/L for samples collected October–March). Associations between hour of collection and benzophenone-3, triclosan, butylparaben and 2,4-dichlorophenol urinary concentrations suggest that exposures

changed considerably throughout the day. We sought to remove the variability in phenol urinary concentrations due to heterogeneity in collection conditions by implementing the 2-step standardization method proposed by Mortamais et al. (2012). However, ICCs obtained with the standardized concentrations were close to those obtained from the unstandardized concentrations, suggesting that study-related variability in sample collection conditions did not strongly influence intra-subject differences in biomarker levels.

Study strengths and limitations

Our study is the first to report phenol concentrations in amniotic fluid and maternal urine collected on the same day. In addition, we examined variability in phenol biomarker concentrations during the prenatal period for a diverse set of environmental phenols, including some that have not been widely studied (triclosan, benzophenone-3 and dichlorophenols). Limitations include the relatively small sample size and the low detection frequency of phenols in amniotic fluid. Additionally, although we made no exclusions on the basis of race/ethnicity, most of our study population was white. Because there is some racial/ethnic variability in exposure patterns, our results may not fully represent variability in the general population.

To better understand the link between maternal and fetal exposures, larger studies with more diverse populations are needed. Collecting more urine samples may improve the characterization of the urine concentration variability during pregnancy.

Conclusions

Our estimates of the reproducibility of benzophenone-3, triclosan, 2,4- and 2,5-dichlorophenols, methyl-, ethyl-, propyl- and butyl-paraben urinary concentrations over the course of the

pregnancy suggest that more than one urine sample is needed to reasonably represent the individual's exposure across pregnancy to these compounds or their precursors. As previously reported (Braun et al. 2011; Braun et al. 2012), we observed low reproducibility in bisphenol A urinary concentration among pregnant women, with little evidence of secondary fetal exposure to bisphenol A via amniotic fluid. Given the infrequent detection and much lower concentrations of most biomarkers in amniotic fluid than in urine, our results suggest that amniotic fluid may not be a suitable matrix for assessing fetal exposure to non-persistent phenols using the currently available detection techniques.

References

- Balakrishnan B, Henare K, Thorstensen EB, Ponnampalam AP, Mitchell MD. 2010. Transfer of bisphenol a across the human placenta. *Am J Obstet Gynecol* 202:393.e391-397.
- Bradman A, Barr DB, Henn BGC, Drumheller T, Curry C, Eskenazi B. 2003. Measurement of pesticides and other toxicants in amniotic fluid as a potential biomarker of prenatal exposure: A validation study. *Environ Health Perspect* 111:1779-1782.
- Braun JM, Kalkbrenner AE, Calafat AM, Bernert JT, Ye X, Silva MJ, et al. 2011. Variability and predictors of urinary bisphenol a concentrations during pregnancy. *Environ Health Perspect* 119:131-137.
- Braun JM, Smith KW, Williams PL, Calafat AM, Berry K, Ehrlich S, et al. 2012. Variability of urinary phthalate metabolite and bisphenol a concentrations before and during pregnancy. *Environ Health Perspect*.
- Calafat AM, Wong LY, Ye XY, Reidy JA, Needham LL. 2008. Concentrations of the sunscreen agent benzophenone-3 in residents of the united states: National health and nutrition examination survey 2003-2004. *Environ Health Perspect* 116:893-897.
- Casas M, Chevrier C, Hond ED, Fernandez MF, Pierik F, Philippat C, et al. 2013. Exposure to brominated flame retardants, perfluorinated compounds, phthalates and phenols in european birth cohorts: Enrieco evaluation, first human biomonitoring results, and recommendations. *Int J Hyg Environ Health* 216:230-242.
- Chevrier C, Petit C, Philippat C, Mortamais M, Slama R, Rouget F, et al. 2012. Maternal urinary phthalates and phenols and male genital anomalies. *Epidemiology* 23:353-356.
- Chou WC, Chen JL, Lin CF, Chen YC, Shih FC, Chuang CY. 2011. Biomonitoring of bisphenol a concentrations in maternal and umbilical cord blood in regard to birth outcomes and adipokine expression: A birth cohort study in taiwan. *Environ Health* 10:94.
- Edlow AG, Chen M, Smith NA, Lu C, McElrath TF. 2012. Fetal bisphenol a exposure: Concentration of conjugated and unconjugated bisphenol a in amniotic fluid in the second and third trimesters. *Reprod Toxicol* 34:1-7.
- Engel SM, Levy B, Liu ZS, Kaplan D, Wolff MS. 2006. Xenobiotic phenols in early pregnancy amniotic fluid. *Reprod Toxicol* 21:110-112.

- Fenichel P, Dechaux H, Harthe C, Gal J, Ferrari P, Pacini P, et al. 2012. Unconjugated bisphenol a cord blood levels in boys with descended or undescended testes. *Hum Reprod* 27:983-990.
- Ikezuki Y, Tsutsumi O, Takai Y, Kamei Y, Taketani Y. 2002. Determination of bisphenol a concentrations in human biological fluids reveals significant early prenatal exposure. *Hum Reprod* 17:2839-2841.
- Janjua NR, Frederiksen H, Skakkebaek NE, Wulf HC, Andersson AM. 2008. Urinary excretion of phthalates and paraben after repeated whole-body topical application in humans. *Int J Androl* 31:118-130.
- Lubin JH, Colt JS, Camann D, Davis S, Cerhan JR, Severson RK, et al. 2004. Epidemiologic evaluation of measurement data in the presence of detection limits. *Environ Health Perspect* 112:1691-1696.
- Meeker JD, Cantonwine DE, Rivera-Gonzalez LO, Ferguson KK, Mukherjee B, Calafat AM, et al. 2013. Distribution, variability, and predictors of urinary concentrations of phenols and parabens among pregnant women in puerto rico. *Environ Sci Technol* 47:3439-3447.
- Modena AB, Fieni S. 2004. Amniotic fluid dynamics. *Acta Biomed* 75 Suppl 1:11-13.
- Mortamais M, Chevrier C, Philippat C, Petit C, Calafat AM, Ye X, et al. 2012. Correcting for the influence of sampling conditions on biomarkers of exposure to phenols and phthalates: A 2-step standardization method based on regression residuals. *Environ Health* 11:29.
- Philippat C, Mortamais M, Chevrier C, Petit C, Calafat A, Ye X, et al. 2012. Exposure to phthalates and phenols during pregnancy and offspring size at birth. *Environ Health Perspect* 120:464-470.
- Rabe-Hesketh S. 2008. *Multilevel and longitudinal modeling using stata* 2nd ed. College Station: Texas:Stata Press.
- Rothman KJ, Greenland S, Lash TL. 2008. *Modern epidemiology*. 3rd ed. Philadelphia, PA:Lippincott.
- Smith KW, Braun JM, Williams PL, Ehrlich S, Correia KF, Calafat AM, et al. 2012. Predictors and variability of urinary paraben concentrations in men and women, including before and during pregnancy. *Environ Health Perspect*.
- Soni MG, Carabin IG, Burdock GA. 2005. Safety assessment of esters of p-hydroxybenzoic acid (parabens). *Food Chem Toxicol* 43:985-1015.

- Summers P, Bena J, Arrigain S, Alexis AF, Cooper K, Bordeaux JS. 2011. Sunscreen use: Non-hispanic blacks compared with other racial and/or ethnic groups. *Arch Dermatol* 147:863-864.
- Teeguarden JG, Calafat AM, Ye X, Doerge DR, Churchwell MI, Gunawan R, et al. 2011. Twenty-four hour human urine and serum profiles of bisphenol a during high-dietary exposure. *Toxicol Sci* 123:48-57.
- Underwood MA, Gilbert WM, Sherman MP. 2005. Amniotic fluid: Not just fetal urine anymore. *J Perinatol* 25:341-348.
- Volkel W, Colnot T, Csanady GA, Filser JG, Dekant W. 2002. Metabolism and kinetics of bisphenol a in humans at low doses following oral administration. *Chem Res Toxicol* 15:1281-1287.
- Whitehead TP, Nuckols JR, Ward MH, Rappaport SM. 2012. Carpet-dust chemicals as measures of exposure: Implications of variability. *Emerg Themes Epidemiol* 9:2.
- Wolff MS, Engel SM, Berkowitz GS, Ye X, Silva MJ, Zhu C, et al. 2008. Prenatal phenol and phthalate exposures and birth outcomes. *Environ Health Perspect* 116:1092-1097.
- Woodruff TJ, Zota AR, Schwartz JM. 2011. Environmental chemicals in pregnant women in the united states: Nhanes 2003-2004. *Environ Health Perspect* 119:878-885.
- Yamada H, Furuta I, Kato EH, Kataoka S, Usuki Y, Kobashi G, et al. 2002. Maternal serum and amniotic fluid bisphenol a concentrations in the early second trimester. *Reprod Toxicol* 16:735-739.
- Ye X, Bishop AM, Reidy JA, Needham LL, Calafat AM. 2007. Temporal stability of the conjugated species of bisphenol a, parabens, and other environmental phenols in human urine. *J Expo Sci Environ Epidemiol* 17:567-572.
- Ye XY, Zsuzsanna K, Needham LL, Calafat AM. 2005. Quantification of urinary conjugates of bisphenol a, 2,5-dichlorophenol, and 2-hydroxy-4-methoxybenzophenone in humans by online solid phase extraction-high performance liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem* 383:638-644.

Table 1: Characteristics of 71 women with three urine measurements during pregnancy.
SARAEH study, 2005 to 2008.

Characteristic	N	% -
Maternal age (years)		
25 to 29.9	5	7
30 to 34.9	21	30
35 to 40.9	45	63
Race/ethnicity		
White	51	72
Black	5	7
Hispanic	11	15
Asian	3	4
American Indian	1	1
Pre-pregnancy BMI (kg/m ²)		
17.8 to 18.4	3	4
18.5 to 24.9	53	75
25 to 41	15	21
Maternal education		
High school	9	13
Any college	23	32
Graduate school, graduate degree	39	55
Conception methods		
Reproductive Techniques	3	4
Natural	68	96
Amniotic fluid sampling reason		
Advanced Maternal Age	49	69
Abnormal Genetic Screen	10	14
Elective	11	15
Other	1	1

Table 2: Distribution of phenol concentrations ($\mu\text{g/L}$) measured in amniotic fluid and in urine collected at three time periods during pregnancy among women of the SARAHEH study, 2005 to 2008.

Analyte	Urine (n=71) ^a												Amniotic fluid (n=69) ^b						Ratio of first urine sample : amniotic fluid		
	1 st sample				2 nd sample				3 rd sample				%>LOD	5 th	50 th	95 th	5 th	50 th	95 th		
	LOD	%>LOD	5 th	50 th	95 th	%>LOD	5 th	50 th	95 th	%>LOD	5 th	50 th								95 th	
Phenols																					
($\mu\text{g/L}$)																					
2,4-DCP ^c	0.2	68	<LOD	0.6	5.7	94	<LOD	1.1	8.3	93	<LOD	1.0	14	1 ^c	<LOD	<LOD	LOD	1.4	1.4	9	
2,5-DCP	0.2	100	0.8	7.5	178	100	1.8	12.8	364	100	1.7	14.2	466	16	<LOD	LOD	5.2	3.5	297	1756	
BPA	0.4	62	<LOD	0.6	7.0	92	<LOD	1.8	9.5	82	<LOD	1.2	5.0	3	<LOD	<LOD	LOD	1.4	12	85	
BP3	0.4	100	4.1	53.5	2220	100	5.7	77	6740	100	8.9	76.1	3180	61	<LOD	0.8	15.7	8	110	1648	
TCS	2.3	79	<LOD	6.5	789	75	<LOD	15.4	734	80	<LOD	16.2	514	6	<LOD	<LOD	19.4	1.4	16	354	
MP	1.0	100	9.6	100	985	100	19	272	1670	100	9.7	156	1830	42	<LOD	<LOD	3.3	23	196	827	
EP	1.0	59	<LOD	2.8	58	65	<LOD	4.5	347	59	<LOD	1.7	144	0	<LOD	<LOD	<LOD	3.5	30	304	
PP	0.2	100	0.5	28.7	424	100	1.6	45.6	531	99	0.8	36.5	589	58	<LOD	0.3	1.4	7	116	2130	
BP	0.2	70	<LOD	1.5	26	85	<LOD	1.9	56.3	73	<LOD	1.7	58.2	6	<LOD	<LOD	0.3	1.4	22	179	
Markers of urine dilution																					
Creatinine (mg/dL) ^d	3.5	100	13.7	44.1	256	100	15.3	86.8	221	100	16	90.9	175								
Specific gravity			1.003	1.008	1.026		1.003	1.02	1.03		1.004	1.015	1.026								

^a Concentrations not corrected for urine dilution, concentrations < LOD were replaced by $\text{LOD}/\sqrt{2}$. -

^b Concentrations < LOD were replaced by instrumental reading values. -

^c 2,4-DCP was only assessed in 11 amniotic fluid samples. -

^d n = 60, creatinine concentration was missing for 11 urine samples from the 1st sampling. -

Abbreviations: BP: butylparaben, BPA: bisphenol A, BP3: benzophenone-3, DCP: dichlorophenol, EP: ethylparaben, MP: methylparaben, PP: propylparaben, TCS: triclosan. -

Table 3: Adjusted associations of ln-transformed amniotic fluid concentrations of benzophenone-3 (BP3) and propylparaben (PP)^a with potential predictors among 68 pregnant women from the SARAEH study, 2005 to 2008.

Characteristics	n	Change in ln-BP3 concentration (µg/L)	Estimated BP3 concentration (GM, µg/L)	p-trend ^b	Change in ln-PP concentration (µg/L)	Estimated PP concentration (GM, µg/L)	p-trend ^b
		β (95% CI)			β (95% CI)		
Urine concentration (ln-unit, µg/L) ^c		0.52 (0.3, 0.7)			0.36 (0.2, 0.5)		
Time of amniotic fluid storage at -20 C (weeks)				0.35			0.35
≤ 4	17	Reference	0.71		Reference	0.20	
4 to 16	12	-0.28 (-1.6, 1.0)	0.54		0.93 (0.0, 1.9)	0.50	
16 to 20	14	-0.39 (-1.7, 0.9)	0.48		0.02 (-0.9, 1.0)	0.20	
> 20	27	-0.61 (-1.9, 0.7)	0.39		-0.34 (-1.3, 0.6)	0.14	
Gestational age at sampling (weeks)				0.71			0.49
15 to 16.9	31	Reference	0.51		Reference	0.20	
17 to 18.9	26	0.11 (-0.7, 0.9)	0.57		-0.18 (-0.8, 0.4)	0.17	
19 to 23	11	-0.35 (-1.8, 1.1)	0.36		0.55 (-0.4, 1.5)	0.35	
Maternal age (year)				0.09			0.56
<31	10	Reference	1.03		Reference	0.18	
31 to 35.9	24	-0.59 (-1.8, 0.7)	0.58		0.06 (-0.9, 1.0)	0.20	
36 to 40	34	-1.04 (-2.3, 0.2)	0.37		0.20 (-0.7, 1.1)	0.23	
BMI				0.67			0.34
< 25	53	Reference	0.53		Reference	0.23	
≥ 25	15	-0.23 (-1.3, 0.9)	0.42		-0.39 (-1.2, 0.4)	0.15	
Race/ethnicity							
White	48	Reference	0.63		Reference	0.20	
Non-white	20	-0.77 (-1.9, 0.3)	0.29		0.16 (-0.6, 0.9)	0.23	
Pregnancy complications related to placenta ^d							
No	58	Reference	0.50		Reference	0.23	
Yes	10	-0.05 (-1.3, 1.2)	0.48		-0.79 (-1.7, 0.1)	0.11	
Baby sex							
Boy	36	Reference	0.59		Reference	0.27	
Girl	32	-0.37 (-1.2, 0.4)	0.41		-0.56 (-1.2; 0.1)	0.15	

^a Phenols detected in at least 50% of the amniotic fluid samples. -

^b p-trend was estimated using categorical variables whose values corresponded to the category-specific - median value (Rothman et al. 2008). -

^c Corrected for specific gravity. -

^d Including oligohydroamnios, choriarniotis, small placenta, placenta praevia and pre-eclampsia.

This analysis was restricted to 68 pregnant women because the child's sex was missing for one woman.

Abbreviations: AF: amniotic fluid, BP3: benzophenone-3, PP: propyl paraben, GM: geometric mean

Table 4: Intraclass Correlation Coefficients for phenol biomarker concentrations measured in urine samples collected at several times during pregnancy. SARAHEH study, 2005 to 2008.

Analyte	Specific gravity corrected concentrations ^a	Creatinine corrected concentrations ^b	Uncorrected concentrations ^a	Concentrations standardized for collection conditions ^{a,c}
2,4-dichlorophenol	0.60	0.48	0.47	0.60
2,5-dichlorophenol	0.61	0.62	0.52	0.60
Bisphenol A	0.11	0.14	0.23	0.15
Benzophenone-3	0.62	0.70	0.57	0.61
Triclosan	0.58	0.61	0.56	0.60
Methylparaben	0.61	0.61	0.52	0.62
Ethylparaben	0.48	0.44	0.48	0.48
Propylparaben	0.55	0.54	0.51	0.54
Butylparaben	0.56	0.64	0.54	0.61

^a n = 213 samples from 71 women. -

^b n = 180 samples from 60 women with creatinine measured in all three urine samples. -

^c All concentrations were standardized for specific gravity. Bisphenol A concentration was additionally standardized for gestational age at sampling and sampling hour. Butylparaben, triclosan and 2,4-dichlorophenol were additionally standardized for hour of sampling. Benzophenone-3 and methylparaben were additionally standardized for hour and season of sampling, propylparaben for season of sampling and ethylparaben for day of sampling. -

Figure legend

Figure 1. Box plots of the distribution of specific-gravity corrected urinary phenol concentrations according to undetected or detectable levels of phenol metabolites in amniotic fluid collected on the same day. In general urinary concentrations were higher in women with detectable amniotic fluid concentrations. Lower whisker represents the lowest value within 1.5 IQR of the 25th percentile. The bottom of box is the 25th percentile, the line is the median, and the top of box is the 75th percentile. Upper whisker is the highest value within 1.5 IQR of the 75th percentile. Dots represent outlier values. Distributions are not shown for bisphenol A, 2,4-dichlorophenol, or ethyl paraben because they were detected in only 2, 1, and 0 amniotic fluid samples, respectively.

