



ENVIRONMENTAL
HEALTH
PERSPECTIVES

<http://www.ehponline.org>

**High-Density Livestock Production and Molecularly
Characterized MRSA Infections in Pennsylvania**

**Joan A. Casey, Bo Shopsin, Sara E. Cosgrove,
Keeve E. Nachman, Frank C. Curriero, Hannah R. Rose,
and Brian S. Schwartz**

<http://dx.doi.org/10.1289/ehp.1307370>

Received: 15 July 2013

Accepted: 6 February 2014

Advance Publication: 7 February 2014

High-Density Livestock Production and Molecularly Characterized MRSA Infections in Pennsylvania

Joan A. Casey,^{1,2} Bo Shopsin,³ Sara E. Cosgrove,⁴ Keeve E. Nachman,^{1,2} Frank C. Curriero,^{1,5} Hannah R. Rose,³ and Brian S. Schwartz^{1,6,7}

¹Department of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA; ²Center for a Livable Future, Johns Hopkins University, Baltimore, Maryland, USA; ³ Department of Medicine, New York University School of Medicine, New York, New York, USA; ⁴Division of Infectious Disease, Johns Hopkins School of Medicine, Baltimore, Maryland, USA; ⁵Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA; ⁶Department of Medicine, Johns Hopkins School of Medicine, Baltimore, Maryland, USA; ⁷Center for Health Research, Geisinger Health System, Danville, Pennsylvania, USA

Correspondence to Brian S. Schwartz, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe Street, Room W7041, Baltimore, MD 21205 USA. E-mail: bschwartz@jhsph.edu.

Running title: MRSA subgroups and high-density livestock production

Acknowledgments: D.G. Mercer, J.J. DeWalle, M.S. Krutsick, and S.H. Brandau participated in study management, data collection, and patient geocoding. D. Altman assisted with microbiologic testing. This research was supported by the New York University-Geisinger Seed Grant Program, the Johns Hopkins Center for a Livable Future, the Johns Hopkins Sommer Scholarship and the National Institute of Environmental Health Sciences (NIEHS; grant ES07141).

Competing Financial Interests: The authors declare that they have no actual or potential competing financial interests.

Abstract

Background: European studies suggest that living near high-density livestock production increases the risk of sequence type ST398 methicillin-resistant *Staphylococcus aureus* (MRSA) colonization. To our knowledge, no studies have evaluated associations between livestock production and human infection by other strain types.

Objectives: We evaluated associations between MRSA molecular subgroups and high-density livestock production.

Methods: We conducted a yearlong 2012 prospective study on a stratified random sample of patients with culture-confirmed MRSA infection; we oversampled patients from the Geisinger Health System with exposure to high-density livestock production in Pennsylvania. Isolates were characterized using *S. aureus* protein A (*spa*) typing and detection of Panton-Valentine leukocidin and *scn* genes. Patients with one of two specific MRSA strains were compared to patients with all other strains of MRSA isolates using logistic regression that accounted for the sampling design, for two different exposure models: one based on the location of the animals (livestock model) and the other on crop field application of manure (crop field model).

Results: Of 196 MRSA isolates, we identified 30 *spa* types, 47 PVL-negative and 15 *scn*-negative isolates, and no ST398 MRSA. Compared with quartiles 1-3 combined, the highest quartiles of swine livestock and dairy/veal crop field exposures were positively associated with community-onset-PVL-negative MRSA (CO-PVL-negative MRSA versus all other MRSA), with adjusted odds ratios of 4.24 (95% CI: 1.60, 11.25) and 4.88 (95% CI: 1.40, 17.00), respectively. The association with CO-PVL-negative MRSA infection increased across quartiles of dairy/veal livestock exposure (trend $p = 0.05$).

Conclusions: The findings suggest that other MRSA strains, beyond ST398, may be involved in livestock-associated MRSA infection in the U.S.

Introduction

Over the past decade, the incidence of community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA) infection has increased in the United States (Dukic et al. 2013). These CA-MRSA infections cost third-party payers between \$478 and \$2200 million annually (Lee et al. 2013). Beginning in the mid-2000s, European research suggested that a portion of the increased incidence might be attributable to high-density livestock production as studies isolated the same MRSA strains from infected farmers and their livestock (Harrison et al. 2013; Hartmeyer et al. 2010). In Europe, multilocus sequence type 398 (ST398) has been the most common colonizer of livestock, specifically swine, and their owners (Graveland et al. 2010; Lewis et al. 2008). This association has led many to refer to ST398 as livestock-associated MRSA (LA-MRSA) (Price et al. 2012). Importantly, pathways for community transmission have been identified, with MRSA isolated from the air and soil at least 150m from swine facilities (Gibbs et al. 2006; Schulz et al. 2012) and from meat processing and consumption (Molla et al. 2012; Waters et al. 2011).

To our knowledge, no North American studies have evaluated residence in rural communities as a risk factor for MRSA infection of specific molecular types. Four studies of colonization in farmers have disparate findings. The first reported ST398 as the only type colonizing pigs and farmers (Smith et al. 2009); the second found mainly ST398, but also ST5 (Khanna et al. 2008); while two more recent studies identified ST5 (t002 by *spa* typing) (Monecke et al. 2011), as the primary colonizer of pigs and of veterinary students visiting farms (Frana et al. 2013; Molla et al. 2012). In the U.S., ST5 has been considered a healthcare-associated MRSA clone, but has begun to appear in the community in those without healthcare risk factors (Klevens et al. 2006). Nearly all MRSA isolates associated with livestock colonization lack the genes encoding Panton-

Valentine leukocidin (PVL) (Smith et al. 2009; Sunde et al. 2011). Lastly, recent studies have reported that *scn*, the gene encoding staphylococcal complement inhibitor (SCIN), is often absent in MRSA strains that colonize livestock (McCarthy et al. 2011; Sung et al. 2008; Verkaik et al. 2011).

In a prior study in Pennsylvania using electronic health records (EHR) from 2005-2010 from a large health system we reported associations between CA-MRSA infection and residential proximity to high-density livestock operations and the crop fields to which manure was applied (Casey et al. 2013). Due to the retrospective study design, we were unable to obtain MRSA isolates from patients. The objectives of the present study were to: (1) prospectively collect MRSA isolates from patients residing in communities with and without high-density livestock production; (2) characterize these isolates by *spa* typing and polymerase chain reaction (PCR) for the presence of *lukF-lukS* genes (encoding PVL) and for the *scn* gene; and (3) assess associations of high-density livestock production with the molecular subgroups.

Methods

Setting, study design, and participants

The study area was a 38 county region of central and northeast Pennsylvania with 3.8 million inhabitants in which Geisinger Health System provides primary care services from 41 community practice clinics (Figure 1). The primary care population is representative of the region's population (Casey et al. 2013). We compared patients infected with two specific MRSA strains to patients infected with all other strains as described below. Institutional review boards at the Geisinger Health System and the Johns Hopkins Bloomberg School of Public Health approved the study and waived informed consent.

Patients with MRSA infection identified by culture between January 1, 2012 and December 31, 2012, who had a Geisinger primary care provider, were eligible for inclusion. The cases were stratified as community-onset or hospital-onset and as residing in an area of high-density livestock production or not. Community-onset infections were patients with a positive culture collected in the outpatient setting or ≤ 2 days after hospitalization. Hospital-onset infections were patients with a positive culture collected > 2 days after hospital admission (David and Daum 2010). Onset date was defined as the date the culture was ordered. Patients living in a zip code with a high-density livestock operation or a crop field to which manure from such an operation was applied were defined as residing in an area of high-density livestock production, as described in detail below. Using these case definitions we completed a random stratified sample (a sample was evaluated due to budgetary limitations) to select cases into four *a priori* categories (Table 1). We over-selected patients that resided in areas of high-density livestock production to improve the power to detect associations.

Patient data collection

MRSA isolates (one per patient) collected in clinical practice were sent to a central laboratory, grown on conventional blood agar, and stored at 4°C. A research assistant selected appropriate isolates and obtained samples for transport using sterile swab applicators (BBL Cultureswabs; Becton Dickinson). A total of 203 unique isolates collected from sites of infection were sent unrefrigerated to New York University Langone Medical Center for genotypic testing, where isolates were sub-cultured onto agar plates and incubated overnight at 37°C. DNA was then extracted by mechanical lysis. We obtained information on patient demographics, inpatient, outpatient, and emergency department encounters, procedures, medication orders, and laboratory data from the EHR. Antimicrobial resistance testing on clindamycin, erythromycin, gentamicin,

rifampin, tetracycline, trimethoprim-sulfamethoxazole (TMP-SMZ), and vancomycin was also available from the EHR. Multidrug resistance was reported as resistance to ≥ 3 unique antimicrobials.

Genotyping

DNA sequence analysis of the protein A gene variable repeat region (*spa* typing) was completed using the primers TIGR-F (5'-GCCAAAGCGCTAACCTTTTA-3') and TIGR-R (5'-TCCAGCTAATAACGCTGCAC-3') (Shopsin et al. 1999). *Spa* types were assigned a StaphType (e.g., t008, t002) using the Ridom SpaServer database available at <http://www.spaserver.ridom.de>. We subsequently mapped *spa* types to clonal complex categories (clonal complex 5 or 5 like, 8 or 8 like, or other) using the Ridom SpaServer *spa* to multilocus sequence type (MLST) database available at <http://spa.ridom.de/mlst.shtml>. We also checked for the presence of PVL genes using *S. aureus* ATCC 29775 as a reference strain and primers as previously described (Lina et al. 1999). The presence of *scn* was determined by an additional PCR (Sung et al. 2008).

Prior literature from northern Europe suggested that ST398 would be a cause of human infection in those exposed to high-density livestock production (Hartmeyer et al. 2010; Wulf et al. 2012), but we did not identify any ST398 infections. Therefore, before beginning statistical analysis, we identified two other strains of interest. Most CA-MRSA contains PVL genes (David and Daum 2010), while most MRSA (both ST5 and ST398) isolated from humans and animals with exposure to high-density livestock production in North America has been PVL-negative (Frana et al. 2013; Price et al. 2012; Tattevin et al. 2012). Therefore, we selected community-onset PVL-negative (CO-PVL-negative) MRSA isolates as a unique group with potential ties to high-density livestock production. We created a second molecular subgroup based on recent evidence

that only about 25% of animal *S. aureus* isolates carry the *scn* gene, while human isolates carry it about 90% of the time (Sung et al. 2008; Verkaik et al. 2011). To complete analysis we created two case groups: (1) CO-PVL-negative MRSA and (2) *scn*-negative MRSA, and compared these groups to patients with all other MRSA types.

Geospatial estimates of high-density livestock exposure

Pennsylvania Act 38 of 2005 requires nutrient management plans (NMPs) to regulate concentrated animal operations (CAOs), operations where livestock density exceeds two animal equivalent units (AEUs, 1000 pounds of animal weight) per acre and where the total number of AEUs exceeds eight. NMPs detail appropriate manure handling, storage, and land application to crop fields, both on-site where the animals are raised and off-site. Plans also contain data on operation location and acreage and livestock type. In addition, the Pennsylvania Department of Environmental Protection requires NMPs from the largest operations called concentrated animal feeding operations (>1000 AEUs, or 301-1000 AEUs and a CAO, or per federal regulation) (Environmental Protection Agency 2012). For our analysis, we defined a high-density livestock operation as: (1) >2 AEUs per acre and >8 AEUs total; or (2) >300 AEUs total.

We used ArcGIS (version 10.0, Esri) to geocode livestock operations, treated crop fields, and patients at their home address. While patients were sampled using presence of high-density livestock production in their zip code, this was not used to generate exposure metrics. We created two exposure metrics for livestock operations (swine or dairy/veal):

$$\text{Exposure for patient } j = \sum_{i=1}^n (a_i / d_{ij}^2), \quad [1]$$

where n is the number of operations, a_i is the AEUs of livestock at operation i , and d_{ij}^2 is the squared distance (in meters) between operation i and patient j . In addition, we created two

metrics for exposure to treated crop fields (swine or dairy/veal manure applications) during the season of infection:

$$\text{Exposure for patient } j = \sum_{k=1}^n (c_k / d_{kj}^2), \quad [2]$$

where n is the number of treated crop fields, c_k is the concentration of manure (gallons per square meter) applied to field k during the season of diagnosis, and d_{kj}^2 is the squared distance (in meters) between field centroid k and patient j (Casey et al. 2013). We modeled the exposure variables as quartiles to allow for non-linear associations and for ease of interpretability.

Statistical methods

A multivariable logistic regression model was used to estimate associations between exposures and outcomes (CO-PVL-negative MRSA or *scn*-negative MRSA compared with all other MRSA strains combined). By study design, we oversampled cases living in zip codes with high-density livestock production. To account for this sampling design while obtaining unbiased regression estimates and robust standard errors, each individual was assigned a sample weight (weight = total MRSA patients identified in a stratum / total included in the present analysis) (Table 1), a measure of the number of patients represented by each sampled individual (Vittinghoff et al. 2005). We used the p -value from Fisher's exact test to compare molecular profiles of the two subgroups of interest to all other MRSA isolates (Table 2).

Models were constructed with attention to the small sample size. Our primary models compared the highest quartile of exposure to livestock or treated crop fields to all other quartiles combined. In addition, we modeled quartiles of exposure as an ordinal variable (0, 1, 2, 3) and used the p -value of the resulting coefficient as a test of linear trend. For CO-PVL-negative MRSA we also estimated associations with individual quartiles of exposure relative to the lowest quartile, but

numbers of observations were too small to run comparable models for *scn*-negative MRSA. Similarly, adjustment for confounders selected based on *a priori* information (Casey et al. 2013a and 2013b) was limited to models of CO-PVL-negative MRSA. Specifically, we adjusted for sex, age (continuous), ever-smoking status, season of infection [winter (December-February) vs. all other seasons combined], physician order for an antibiotic in the 365 to 14 days preceding diagnosis, and ever-receiving Medical Assistance as health insurance, as a surrogate for low individual socioeconomic status (Bratu et al. 2006; Casey et al. 2013). We were unable to adjust models for race/ethnicity and isolate source because data were off-support (Oakes 2006). We did not adjust for location of onset because location was part of the CO-PVL-negative MRSA definition. Statistical analyses were performed using Stata 11.2 (StataCorp) using the svy commands for weighted regression and R version 3.0.0 (R Foundation for Statistical Computing).

Results

Study population characteristics

From January 1 to December 31, 2012 we collected data on 203 MRSA infections (from a pool of 1128 patients with MRSA infection) from patients in 20 counties in Pennsylvania. The analysis included 196 cases (Figure 1) after exclusion of isolates lacking the *mecA* gene ($n = 4$) and patients who could not be geocoded or resided outside the study area ($n = 3$). We identified 30 unique Ridom *spa* types, 47 PVL-negative isolates and 15 *scn*-negative isolates, and no ST398 MRSA (Table 2). Four isolates were both CO-PVL-negative and *scn*-negative. Patients with CO-PVL-negative MRSA ($n = 23$) as well as patients with *scn*-negative MRSA ($n = 15$) were older than patients in the all other MRSA group ($n = 162$) (Table 3). Patients with the two MRSA strains of interest were less likely to reside in cities than were patients with all other

MRSA strains (Table 3). Patients in the third and fourth quartiles of swine livestock exposure were more likely than those in the first and second quartiles to be diagnosed in the fall (Table 3). About 95% of patients with MRSA onset in the winter fell into the first or second quartile of swine and dairy/veal crop field exposure (data not shown).

Molecular testing

We identified 30 *spa* types, but two predominated: t008, n = 122 (62.2%) and t002, n = 27 (13.8%) (Table 2). Nearly all t008, n = 116 (95.1%), was community-onset, while the majority of t002, n = 17 (63.0%), was hospital-onset. Isolates lacking *scn* (n = 15) were distributed across multiple *spa* types, and 80% (n = 12) were community-onset. CO-PVL-negative isolates were mostly *spa* types associated with clonal complex 5.

Antimicrobial susceptibility

CO-PVL-negative strains were more often resistant to antibiotics commonly used to treat MRSA infection (i.e. clindamycin, tetracycline, and TMP-SMZ) than all other CO-PVL-positive MRSA strains. The 30 HO-MRSA isolates were resistant to more antibiotics compared to CO-MRSA: clindamycin (83.3% vs. 12.7%), tetracycline (6.7% vs. 3.6%), TMP-SMZ (3.3% vs. 1.2%), and multidrug (83.3% vs. 14.5%). Resistance patterns did not appear to be associated with agriculture exposure variables, but tetracycline-resistance in a small number of isolates (n = 4) was associated with infection by CO-PVL-negative MRSA ($p = 0.02$).

Associations with livestock and crop field manure exposure

High-density livestock production was not associated with *scn*-negative MRSA infection in unadjusted analysis (Table 4). Swine livestock exposure was associated with CO-PVL-negative MRSA in unadjusted and adjusted analysis (Table 4), which appeared driven by the association

in the fourth quartile of exposure (Figure 2). There was a positive but non-significant association with the dichotomized dairy/veal livestock exposure variable (OR = 2.42; 95%CI: 0.85, 6.88). There was a trend ($p = 0.05$) of increasing odds of CO-PVL-negative MRSA across quartiles of dairy/veal livestock exposure; however, all odds ratios were increasingly imprecise with higher exposure for individual versus lowest quartile exposures (Figure 2). Those in the highest quartile of swine crop field exposure had an odds ratio of 2.38 (95% CI: 0.78, 7.28) of being a CO-PVL-negative MRSA case compared to quartiles 1-3 (Table 4). Dichotomous dairy/veal crop field exposure was associated with CO-PVL-negative MRSA in adjusted analyses (OR = 4.88; 95% CI: 1.40, 17.00) (Table 4). Neither crop field variable evidenced a statistically significant trend across quartiles (Figure 2). Associations strengthened slightly as we added sex, age, ever-smoking, antibiotic order, and Medical Assistance to the models. Season of onset was a strong confounder in the crop field exposure models (e.g., the unadjusted OR of 0.79; 95% CI: 0.20, 3.09 increased to 2.38; 95% CI: 0.78, 7.28 [Table 4] for the swine crop field model, primarily due to the inclusion of season).

Discussion

This analysis compared patients infected by two specific MRSA strains to patients infected by all other MRSA strains and found a number of associations with high-density livestock production. The livestock model incorporated AEU's at the farming operation and distance to the patient residence, while the crop field model incorporated manure volume, crop field area, and distance to the patient residence. Although adjusted estimates were imprecise due to small numbers, each exposure variable was associated with increased odds of CO-PVL-negative MRSA infection, and three of four were statistically significant. Specifically, there were increased odds comparing the fourth quartiles to the other three quartiles of both swine livestock exposure and dairy/veal crop

field exposure. There was also a significant trend of increasing odds across the four quartiles for dairy/veal livestock exposure. For the fourth exposure variable, swine crop field, the odds ratio in the fourth quartile versus the other three was elevated, but the estimate was not statistically significant. Unexpectedly, we did not identify any patients with MRSA infection due to ST398, the MRSA subtype that previously has been associated with livestock operations, mainly based on studies of colonization in Europe. This raises two issues. First, to understand risk factors for MRSA infection we need to study MRSA infection rather than colonization. Second, while ST398 may be prevalent among swine and swine farmers in the U.S. other strains found on farms may actually be more important for infection.

When we did not identify any infections due to ST398, we evaluated associations of livestock production with two other MRSA strains compared to patients infected by other MRSA strains. To our knowledge, only one U.S. study has reported PVL-positive MRSA isolated from swine (Osadebe et al. 2013). Associations with CO-PVL-negative MRSA were of interest because many highly successful human epidemic MRSA clones, like CC5, do not contain PVL genes and have recently been isolated from pigs and pig farmers in the U.S. (Frana et al. 2013; Molla et al. 2012; Osadebe et al. 2013). We observed eight tetracycline-resistant isolates, four of which were CO-PVL-negative MRSA. This resistance could plausibly be a marker of high-density livestock production as tetracycline is the antibiotic most commonly sold for use in U.S. farm animals and isolates resistant to tetracycline have been observed in workers at industrial livestock operations (Rinsky et al. 2013; U.S. Food and Drug Administration 2010). Additionally, only 6.7% of HO-MRSA was resistant to tetracycline, suggesting that the association of CO-PVL-negative MRSA is not likely attributable to the healthcare system. Likewise, we selected *scn*-negative MRSA strains because we expected MRSA associated with livestock production to lack *scn* (McCarthy

et al. 2011; Sung et al. 2008; Verkaik et al. 2011). Previous work showing selection against *scn* in animal hosts dealt primarily with ST398 (McCarthy et al. 2011), a strain of MRSA already adapted to animals (Price et al. 2012). Since the *scn* gene produces a protein that hinders phagocytosis of *S. aureus* by human neutrophils (Sung et al. 2008) this work might not translate directly to human-adapted strains of MRSA (e.g., ST5 or ST8), especially in the study of infection rather than colonization.

A few U.S. studies have molecularly characterized MRSA isolated from agricultural workers (Frana et al. 2013; Molla et al. 2012; Osadebe et al. 2013; Smith et al. 2013; Smith et al. 2009). As those studies only described MRSA carriage in persons with direct contact with swine, they may have low utility regarding inference about MRSA infection in the community. Two studies reported transient carriage (<24 hours) after short-term exposure to pigs and veal calves (Frana et al. 2013; van Cleef et al. 2011). Even if carriage persists after exposure, and the risk of infection increases, infection is not assured (Davis et al. 2004).

The body of literature on MRSA carriage is also relevant to the absence of ST398 among our MRSA isolates. While there is ample evidence that those in contact with livestock have increased prevalence of ST398 MRSA colonization (Graveland et al. 2010; Smith et al. 2009; Voss et al. 2005), data suggests ST398 MRSA infection is rare (Golding et al. 2010; Salmenlinna et al. 2010; Wulf et al. 2012). Strains of MRSA adapted primarily to animals, like ST398, may have diminished ability to infect humans (Graveland et al. 2011; Price et al. 2012). A recent Italian study reported a much greater proportion of ST398 among MRSA-colonized patients, compared to those with a MRSA infection where ST8 was most prevalent (Monaco et al. 2013).

Confounding by occupation or socioeconomic status may have impacted our associations. We could not obtain patient occupation from the EHR, and therefore cannot exclude the possibility that some CO-PVL-MRSA cases were occupational versus community acquired. We adjusted for Medical Assistance, a surrogate for low socioeconomic status that was associated with MRSA infection in our previous study (Casey et al. 2013), but we acknowledge that there may be residual confounding due to socioeconomic status. Since the CO-PVL-negative MRSA group had several characteristics of healthcare-associated MRSA [(HA-MRSA), e.g., older age, antibiotic resistance, lack of PVL genes, many *spa* type t002] the association with high-density livestock production could have arisen if HA-MRSA was more common in rural areas. However, in a previous study conducted in this region HA-MRSA was not more common in rural areas (Casey et al. 2013). Also, HA-MRSA in the current analysis defined by common epidemiologic criteria (Morrison et al. 2006) was not associated with high-density livestock production (data not shown). We adjusted models for season of onset because, as expected, MRSA patients with onset in the winter had low exposure to treated crop fields since manure application is restricted in the winter.

The novel observations in this study require replication. We completed analysis on a limited number of patients with a MRSA infection and while we incorporated several aspects of operations and crop fields into our individual-level exposure estimates, we did not collect environmental samples. Lastly, we could not calculate incidence rates because we only included patients with MRSA infection.

Conclusions

To our knowledge, no previous studies in the U.S. have evaluated associations between patients infected with specific MRSA strains and high-density livestock production. We identified associations of CO-PVL-negative MRSA with swine and dairy/veal livestock operations and application of dairy/veal manure to crop fields. Our findings were based on small numbers of observations and indirect estimates of exposure, but if confirmed, have important implications for the role of livestock operations in the MRSA epidemic, the different roles of colonization and infection studies, and the identification of other MRSA strains that may be arising from high-density livestock production. To help estimate the public health burden that high-density livestock production may place on the U.S. healthcare system, future studies should use MRSA infection, along with colonization, as their outcomes of interest. Our study also indicates the need to look beyond MRSA ST398 when investigating MRSA infection associated with high-density livestock production in the U.S.

References

- Bratu S, Landman D, Gupta J, Trehan M, Panwar M, Quale J. 2006. A population-based study examining the emergence of community-associated methicillin-resistant *Staphylococcus aureus* USA300 in New York City. *Ann Clin Microbiol Antimicrob* 5:29.
- Casey JA, Curriero FC, Cosgrove SE, Nachman KE, Schwartz BS. 2013. High-density livestock operations, crop field application of manure, and risk of community-associated methicillin-resistant *Staphylococcus aureus* infection in Pennsylvania. *JAMA Intern Med* 10.1001/jamainternmed.2013.10408 [Online 16 September 2013].
- David MZ, Daum RS. 2010. Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev* 23:616-687.
- Davis KA, Stewart JJ, Crouch HK, Florez CE, Hospenthal DR. 2004. Methicillin-resistant *Staphylococcus aureus* (MRSA) nares colonization at hospital admission and its effect on subsequent MRSA infection. *Clin Infect Dis* 39(6):776-782.
- Dukic VM, Lauderdale DS, Wilder J, Daum RS, David MZ. 2013. Epidemics of community-associated methicillin-resistant *Staphylococcus aureus* in the United States: a meta-analysis. *PLoS One* 8(1):e52722.
- Environmental Protection Agency. 2012. Regulatory Definitions of Large CAFOs, Medium CAFOs, and Small CAFOs. Available: http://www.epa.gov/npdes/pubs/sector_table.pdf [accessed 10 July 2013].
- Frana TS, Beahm AR, Hanson BM, Kinyon JM, Layman LL, Karriker LA, et al. 2013. Isolation and characterization of methicillin-resistant *Staphylococcus aureus* from pork farms and visiting veterinary students. *PLoS One* 8(1):e53738.
- Gibbs SG, Green CF, Tarwater PM, Mota LC, Mena KD, Scarpino PV. 2006. Isolation of antibiotic-resistant bacteria from the air plume downwind of a swine confined or concentrated animal feeding operation. *Environ Health Perspect* 114(7):1032-1037.
- Golding GR, Bryden L, Levett PN, McDonald RR, Wong A, Wylie J, et al. 2010. Livestock-associated methicillin-resistant *Staphylococcus aureus* sequence type 398 in humans, Canada. *Emerg Infect Dis* 16(4):587-594.

- Graveland H, Wagenaar JA, Bergs K, Heesterbeek H, Heederik D. 2011. Persistence of livestock associated MRSA CC398 in humans is dependent on intensity of animal contact. *PLoS One* 6(2):e16830.
- Graveland H, Wagenaar JA, Heesterbeek H, Mevius D, van Duijkeren E, Heederik D. 2010. Methicillin-resistant *Staphylococcus aureus* ST398 in veal calf farming: human MRSA carriage related with animal antimicrobial usage and farm hygiene. *PLoS One* 5(6):e10990.
- Harrison EM, Paterson GK, Holden MT, Larsen J, Stegger M, Larsen AR, et al. 2013. Whole genome sequencing identifies zoonotic transmission of MRSA isolates with the novel *mecA* homologue *mecC*. *EMBO molecular medicine* 5(4):509-515.
- Hartmeyer GN, Gahrn-Hansen B, Skov RL, Kolmos HJ. 2010. Pig-associated methicillin-resistant *Staphylococcus aureus*: family transmission and severe pneumonia in a newborn. *Scand J Infect Dis* 42(4):318-320.
- Khanna T, Friendship R, Dewey C, Weese JS. 2008. Methicillin resistant *Staphylococcus aureus* colonization in pigs and pig farmers. *Vet Microbiol* 128(3-4):298-303.
- Klevens RM, Morrison MA, Fridkin SK, Reingold A, Petit S, Gershman K, et al. 2006. Community-associated methicillin-resistant *Staphylococcus aureus* and healthcare risk factors. *Emerg Infect Dis* 12:1991-1993.
- Lee BY, Singh A, David MZ, Bartsch SM, Slayton RB, Huang SS, et al. 2013. The economic burden of community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA). *Clin Microbiol Infect* 19(6):528-536.
- Lewis HC, Molbak K, Reese C, Aarestrup FM, Selchau M, Sorum M, et al. 2008. Pigs as source of methicillin-resistant *Staphylococcus aureus* CC398 infections in humans, Denmark. *Emerg Infect Dis* 14(9):1383-1389.
- Lina G, Piemont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, et al. 1999. Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis* 29(5):1128-1132.
- McCarthy AJ, Witney AA, Gould KA, Moodley A, Guardabassi L, Voss A, et al. 2011. The distribution of mobile genetic elements (MGEs) in MRSA CC398 is associated with both host and country. *Genome biology and evolution* 3:1164-1174.

- Molla B, Byrne M, Abley M, Mathews J, Jackson CR, Fedorka-Cray P, et al. 2012. Epidemiology and genotypic characteristics of methicillin-resistant *Staphylococcus aureus* strains of porcine origin. *J Clin Microbiol* 50(11):3687-3693.
- Monaco M, Pedroni P, Sanchini A, Bonomini A, Indelicato A, Pantosti A. 2013. Livestock-associated methicillin-resistant *Staphylococcus aureus* responsible for human colonization and infection in an area of Italy with high density of pig farming. *BMC Infect Dis* 13(1):258.
- Monecke S, Coombs G, Shore AC, Coleman DC, Akpaka P, Borg M, et al. 2011. A field guide to pandemic, epidemic and sporadic clones of methicillin-resistant *Staphylococcus aureus*. *PLoS One* 6(4):e17936.
- Morrison MA, Hageman JC, Klevens RM. 2006. Case definition for community-associated methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 62(2):241.
- Oakes JM. 2006. Commentary: advancing neighbourhood-effects research--selection, inferential support, and structural confounding. *Int J Epidemiol* 35(3):643-647.
- Osadebe LU, Hanson B, Smith TC, Heimer R. 2013. Prevalence and characteristics of *Staphylococcus aureus* in Connecticut swine and swine farmers. *Zoonoses Public Health* 60(3):234-243.
- Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. 2012. *Staphylococcus aureus* CC398: host adaptation and emergence of methicillin resistance in livestock. *mBio* 3(1):e00305-00311.
- Rinsky JL, Nadimpalli M, Wing S, Hall D, Baron D, Price LB, et al. 2013. Livestock-associated methicillin and multidrug resistant *Staphylococcus aureus* is present among industrial, not antibiotic-free livestock operation workers in North Carolina. *PLoS One* 8(7):e67641.
- Salmenlinna S, Lyytikäinen O, Vainio A, Myllyniemi AL, Raulo S, Kanerva M, et al. 2010. Human cases of methicillin-resistant *Staphylococcus aureus* CC398, Finland. *Emerg Infect Dis* 16(10):1626-1629.
- Schulz J, Friese A, Klees S, Tenhagen BA, Fetsch A, Rosler U, et al. 2012. Longitudinal study of the contamination of air and of soil surfaces in the vicinity of pig barns by livestock-associated methicillin-resistant *Staphylococcus aureus*. *Appl Environ Microbiol* 78(16):5666-5671.

- Shopsin B, Gomez M, Montgomery SO, Smith DH, Waddington M, Dodge DE, et al. 1999. Evaluation of protein A gene polymorphic region DNA sequencing for typing of *Staphylococcus aureus* strains. *J Clin Microbiol* 37(11):3556-3563.
- Smith TC, Gebreyes WA, Abley MJ, Harper AL, Forshey BM, Male MJ, et al. 2013. Methicillin-Resistant *Staphylococcus aureus* in Pigs and Farm Workers on Conventional and Antibiotic-Free Swine Farms in the USA. *PLoS One* 8(5):e63704.
- Smith TC, Male MJ, Harper AL, Kroeger JS, Tinkler GP, Moritz ED, et al. 2009. Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine and swine workers. *PLoS One* 4(1):e4258.
- Sunde M, Tharaldsen H, Marstein L, Haugum M, Norstrom M, Jacobsen T, et al. 2011. Detection of methicillin-resistant *Staphylococcus aureus* sequence type 8 in pigs, production environment, and human beings. *J Vet Diagn Invest* 23(2):348-350.
- Sung JM, Lloyd DH, Lindsay JA. 2008. *Staphylococcus aureus* host specificity: comparative genomics of human versus animal isolates by multi-strain microarray. *Microbiology* 154(Pt 7):1949-1959.
- Tattevin P, Schwartz BS, Graber CJ, Volinski J, Bhukhen A, Bhukhen A, et al. 2012. Concurrent epidemics of skin and soft tissue infection and bloodstream infection due to community-associated methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis* 55(6):781-788.
- U.S. Food and Drug Administration. 2010. Summary report on antimicrobials sold or distributed for use in food-producing animals. Available: <http://www.fda.gov/downloads/ForIndustry/UserFees/AnimalDrugUserFeeActADUFA/UCM277657.pdf> [accessed 10 September 2013].
- van Cleef BA, Graveland H, Haenen AP, van de Giessen AW, Heederik D, Wagenaar JA, et al. 2011. Persistence of livestock-associated methicillin-resistant *Staphylococcus aureus* in field workers after short-term occupational exposure to pigs and veal calves. *J Clin Microbiol* 49(3):1030-1033.
- Verkaik NJ, Benard M, Boelens HA, de Vogel CP, Nouwen JL, Verbrugh HA, et al. 2011. Immune evasion cluster-positive bacteriophages are highly prevalent among human *Staphylococcus aureus* strains, but they are not essential in the first stages of nasal colonization. *Clin Microbiol Infect* 17(3):343-348.

- Vittinghoff E, Glidden D, Shiboski S, McCulloch C. 2005. Regression Methods in Biostatistics: Linear, Logistic, Survival, and Repeated Measures Models. New York, NY: Springer Science+Business Media, Inc.
- Voss A, Loeffen F, Bakker J, Klaassen C, Wulf M. 2005. Methicillin-resistant *Staphylococcus aureus* in pig farming. Emerg Infect Dis 11(12):1965-1966.
- Waters AE, Contente-Cuomo T, Buchhagen J, Liu CM, Watson L, Pearce K, et al. 2011. Multidrug-Resistant *Staphylococcus aureus* in US Meat and Poultry. Clin Infect Dis 52(10):1227-1230.
- Wulf MW, Verduin CM, van Nes A, Huijsdens X, Voss A. 2012. Infection and colonization with methicillin resistant *Staphylococcus aureus* ST398 versus other MRSA in an area with a high density of pig farms. Eur J Clin Microbiol Infect Dis 31(1):61-65.

Table 1. Patients identified and included in the present analysis according to MRSA subtype and exposure.

Group	High density livestock production in zip code	No high density livestock production in zip code	Total
CO-MRSA			
Patients included, n (%)	70 (42.2)	96 (57.8)	166
Total patients, n (%)	321 (32.5)	667 (67.5)	988
Weight ^a	4.59	6.95	
HO-MRSA			
Patients included, n (%)	11 (36.7)	19 (63.3)	30
Total patients, n (%)	54 (22.5)	186 (77.5)	240
Weight ^a	4.91	9.79	
Total patients included	81	115	196
Total patients identified	375	853	1128

Abbreviations: CO-MRSA, community-onset; HO, hospital-onset.

^aWeight calculated as the total number of patients identified in each stratum divided by the number of patients sampled from that stratum for the present analysis

Table 2. Onset location, Ridom *spa* types, and PVL presence by MRSA subgroup (n [%]).

Characteristic	CO-PVL-negative ^a , n = 23	<i>scn</i> -negative ^a , n = 15	All other MRSA, n = 162
Community-onset	23 (100) ^b	12 (80.0)	135 (83.3)
Clonal complex 5 or 5-like			
t002	9 (39.1) ^b	1 (6.7)	17 (10.5)
t010	2 (8.7) ^b	0	0
t045	0	1 (6.7)	1 (0.6)
t062	1 (4.3)	0	0
t088	3 (13.0) ^b	1 (6.7)	0
t105	1 (4.3)	1 (6.7)	2 (1.2)
t306	1 (4.3)	0	0
t6614	1 (4.3)	1 (6.7)	0
Other ^c	0	0	3 (1.8)
Clonal complex 8 or 8-like			
t008	0 ^b	7 (46.7)	115 (71.0)
t024	0	1 (6.7)	6 (3.7)
t068	0	0	2 (1.2)
t121	0	1 (6.7)	2 (1.2)
t622	0	0	3 (1.9)
Other ^d	0	0	7 (4.3)
Other clonal complex			
t125	1 (4.3)	1 (6.7)	0
t948	1 (4.3)	0	0
t9964	1 (4.3)	0	0
t11970	1 (4.3)	0	0
Other ^e	0	0	3 (1.9)
Novel ^f	1 (4.3)	0	1 (0.6)
PVL-positive	0 ^b	9 (60.0) ^b	140 (86.4)

Abbreviations: CO-PVL-negative, community-onset Panton-Valentine leukocidin-negative. Percentages do not add to one due to rounding.

^a4 patients were in both the CO-PVL-negative group and the *scn*-negative group. ^b*p*-value < 0.05 comparing two subgroups of interest to all other MRSA using Fisher's exact test. ^cOne each of t437, t539, and t85. ^dOne each of t064, t206, t211, t304, t681, t692, t1610. ^eOne each of t216, t316, t11971. ^fNovel *spa* type.

Table 3. Patient demographic and clinical characteristics of MRSA subgroups by swine livestock exposure quartile^a (data are n [%] unless otherwise indicated).

Characteristic	CO-PVL-negative, n = 23 ^b	<i>scn</i> -negative, n = 15 ^b	Other MRSA, n = 162	Swine quartile 1	Swine quartile 2	Swine quartile 3	Swine quartile 4
Male	15 (65.2)	11 (73.3)	85 (52.5)	28 (26.2)	25 (23.4)	24 (22.4)	30 (28.0)
Age at infection or visit, median (interquartile range), years	54.2 (32.3-75.2)	52.8 (14.6-75.1)	32.8 (14.4-55.2)	46.0 (17.3-62.4)	33.6 (22.5-54.7)	31.1 (9.8-51.7)	37.5 (13.6-59.1)
Race/ethnicity ^c							
Non-Hispanic White	22 (95.7)	15 (100)	155 (95.7)	44 (23.4)	49 (26.1)	48 (25.5)	47 (25.0)
Non-Hispanic Black	1 (4.4)	0	6 (3.7)	5 (71.4)	0	0	2 (28.6)
Other	0	0	1 (0.6)	0	0	1 (100)	0
Never-smoker	16 (69.6)	10 (66.7)	139 (85.8)	39 (23.9)	43 (26.4)	40 (24.5)	41 (25.2)
Season of onset							
Winter	7 (30.4)	3 (20.0)	27 (16.7)	12 (34.3)	7 (20.0)	7 (20.0)	9 (25.7)
Spring	5 (21.7)	4 (26.7)	37 (22.8)	16 (34.8)	13 (28.3)	9 (19.6)	8 (17.4)
Summer	7 (30.4)	5 (33.3)	39 (24.1)	10 (20.4)	16 (32.7)	12 (24.5)	11 (22.5)
Fall	4 (17.4)	3 (20.0)	59 (36.4)	11 (16.7)	13 (19.7)	21 (31.8)	21 (31.8)
Antibiotic prescription in prior year	5 (21.7)	6 (40.0)	65 (40.1)	21 (27.6)	20 (26.3)	15 (19.7)	20 (26.3)
Source							
Skin/soft tissue	19 (82.6)	11 (73.3)	141 (87.0)	36 (25.9)	39 (28.1)	28 (20.1)	36 (25.9)
Respiratory	0	2 (13.3)	15 (9.3)	6 (35.3)	4 (23.5)	4 (23.5)	3 (17.7)
Bone	2 (8.7)	1 (6.7)	2 (1.2)	1 (25.0)	0	2 (50.0)	1 (25.0)
Other	2 (8.7)	1 (6.7)	4 (2.5)	0	3 (42.9)	2 (28.6)	2 (28.6)
Antibiotic resistance ^d							
Clindamycin	12 (52.2)	2 (13.3)	32 (19.8)	10 (21.7)	11 (23.9)	13 (28.3)	12 (26.1)
Erythromycin	18 (78.3)	12 (80.0)	149 (92.0)	43 (24.4)	46 (26.1)	46 (26.1)	41 (23.3)
Tetracycline	4 (17.4)	1 (6.7)	4 (2.5)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)
TMP-SMZ	1 (4.4)	1 (6.7)	2 (1.2)	2 (66.7)	1 (33.3)	0	0
Multidrug ^e	14 (60.9)	3 (20.0)	33 (20.4)	12 (24.5)	11 (22.5)	14 (28.6)	12 (24.5)

Characteristic	CO-PVL-negative, n = 23 ^b	<i>scn</i> -negative, n = 15 ^b	Other MRSA, n = 162	Swine quartile 1	Swine quartile 2	Swine quartile 3	Swine quartile 4
Community type							
City	1 (4.4)	0	23 (14.2)	11 (45.8)	3 (12.5)	9 (37.5)	1 (4.2)
Borough	11 (47.8)	5 (33.3)	44 (27.2)	15 (25.9)	15 (25.9)	8 (13.8)	20 (34.5)
Township	11 (47.8)	10 (66.7)	95 (58.6)	23 (20.2)	31 (27.2)	32 (28.1)	28 (24.6)
Medical Assistance							
Never received	15 (65.2)	10 (66.7)	127 (78.4)	35 (23.3)	37 (24.7)	41 (27.3)	37 (24.7)
Location of onset							
Community	27 (100)	15 (83.3)	135 (83.3)	41 (24.7)	43 (25.9)	40 (24.1)	42 (25.3)
Hospital	0	3 (16.7)	27 (16.7)	8 (26.7)	6 (20.0)	9 (30.0)	7 (23.3)
Number of patients with high density livestock production in their residential zip code	9 (39.1)	7 (46.7)	67 (41.4)	1 (1.2)	17 (21.0)	25 (30.9)	38 (46.9)

Abbreviations: AEU, animal equivalent unit; CO-PVL-negative, community-onset Pantone-Valentine leukocidin-negative; TMP-SMZ, trimethoprim-sulfamethoxazole.

^aQuartile 1: swine livestock exposure <6.20 AEU/km²; quartile 2: 6.20-16.00 AEU/km²; quartile 3: 16.01-33.40 AEU/km²; quartile 4: >33.41 AEU/km². ^b4 patients were in both the CO-PVL-negative group and the *scn*-negative group. ^cRace/ethnicity was missing for 2 members of the all other isolates comparison group. ^dResistance to gentamicin was observed in one CO-PVL-negative isolate and intermediate resistance to rifampin was observed in one all other MRSA isolate. ^eResistance to ≥3 unique antimicrobials.

Table 4. Unadjusted and adjusted^a associations of dichotomous exposures with CO-PVL-negative MRSA and *scn*-negative MRSA compared to patients with all other MRSA.

Exposure	CO-PVL-negative ^b / All other MRSA (n)	CO-PVL-negative Unadjusted OR (95% CI)	CO-PVL-negative Adjusted OR ^c (95% CI)	<i>scn</i> -negative ^b / All other MRSA (n)	<i>scn</i> -negative Unadjusted OR (95% CI)
Livestock					
Swine					
Quartiles 1, 2, 3	12/124	1.0	1.0	12/125	1.0
Quartile 4 ^d	11/37	3.55 (1.41, 8.94)	4.24 (1.60, 11.25)	3/37	0.79 (0.20, 3.09)
Dairy/veal					
Quartiles 1, 2, 3	15/122	1.0	1.0	13/122	1.0
Quartile 4 ^e	8/40	1.57 (0.61, 4.05)	2.42 (0.85, 6.88)	2/40	0.47 (0.10, 2.24)
Treated crop field					
Swine					
Quartiles 1, 2, 3	15/121	1.0	1.0	14/121	1.0
Quartile 4 ^f	8/41	1.55 (0.59, 4.05)	2.38 (0.78, 7.28)	1/41	0.18 (0.02, 1.43)
Dairy/veal					
Quartiles 1, 2, 3	14/123	1.0	1.0	13/123	1.0
Quartile 4 ^g	9/39	2.28 (0.89, 5.83)	4.88 (1.40, 17.00)	2/39	0.46 (0.10, 2.20)

Abbreviations: AEU, animal equivalent unit; CO-PVL-negative, community-onset Panton-Valentine leukocidin-negative.

^aAdjusted estimates are not reported for the *scn*-negative group due to size limitations. ^b4 patients were in both the CO-PVL-negative group and the *scn*-negative group. ^cAdjusted for sex, age, ever-smoking status, antibiotic order in the 365 to 14 days prior to infection, season, and receipt of Medical Assistance. ^dQuartile 4: ≥ 33.41 AEU/km². ^eQuartile 4: ≥ 49.30 AEU/km². ^fQuartile 4: ≥ 43310 gal/m²/km². ^gQuartile 4: ≥ 118393 gal/m²/km².

Figure Legends

Figure 1. High-density livestock production and MRSA infection, Geisinger Health Care System, Pennsylvania. Abbreviations: PA, Pennsylvania; PVL, Panton-Valentine leukocidin.

Figure 2. Association of swine and dairy/veal seasonal crop field quartiles and livestock quartiles with CO-PVL-negative MRSA status from adjusted models. Adjusted for sex, age, ever-smoking status, antibiotic order in the 365 to 14 days prior to infection, season, and receipt of Medical Assistance. Abbreviations: CO-PVL-negative, community-onset Panton-Valentine leukocidin-negative. * p -value for linear trend ≤ 0.05 (quartiles included as a single variable with values 1, 2, 3, and 4).

Figure 1




Figure 2

