

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Acrolein Exposure in U.S. Tobacco Smokers and Non-Tobacco Users: NHANES 2005-2006

K. Udeni Alwis, B. Rey deCastro, John C. Morrow, and Benjamin C. Blount

Table of Contents

Table S1. Sample-weighted median (25th, 75th percentile) urinary 3HPMA concentration $\mu\text{g/g}$ creatinine by age, sex, and race/ethnicity among NHANES 2005-2006 tobacco smokers and non-tobacco users (sample size unweighted).

Table S2. Sample-weighted median (25th, 75th percentile) urinary CEMA concentration $\mu\text{g/g}$ creatinine by age, sex, and race/ethnicity among NHANES 2005-2006 tobacco smokers and non-tobacco users (sample size unweighted).

Table S3. Sample-weighted multiple regression slopes for urinary 3HPMA and CEMA concentrations ($\mu\text{g/L}$) among NHANES 2005-2006 non-tobacco users.

Table S4. Sample-weighted multiple regression slopes for urinary 3HPMA and CEMA concentrations ($\mu\text{g/L}$) among NHANES 2005-2006 exclusive cigarette smokers with serum cotinine > 10 ng/mL.