

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Adverse Associations of both Prenatal and Postnatal Exposure to Organophosphorous Pesticides with Infant Neurodevelopment in an Agricultural Area of Jiangsu Province, China

Ping Liu, Chunhua Wu, Xiuli Chang, Xiaojuan Qi, Minglan Zheng, and Zhijun Zhou

Table of Contents

Table S1. The distributions of maternal and child urinary DAP concentrations (nmol/L)

Table S2. DQ scores and proportions delayed according to the GDS ($n=310$)

Table S3. Association of maternal urinary DAP metabolites during pregnancy with fetal growth ($n=310$)

Table S4. Change in DQ scores in children tested at 2 years of age, for a 10-fold increase in maternal DAP concentrations during pregnancy (nmol/L)

Table S5. Change in DQ scores in children tested at 2 years of age, for a 10-fold increase in postnatal DAP concentrations (nmol/L)

Table S6. The comparison of the associations between DQ scores and total DAP concentrations measured in the prenatal vs. postnatal periods.