

**Note to Readers:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Traffic-Related Air Pollution, Noise at School, and Behavioral Problems in Barcelona Schoolchildren: A Cross-Sectional Study**

Joan Forns, Payam Dadvand, Maria Foraster, Mar Alvarez-Pedrerol, Ioar Rivas, Mònica López-Vicente, Elisabet Suades-Gonzalez, Raquel Garcia-Esteban, Mikel Esnaola, Marta Cirach, James Grellier, Xavier Basagaña, Xavier Querol, Mònica Guxens, Mark J. Nieuwenhuijsen, and Jordi Sunyer

#### **Table of Contents**

**Table S1.** Bivariate analysis between SDQ total difficulties score and ADHD-DSM-IV and sociodemographic characteristics at school level (n=39)

**Table S2.** Multilevel Variance Inflation Factors (MVIFS), parameter estimates, standard errors, and t-values for the multilevel model

**Table S3.** Effect modification of noise in the association between SDQ total difficulties score and ADHD symptomatology (from ADHD-DSM-IV) and TRAPs exposure at school ( $\mu\text{g}/\text{m}^3$ ) as continuous variable (based on an IQR increase)

**Table S4.** Adjusted mean ratio (aMRs and 95% CIs) of SDQ total difficulties score and ADHD symptomatology (from ADHD-DSM-IV) for black carbon exposure at school ( $\mu\text{g}/\text{m}^3$ ) and at home address ( $\mu\text{g}/\text{m}^3$ ) as continuous variable (based on an IQR increase)

**Table S5.** Adjusted mean ratio (aMRs 95% CIs) of SDQ total difficulties score and ADHD symptomatology (from ADHD-DSM-IV) for TRAPs exposure at school ( $\mu\text{g}/\text{m}^3$ ) and noise at school (dB) as continuous variable (based on an IQR increase) restricted to these children attending the same school since 3 years and before (n=2.192)

**Figure S1.** Adjusted mean ratio (aMRs 95% CIs) of SDQ subscales (emotion symptoms, conduct disorders, hyperactivity/inattention, peer relationship problems and prosocial

behavior) for indoor and outdoor TRAPs exposure at school ( $\mu\text{g}/\text{m}^3$ ) and noise at school (dB) as continuous variable (based on an IQR increase):

**Figure S2.** Adjusted mean ratio (aMRs 95% CIs) of ADHD-DSM-IV list (Inattention and hyperactivity) for indoor and outdoor TRAPs exposure at school ( $\mu\text{g}/\text{m}^3$ ) and noise at school (dB) as continuous variable (based on an IQR increase):