

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Ingested Nitrate and Breast Cancer in the Spanish Multicase-Control Study on Cancer (MCC-Spain)**

Nadia Espejo-Herrera, Esther Gracia-Lavedan, Marina Pollan, Nuria Aragonés, Elena Boldo, Beatriz Perez-Gomez, Jone M. Altzibar, Pilar Amiano, Ana Jiménez Zabala, Eva Ardanaz, Marcela Guevara, Antonio J. Molina, Juan Pablo Barrio, Ines Gómez-Acebo, Adonina Tardón, Rosana Peiró, M<sup>a</sup> Dolores Chirlaque, Margarita Palau, Montse Muñoz, Laia Font-Ribera, Gemma Castaño-Vinyals, Manolis Kogevinas, and Cristina M. Villanueva

#### **Table of Contents**

**Table S1.** General characteristics of analyzed compared to excluded population

**Table S2.** Average waterborne ingested nitrate in different exposure periods and breast cancer association (OR and 95%CI) by menopausal status

**Table S3.** Interaction of waterborne ingested nitrate with relevant covariables and breast cancer association (OR and 95%CI) among premenopausal women

**Table S4.** Waterborne ingested nitrate and breast cancer association (OR and 95%CI) stratified by other covariables among pre and postmenopausal women

**Table S5.** Breast cancer association (OR and 95%CI) with dietary ingested nitrate (mg/day) from different sources by menopausal status

**Figure S1.** Exposure-response relationship between waterborne nitrate intake (mg/day) from age 18 to 2 years before the interview and breast cancer association. Generalized additive models (GAMs) by study area