

**ehp**

**ENVIRONMENTAL  
HEALTH  
PERSPECTIVES**

ehponline.org

# **Intake to Production Ratio: A Measure of Exposure Intimacy for Manufactured Chemicals**

**William W. Nazaroff, Charles J. Weschler, John C. Little,  
Elaine A. Cohen Hubal**

**<http://dx.doi.org/10.1289/ehp.1204992>**

**Online 25 September 2012**



**NIEHS**  
National Institute of  
Environmental Health Sciences

National Institutes of Health  
U.S. Department of Health and Human Services

## **Intake to Production Ratio: A Measure of Exposure Intimacy for Manufactured Chemicals**

William W. Nazaroff<sup>1\*</sup>, Charles J. Weschler<sup>2,3</sup>, John C. Little<sup>4</sup>, Elaine A. Cohen Hubal<sup>5</sup>

<sup>1</sup> Department of Civil and Environmental Engineering, University of California, Berkeley, California, USA

<sup>2</sup> Environmental and Occupational Health Sciences Institute, University of Medicine & Dentistry of New Jersey & Rutgers University, Piscataway, New Jersey, USA

<sup>3</sup> International Centre for Indoor Environment & Energy, Technical University of Denmark, Lyngby, DK

<sup>4</sup> Department of Civil and Environmental Engineering, Virginia Tech, Blacksburg, Virginia, USA

<sup>5</sup> National Center for Computational Toxicology, Environmental Protection Agency, Research Triangle Park, North Carolina, USA

\* Corresponding author at: Department of Civil and Environmental Engineering, 425 Davis Hall, University of California, Berkeley CA 94720-1710 USA. Tel.: +1 510 642 1040. E-mail address: nazaroff@ce.berkeley.edu.

**Keywords:** bioavailability, bisphenol A, *p*-dichlorobenzene, pentachlorophenol, phthalate, screening, triclosan

**Short title:** Intake to production ratio

### **Acknowledgments**

This paper was developed as an outgrowth of two workshops organized by the Indoor Air Institute on the topic of semivolatile organic compounds in indoor environments. Funding for the workshops was provided to the Indoor Air Institute by the EPA Office of Research and Development under Order for Services No. EP-11-0-000172 and by a grant from the American Chemistry Council Long-Range Research Initiative. Funding in support of the workshops was also provided to the Building Ecology Research Group by the EPA Office of Air and Radiation, Indoor Environment Division, under CADMUS Prime Contract No. EP-D-08-096, Task Order Agreement No. IED069-BERG-1. We thank Hal Levin for his contributions to organizing the

workshops. The views expressed in this article are those of the authors and do not necessarily reflect the views or policies of the US EPA. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

### Competing Financial Interests Declaration

The authors have no competing financial interests.

### Abbreviations and Definitions

AM — arithmetic mean

$AM_{ue}$  — mean analyte level excreted in urine ( $\mu\text{g/g}$  creatinine)

BBzP — butyl benzyl phthalate

BPA — bisphenol A

$C_{50}, C_{75}, C_{90}, C_{95}$  — 50<sup>th</sup>, 75<sup>th</sup>, 90<sup>th</sup>, 95<sup>th</sup> percentile concentration values

CDC — Centers for Disease Control

CE — population mean creatinine excretion rate (1.54 g/d per person)

DCB — para-dichlorobenzene

DCP — 2,5-dichlorophenol

DEHP — di(2-ethylhexyl) phthalate

DEP — diethyl phthalate

DiBP — di(isobutyl) phthalate

DnBP — di(*n*-butyl) phthalate

EPA — (US) Environmental Protection Agency

$f_{ue}$  — urinary excretion fraction for an analyte

GM — geometric mean

GSD — geometric standard deviation

$I$  — daily average intake ( $\mu\text{g/d}$  per person)

iF — intake fraction

IPR — intake-to-production ratio

IUR — Inventory Update Reporting

MBzP — mono-benzyl phthalate

MECPP — mono-(2-ethyl-5-carboxypentyl) phthalate

MEHHP — mono-(2-ethyl-5-hydroxyhexyl) phthalate

MEHP — mono-2-ethylhexyl phthalate

MEOHP — mono-(2-ethyl-5-oxohexyl) phthalate

MEP — monoethyl phthalate

MiBP — mono-isobutyl phthalate

MnBP — mono-*n*-butyl phthalate

MP — methyl paraben

MW — molecular weight

NTP — National Toxicology Program

*P* — chemical production rate (g/d per person)

ppm — parts per million

$r^2$  — coefficient of determination

SVOC — semivolatile organic compound

US — United States

## Abstract

*Background:* Limited data are available to assess human exposure to thousands of chemicals currently in commerce. Information that relates human intake of a chemical to its production and use can help inform understanding of mechanisms and pathways that control exposure and support efforts to protect public health.

*Objectives:* We introduce the intake-to-production ratio (IPR) as an economy-wide quantitative indicator of the extent to which chemical production results in human exposure.

*Methods:* The IPR is evaluated as the ratio of two terms: aggregate rate of chemical uptake in a human population (inferred based on urinary excretion data) divided by the rate at which that chemical is produced in or imported into that population's economy. We use biomonitoring data from the US Centers for Disease Control along with chemical manufacturing data reported by the US Environmental Protection Agency as well as other published data to estimate the IPR for nine chemicals in the United States. Results are reported in units of parts per million (ppm), where one ppm indicates one gram of chemical uptake for every million grams of economy-wide use.

*Results:* Estimated IPR values for the studied compounds span many orders of magnitude from a low of 0.6 ppm for bisphenol A to a high of greater than 180,000 ppm for methyl paraben. Intermediate results are obtained for five phthalates and two chlorinated aromatic compounds: 120 ppm for butyl benzyl phthalate, 670 ppm for di(2-ethylhexyl) phthalate, 760 ppm for di(*n*-butyl) phthalate, 1040 ppm for para-dichlorobenzene, 6800 ppm for di(isobutyl) phthalate, 7700 ppm for diethyl phthalate, and in the range 8000-24,000 ppm for triclosan.

*Conclusion:* The IPR provides an aggregate metric of exposure intensity for characterizing population-level exposure to synthesized chemicals.

## Introduction

Exposures to manufactured chemicals can pose health risks (Collins et al. 2008; Rudén and Hansson 2010). Effective management of these risks requires knowledge not only about chemical toxicities but also about human exposures. Tracing exposure pathways from sources to receptors is challenging. Innovative approaches are needed to leverage exposure information that is available for only a small subset of the thousands of chemicals in commerce (Egeghy et al. 2012). Multimedia fate and transport models have been combined with metrics such as the intake fraction for large-scale assessments (Bennett et al. 2002a). Proximity between chemical releases and exposed populations is emerging as a parameter of key influence for assessing human intake (Manneh et al. 2010). Methods for characterizing the degree of *exposure intimacy* of human populations to manufactured chemicals could improve understanding of exposure pathways and assist in studies that aim to understand biologically relevant exposures (i.e., those associated with a disease) (Cohen Hubal 2009).

The existence of two national data sets in the United States creates an opportunity to compute a new aggregate exposure metric for certain chemicals, including several species that are of current environmental health interest. The Chemical Data Reporting system (formerly known as Inventory Update Reporting), maintained by the US Environmental Protection Agency, collects and manages data concerning the manufacture and importation of industrial chemicals (USEPA 2012). The National Report on Human Exposure to Environmental Chemicals, a project of the US Centers for Disease Control, employs biomonitoring to characterize exposure of the US population to environmental chemicals (CDC 2012). We use data from these two systems to compute a dimensionless measure of exposure intensity, the “intake to production ratio” (IPR). In this paper, IPR values are estimated for the US population for nine organic chemicals: methyl

paraben, triclosan, five phthalates, para-dichlorobenzene and bisphenol A. The resulting IPR values span a remarkably broad range — more than five orders of magnitude for these nine chemicals. We explore the significance of the results, the limitations of the concept, and how the IPR might contribute to an improved understanding of population-level exposure to manufactured chemicals.

## Methods

Aggregate exposure metrics should provide useful information about exposure, should be applicable to many chemicals, and should be economical to implement. Population-level exposure metrics are required to extend molecular and cellular-level insights to inform risk assessment and decision-making (Ankley et al. 2010). One such metric is introduced here: the intake-to-production ratio (IPR). We define the IPR to be the aggregate rate of chemical uptake in a human population (estimated here based on urinary excretion data) divided by the rate at which that chemical is produced in or imported into that population's economy. The IPR is a dimensionless ratio that we report here using a parts-per-million ( $\text{ppm} = 10^{-6}$ ) modifier. An IPR of 1.0 ppm indicates that, on average, for every million grams of the chemical entering the economy, one gram enters the aggregate human population served by that economy. The IPR does not reflect the distribution of exposures within a population: one gram of uptake by one individual yields the same IPR as one milligram of uptake by each of one thousand individuals.

The IPR is intended as a magnitude estimate that characterizes the degree of *exposure intimacy* between a chemical and a population. Knowledge about exposure pathways is not required for estimating IPR values. However, knowledge of the IPR can help inform investigations that aim to elucidate important exposure pathways. While the IPR is simple in concept, data availability is a limiting constraint for its use today. In principle, the IPR might be

developed and applied for many manufactured chemicals and for a variety of economy-based populations. In the current paper, we emphasize chemicals that are commonly used in consumer products, in building materials or furnishings, or in other products that might lead to widely distributed yet close contact with populations. One broad class of interest is semivolatile organic compounds (SVOCs) (Weschler and Nazaroff 2008), many species of which are commonly found indoors (Rudel et al. 2003). Some SVOCs have associated health concerns owing to known or suspected endocrine-disrupting and/or carcinogenic properties (Rudel et al. 2003; Rudel and Perovich 2009). The specific demonstrations in this paper for contemporary exposures are limited to a set of chemicals that do not bioaccumulate and for which urinary excretion, either of the primary compound (whether conjugated or not) or of one or more metabolites, is the dominant biological loss pathway. However, these conditions are not inherent limitations for the IPR metric. We illustrate this point in the discussion by estimating an IPR for pentachlorophenol from an historic interpretation incorporating multimedia fate and transport modeling as a basis for estimating population intake.

### *Production Data*

The specific demonstration in this paper focuses on the general US population. Production data were mainly obtained from the US Environmental Protection Agency's Inventory Update Reporting and Chemical Data Reporting system (USEPA 2012), whose purpose is "to collect quality screening-level, exposure-related information on chemical substances and to make that information available for use by the EPA and, to the extent possible, to the public." Until recently, the system was known as Inventory Update Reporting, and the IUR acronym is used here. The IUR data have been historically reported at four-year intervals, starting in 1986. The most recent IUR data, for 2006, summarize production and imports for calendar year 2005.

Under the reporting rules, “manufacturers and importers producing 25,000 pounds or more of a reportable chemical substance must report the identity of the chemical substance and basic manufacturing information.” In the 2006 report, the IUR compiled data reported by 1541 companies covering 3827 sites and 6200 chemicals (USEPA 2008). It is possible that total production and importation rates are higher than used in this analysis because of the exclusion of operations smaller than the reporting limit. That cause of bias only appears potentially significant for methyl paraben (reported production of less than 0.5 million pounds per year) and di(isobutyl) phthalate (0.5-1 million pounds per year). For all other chemicals, the cumulative production rates exceed by at least two orders of magnitude the reporting limit.

For phthalates, we obtained production data directly from the USEPA IUR. For bisphenol A (BPA), only a lower bound was reported by IUR (specifying that greater than 1 billion lbs per year was produced in each report from 1990 to 2006). A summary risk appraisal for BPA provided a more precise value (NTP 2007). Also, the most recent IUR does not contain data for triclosan. Instead, we used the most recently reported national data, which are from the 1998 IUR (Fang et al. 2010). For the period 1986-1998, triclosan production increased substantially. Allowing that the growth might have continued, we treat the 1998 production rate as a lower bound and estimate an upper bound on production in 2002 by fitting an exponential growth rate to the data for 1986-1998.

In cases where production only is reported to occur within a specified range, we used the geometric mean of the upper and lower bounds of the range as the central best estimate of total production. Because many of the ranges are broad, uncertainty in production dominates total uncertainty of IPR estimates for most chemicals. For chemicals with multiplicatively large ranges, the production uncertainty is estimated as the multiplicative factor that links the central

best estimate to the upper end of the reported range. Details are reported in the Supplemental Material (see Supplemental Material, pp. 1–2).

To estimate the per-capita daily production, the central estimate aggregate annual production was converted to grams per person per day by means of unit conversion and by dividing by the estimated total US population, using US census data for the appropriate year. Detailed results showing per-capita production rate estimates for the period 1986-2006 are presented in Supplemental Material, Tables S1 and S2.

#### *Intake Estimates Based on Urinary Excretion Data*

Biomonitoring-based exposure information is being gathered by the Centers for Disease Control (CDC) for the US population (CDC 2012). The current analyte list comprises 212 chemicals. We used data from the February 2012 update. For the IPR estimates reported here, we have matched – to the extent possible – the timing of urinary excretion sampling with the timing of production data. We used the CDC data for the US population aged 6 and older to estimate arithmetic mean levels of target analytes in urine, expressed in units of  $\mu\text{g}$  of analyte per g of creatinine.

To elaborate, our goal in using the CDC biomonitoring data is to estimate the cumulative excretion rate of the chemical of interest or its key metabolite(s) summed over the entire US population. For each sampling period, the CDC reports geometric means as well as the 50<sup>th</sup>, 75<sup>th</sup>, 90<sup>th</sup>, and 95<sup>th</sup> percentile results for the US population  $\geq 6$  years of age. For the compounds of interest here, the distributions of analyte concentrations in urine exhibit substantial skew. So, while the geometric mean represents well the central tendency of individual excretion rates, it is substantially below the population mean value. To estimate a mean from the tabulated data, we fit lognormal distributions to the four reported percentile values to obtain best-fit geometric

means (GM) and geometric standard deviations (GSD). The fitting was done as a least-squares linear regression of the logarithm of the measured value against the  $z$ -score that corresponds to the indicated percentile value. An  $r^2$  value for the regression was computed as an indicator of goodness of fit of the lognormal form to the 50<sup>th</sup>-95<sup>th</sup> percentile results. Given the fitted GM and GSD, the arithmetic mean (AM) was computed from equation [1], which holds for lognormal distributions:

$$AM = GM \times \exp\left(\frac{[\ln(GSD)]^2}{2}\right) \quad [1]$$

Creatinine-corrected excretion data for the whole sampled population were used for the time period that corresponded most closely to the year for which production data were available. Table 1 summarizes the resulting estimates of arithmetic mean excretion rates of target analytes. The ratio of AM to GM for most of the chemicals varies over a range from 1.5 (MiBP) to 3.2 (MEP). However, for 2,5-dichlorophenol and for triclosan, the ratios are much higher, 11.8 and 13.5, respectively, reflecting the very large ranges of levels in urine, with the 95<sup>th</sup> percentile concentration exceeding the medians by factors of  $\sim 40$  in each case.

We augmented the primary assessment of population intake rates for IPR calculations by repeating the analysis for all years for which analyte excretion data are available. A goal of that exercise was to explore the extent to which the per-capita average excretion rates have been varying with time. The earliest data are from 1999-2000 and for some analytes, monitoring data are available on a biennial basis for five consecutive periods through 2008. For other analytes, biomonitoring began more recently, as in the case of methyl paraben for which the first data are for 2005-2006. The full analysis of per-capita average intake rates based on urinary excretion data is reported in Supplemental Material, Table S3.

We estimated the population average intake rates of parent compounds from the arithmetic mean analyte levels in urine, taking into account empirical data on fractional excretion of the analytes following ingestion exposure and also accounting for the relative molecular weights of the analytes and the parent compounds. For three of the phthalates (BBzP, DEHP, and DnBP) and their associated monoester metabolites, we relied on the summary data for molar excretion rates as reviewed by Wittassek et al. (2011). For the other two phthalates (DiBP and DEP), lacking direct experimental data, we assumed that the fractional excretion of the associated metabolites was the same as for DnBP. For BPA, we assumed complete urinary excretion, based on the report by Dekant and Völkel (2008). For *p*-dichlorobenzene, we also assumed complete excretion of the associated metabolite 2,5-dichlorophenol (DCP) on the basis of an exposure study of Yoshida et al. (2002). For triclosan, we assumed 57% urinary excretion, based on the work of Sandborgh-Englund et al. (2006). We were unable to locate any published information on the fractional urinary excretion of methyl paraben and so we have conducted the analysis on the basis that the upper bound must be complete excretion. Table 2 summarizes the fractional urinary excretion estimates as well as the molecular weights used in the analysis.

Two other data elements are required to complete the analysis: creatinine clearance rates and body masses. For the former, we used 18 mg/kg per day for females and 23 mg/kg per day for males, following similar calculations by Fromme et al. (2007) and Kohn et al. (2000). For body masses, we computed US population average values by applying the age-specific data for 2002 reported in Ogden et al. (2004). The mean was estimated separately for males and for females as the evenly weighted mean for all ages between 6 and 69. The results were 69.2 kg for females and 79.5 kg for males. Combining these factors, we estimated an average urinary creatinine

clearance rate of 1.25 g/d for females and 1.83 g/d for males, or 1.54 g/d as the combined population average.

For the case of a single analyte resulting from a given parent compound, the population-mean daily intake of the parent compound is estimated from equation [2] (Kohn et al. 2000):

$$I = \frac{AM_{ue} \times CE}{f_{ue}} \times \frac{MW_p}{MW_a} \quad [2]$$

where  $I$  is the daily average intake ( $\mu\text{g/d}$  per person),  $AM_{ue}$  is the arithmetic mean level excreted in urine ( $\mu\text{g/g}$  creatinine, Table 1),  $CE$  is the population mean creatinine excretion rate (1.54 g/d per person),  $f_{ue}$  is the urinary excretion fraction (Table 2), and  $MW_p$  and  $MW_a$  are the respective molecular weights of the parent compound and of the urinary analyte (Table 2).

For DEHP, with four analytes, a modified approach was used to obtain an overall estimate:

$$I_{DEHP} = \left( \sum \frac{AM_{ue}}{MW_a} \right) \times \frac{CE \times MW_{DEHP}}{\sum f_{ue}} \quad [3]$$

In applying equation [3], the sums were separately evaluated for the four primary metabolites and the results substituted into the equation to obtain an estimate of DEHP intake.

In using urinary metabolites as quantitative markers of intake, it is important to recognize the possibility that certain metabolites may have multiple potential parent compounds. We have relied on information from the biomonitoring report (CDC 2012), which indicates that the metabolites considered in this study have a single parent compound with one exception. Mono-*(n*-butyl) phthalate (MnBP) is a metabolite of both di-*(n*-butyl) phthalate (DnBP) and butylbenzyl phthalate (BBzP). In the analysis presented here, we have assigned the MnBP data entirely to DnBP intake on the basis of two lines of evidence. First, Anderson et al. (2001) have shown that BBzP is converted to mono-benzyl phthalate (MBzP) with a conversion rate that is more than

10× as large as for MnBP. Second, the average urinary level of MnBP is roughly twice the level of MBzP. Consequently, the error in assuming that all MnBP is attributable to DnBP intake is no larger than 5%, which is negligible in the current context.

Variability in the biomonitoring data provides indicators of uncertainty in the intake estimates. For most chemicals, relative to the uncertainty in published production data, intake uncertainty is small. For the purpose of this paper, we estimate the percentage uncertainty in daily per capita intake as the larger of (a) the square root of the ratio of maximum to minimum in the reported range of geometric mean excretion rates as reported by CDC (2012); and (b) the relative standard deviation in year-to-year arithmetic mean excretion rates. These values are then combined in quadrature with the uncertainty estimates for production data to obtain an overall estimated uncertainty in the IPR (see Supplemental Material, Equation S1). For triclosan and methyl paraben, only lower and upper bounds are determined for the IPR as the data are not well suited for evaluating best point estimates. Details of the uncertainty analysis are reported in the Supplemental Material (see Supplemental Material, pp. 4–6 and Table S4).

## **Results and Discussion**

We estimated IPR values for recent conditions in the United States for nine common manufactured chemicals (Table 3 and Figure 1). A striking feature of the results is the very broad range of IPR values obtained, spanning more than five orders of magnitude from bisphenol A at 0.6 ppm to methyl paraben at >180,000 ppm. The probable explanation for this broad range is that different uses of these chemicals lead to very different degrees of opportunity for exposure, or different levels of exposure intimacy, as expressed through intake into the human body. For example, a main use of bisphenol A is as a starting material in the manufacture of polycarbonate plastics and epoxy resins, accounting for 72% and 21%, respectively of the BPA consumed in

the US in 2003 (NTP 2007). In these products, most of the original BPA is transformed into the polymeric fabric of the material. Small amounts of monomer may be released after manufacture either owing to the presence of residual monomer or as the polycarbonate or epoxy degrades (Kang et al. 2006). Consequently, one expects that human exposure to BPA would be limited to a small fraction of the original BPA used to produce the polycarbonate plastic or the epoxy resins. The end products made from BPA also have widely variable human exposure potential, from high potential in food-can linings to low potential in auto-body parts.

Conversely, the relatively high IPR for triclosan, an antibacterial agent, very likely reflects its major use in personal care products. Many brands of toothpaste, antibacterial soaps and mouthwashes use triclosan as an active ingredient. Exposures as a consequence of ingestion (toothpaste or mouthwash) or dermal permeation (soaps) would be relatively direct. The high IPR for triclosan suggests that these uses lead to intakes that are a meaningful fraction of the total amount of triclosan incorporated into personal care products.

Similarly, methyl paraben is “widely used as a preservative in food, cosmetic and pharmaceutical products” (Soni et al. 2005). Its use in food and in ingested pharmaceuticals would be associated with an expected IPR approaching unity, i.e. 1,000,000 ppm. As a component of cosmetics, there is opportunity for intake via at least three pathways: transdermal permeation, inhalation of the semivolatile compound, and incidental ingestion. Methyl paraben has some natural sources; however, Soni et al. (2005) report that, “paraben intake from natural sources is negligible.”

As illustrated by these cases, the IPR can vary over several orders of magnitude, with the degree of variability depending on the ways in which the chemical is used. A chemical primarily used in products that are ingested, i.e. as a food additive or in pharmaceuticals should have an

IPR approaching 1,000,000 ppm. To elaborate, if all of a manufactured chemical is added to food that is ingested, then the cumulative intake in a human population would be the same as the cumulative production of the chemical. The IPR for a chemical mainly used externally in personal care products might be somewhat smaller, perhaps on the order of 10,000 to 100,000 ppm. The IPR for a chemical that is chiefly used as a constituent of sprays or pesticides applied indoors might be in the range 100 to 10,000 ppm. The IPR for semivolatile organic compounds present as additives in polymeric products, such as plasticizers or flame retardants, might be on the order of 1 to 1000 ppm. We caution that these are rough estimates that are guided in part by the limited IPR investigations reported here as well as prior studies of intake fractions for indoor pollutant emissions (Nazaroff 2008) and research on the dynamic behavior of indoor SVOCs, as reviewed by Weschler and Nazaroff (2008). Further studies that better characterize IPRs, not only in these but also in other use categories, appear warranted.

Among possible applications, we envision that the IPR might be a useful metric for preliminary screening for some chemicals in the following manner. There are on the order of 30,000 commercial chemicals to prioritize in terms of public exposure. Where data permit, the reported or intended use of a compound could inform its assignment to a specific IPR range or bin. Combining available information on annual production with the chemical's assigned IPR range would yield a preliminary estimate of the anticipated range for the population's intake rate for that chemical. Such zero-order screening would allow those compounds with the highest anticipated intake rates to be identified, which could be of value in prioritizing chemicals for more detailed health risk assessments.

As detailed in the Supplemental Material (Table S4), the uncertainty in the estimates reported here is dominated by the lack of precision in the production data. For five of the compounds

studied here (BPA, DEHP, DnBP, DCB, and DEP), production uncertainty contributes a multiplicative uncertainty of 2.2-2.3×. For BBzP and DiBP, the production uncertainty is smaller owing to a narrower reporting range, and particularly in the case of DiBP, the intake uncertainty makes a meaningful contribution to the overall uncertainty estimate. For triclosan and for methyl paraben, we are only able to estimate ranges that bracket the expected IPR values. For triclosan, the range spans a factor of three, accounting for the uncertainty in production in 2002 based on data reported for the period 1986 to 1998. For methyl paraben, the production rate is expressed as an upper bound. The intake estimate represents a lower bound based on the assumption that urinary excretion is complete. The consequence is an IPR estimate (180,000 ppm) that represents a lower bound. The upper bound cannot exceed 1,000,000 ppm, so the IPR for methyl paraben can be bracketed within a range that spans a factor of five. Importantly, the uncertainties, while moderately high, are small compared to the differences in IPR estimates for these chemicals (Figure 1). Consequently, we believe that there is useful information in the IPR estimates despite the moderately high uncertainties.

Because IPR is defined as the ratio of two rates, temporal aspects merit consideration. The IPR is best suited for characterizing exposure to chemicals that move fairly rapidly from manufacturing to human intake. The IPR would also be well suited for cases in which the production rates and intake rates are relatively stable over time. Consider, for example, DEHP as a plasticizer. To the extent that it is used in products such as vinyl flooring, DEHP would be expected to be persistent, contributing to exposure for years or decades after manufacture. Intake in any given year would result not only from the DEHP manufactured in that year but also from DEHP manufactured in prior years that is present in products that are still in use. Interestingly, though, a study of German university students showed strong year-by-year tracking between

economy-wide production data and estimated intake rates for the period 1988-2004 (Helm 2007). Wittassek et al. (2011) have reported an analogous finding for di(*n*-butyl) phthalate (DnBP). We have made estimates of the IPR values for these two chemicals in Germany, based on data extracted from Figure 3 of Wittassek et al. (2011). The results indicate IPR values in the range 30-60 ppm for DEHP and 500-800 ppm for DnBP. The DEHP result is about an order of magnitude lower than our estimate for the US population, potentially reflecting earlier restrictions on the use of DEHP in Europe than in the US (Europa 1999). The DnBP results are about the same for the US and German populations.

There are sufficient biomonitoring and chemical production data in the US to separately estimate IPR values for 2001-2002 and for 2005-2006 for the five phthalates considered. We report the results of that assessment in the Supplemental Material (Table S5). Overall, the results show reasonable consistency over time: the average deviation across the five species between the two time periods is 34%.

A general challenge in using biomonitoring data for IPR assessments is to relate measured levels in biological fluids to rates of intake. The IPR concept is not restricted to chemicals excreted in urine after human intake. To the extent that levels in blood, breath or other media can be used to infer intake, these metrics, sampled in representative subpopulations, could be used to estimate IPRs. For the chemicals considered here, the rapid urinary excretion of the primary chemicals and/or their metabolites enables a relatively simple quantitative description of this relationship. For bioaccumulative compounds, levels measured in body fluids can be translated into intake estimates through pharmacokinetic models (e.g., Hays and Aylward 2012; Lorber 2008; Lorber and Egeghy 2011).

It is also possible to estimate IPR values from historic data. Consider the example of pentachlorophenol, a wood preservative and biocide widely used in the United States prior to regulatory changes in the mid 1980s. Data presented by Hattemer-Frey and Travis (1989) support an IPR estimate for pentachlorophenol in the US population during the late 1980s of 60 ppm, based on annual production of 23 million kg, estimated average intake of 16  $\mu\text{g}/\text{d}$  per person, and a US population at the time of 240 million. In the context of the results presented in Figure 1, this estimate suggests a low to intermediate level of exposure intimacy for the historic uses of pentachlorophenol, higher than current values for bisphenol A and lower than estimated for the phthalates.

The IPR complements another aggregate exposure metric, the intake fraction (iF) (Bennett et al. 2002b). Both are dimensionless ratios quantifying exposure intensity. The numerator in each case represents an intake summed over an exposed population. However, the denominators are distinct, with the denominator of the iF quantifying an environmental release whereas the IPR denominator measures the total quantity of a chemical manufactured and/or imported. The IPR is necessarily applicable only over large (economy-wide) populations whereas the intake fraction can be applied more flexibly across widely variable population sizes. The iF is well suited for characterizing exposure to chemicals that are created incidentally, such as combustion byproducts. The IPR is well suited for characterizing exposure to chemicals that are synthesized and used either neat (e.g., certain pesticides) or as product constituents, especially those that partition significantly among multiple phases.

## Conclusions

Novel insights concerning public exposures to manufactured chemicals can be gained from further development and application of the intake-to-production ratio metric. For example,

collections of IPRs for manufactured chemicals with a range of properties and uses could inform assessment of potential exposures for new or data-poor chemicals. Statistical modeling approaches could be used to capture aggregate information embedded in the IPRs and these results could be combined with predictions from mechanistic models to improve estimates of potential exposure. When combined with rapid toxicity screening tools (Judson et al. 2010; Judson et al. 2011; Rotroff et al. 2010), methods to rapidly estimate potential for human exposure could contribute to improved health-risk-based prioritization of a wide range of chemicals of concern (Cohen Hubal et al. 2010; Egeghy et al. 2012). The chemical/source combinations of greatest concern can then be more comprehensively investigated. Because our understanding of the vast number of sources and wide range of exposure pathways remains rudimentary for many manufactured chemicals, it is clear that an iterative approach will be required. As the mechanisms and pathways controlling the source-to-dose continuum are more clearly elucidated, improved methods to obtain rapid screening-level exposure estimates should emerge.

## References

- Anderson WAC, Castle L, Scotter MJ, Massey RC, Springall C. 2001. A biomarker approach to measuring human dietary exposure to certain phthalate diesters. *Food Addit Contam A* 18:1068-1074.
- Ankley GT, Bennett RS, Erickson RJ, Hoff DJ, Hornung MW, Johnson RD, et al. 2010. Adverse outcome pathways: a conceptual framework to support ecotoxicology research and risk assessment. *Environ Toxicol Chem* 29:730–741.
- Bennett DH, Margni MD, McKone TE, Jolliet O. 2002a. Intake fraction for multimedia pollutants: A tool for life cycle analysis and comparative risk assessment. *Risk Anal* 22:905-918.
- Bennett DH, McKone TE, Evans JS, Nazaroff WW, Margni MD, Jolliet O, et al. 2002b. Defining intake fraction. *Environ Sci Technol* 36:206A-211A.
- CDC [Centers for Disease Control and Prevention]. 2012. National Report on Human Exposure to Environmental Chemicals. Available: <http://www.cdc.gov/exposurereport> [accessed 24 May 2012].
- Cohen Hubal EA. 2009. Biologically relevant exposure science for 21<sup>st</sup> century toxicity testing. *Tox Sci* 111:226-232.
- Cohen Hubal EA, Richard A, Aylward L, Edwards S, Gallagher J, Goldsmith M-R, et al. 2010. Advancing exposure characterization for chemical evaluation and risk assessment. *J Toxicol Env Heal B* 13:299-313.
- Collins FS, Gray GM, Bucher JR. 2008. Transforming environmental health protection. *Science* 319:906-907.
- Dekant W, Völkel W. 2008. Human exposure to bisphenol A by biomonitoring: Methods, results and assessment of environmental exposures. *Toxicol Appl Pharm* 228:114-134.
- Egghy PP, Judson R, Gangwal S, Mosher S, Smith D, Vail J, et al. 2012. The exposure data landscape for manufactured chemicals. *Sci Tot Environ* 414:159-166.

- Europa Press Release. 1999. Ban of phthalates in childcare articles and toys. Available: <http://europa.eu/rapid/pressReleasesAction.do?reference=IP/99/829&format=HTML&aged=1&language=EN&guiLanguage=en> [accessed 11 January 2012].
- Fang J-L, Stingley RL, Beland FA, Harrouk W, Lumpkins DL, Howard P. 2010. Occurrence, efficacy, metabolism, and toxicity of triclosan. *J Environ Sci Heal C* 28: 147-171.
- Fromme H, Gruber L, Schlummer M, Wolz G, Böhmer S, Angerer J, et al. 2007. Intake of phthalates and di(2-ethylhexyl)adipate: Results of the Integrated Exposure Assessment Survey based on duplicate diet samples and biomonitoring data. *Environ Int* 33:1012-1020.
- Hattemer-Frey HA, Travis CC. 1989. Pentachlorophenol: Environmental partitioning and human exposure. *Arch Environ Contam Toxicol* 18:482-489.
- Hays SM, Aylward LL. 2012. Interpreting human biomonitoring data in a public health risk context using Biomonitoring Equivalents. *Int J Hyg Envir Heal* 215:145-148.
- Helm D. 2007. Correlation between production amounts of DEHP and daily intake. *Sci Total Environ* 388:389-391.
- Judson RS, Houck KA, Kavlock RJ, Knudsen TB, Martin MT, Mortensen HM, et al. 2010. In vitro screening of environmental chemicals for targeted testing prioritization: the ToxCast project. *Environ Health Persp* 118:485-492.
- Judson RS, Kavlock RJ, Setzer RW, Cohen Hubal EA, Martin MT, Knudsen TB, et al. 2011. Estimating toxicity-related biological pathway altering doses for high-throughput chemical risk assessment. *Chem Res Toxicol* 24:451-462.
- Kang J-H, Kondo F, Katayama Y. 2006. Human exposure to bisphenol A. *Toxicol* 226:79-89.
- Koch HM, Bolt HM, Preuss R, Angerer J. 2005. New metabolites of di(2-ethylhexyl)phthalate (DEHP) in human urine and serum after single oral doses of deuterium-labelled DEHP. *Arch Toxicol* 79:367-376.
- Kohn MC, Parham F, Masten SA, Portier CJ, Shelby MD, Brock JW, et al. 2000. Human exposure estimates for phthalates. *Environ Health Persp* 108:A440-A442.
- Lorber M. 2008. Exposure of Americans to polybrominated diphenyl ethers. *J Expo Sci Env Epid* 18:2-19.

- Lorber M, Egeghy P. 2011. Simple intake and pharmacokinetic modeling to characterize exposure of Americans to perfluorooctanoic acid, PFOA. *Environ Sci Technol* 45:8006-8014.
- Manneh R, Margni M, Deschenes L. 2010. Spatial variability of intake fractions for Canadian emission scenarios: A comparison between three resolution scales. *Environ Sci Technol* 44:4217-4224.
- Nazaroff WW. 2008. Inhalation intake fraction of pollutants from episodic indoor emissions. *Build Environ* 43:269-277.
- NTP (National Toxicology Program). 2007. NTP-CERHR Expert Panel Report on the Reproductive and Developmental Toxicity of Bisphenol A. Report NTP-CERHR-BPA-07. Available: [cerhr.niehs.nih.gov/evals/bisphenol/BPAFinalEPVF112607.pdf](http://cerhr.niehs.nih.gov/evals/bisphenol/BPAFinalEPVF112607.pdf) [accessed 19 January 2012].
- Ogden CL, Fryar CD, Carroll MD, Flegal KM. 2004. Mean body weight, height, and body mass index, United States 1960-2002. *Advance Data from Vital and Health Statistics* Number 347. Available: [www.cdc.gov/nchs/data/ad/ad347.pdf](http://www.cdc.gov/nchs/data/ad/ad347.pdf) [accessed 19 January 2012].
- Rotroff DM, Wetmore BA, Dix DJ, Ferguson SS, Clewell HJ, Houck KA, et al. 2010. Incorporating human dosimetry and exposure into high-throughput in vitro toxicity screening. *Toxicol Sci* 117:348-358.
- Rudel RA, Camann DE, Spengler JD, Korn LR, Brody JG. 2003. Phthalates, alkylphenols, pesticides, polybrominated diphenyl ethers, and other endocrine-disrupting compounds in indoor air and dust. *Environ Sci Technol* 37:4543-4553.
- Rudel RA, Perovich LJ. 2009. Endocrine disrupting chemicals in indoor and outdoor air. *Atmos Environ* 43:170-181.
- Rudén C, Hansson SO. 2010. Registration, Evaluation, and Authorization of Chemicals (REACH) is but the first step – How far will it take us? Six further steps to improve the European chemicals legislation. *Environ Health Persp* 118:6-10.
- Sandborgh-Englund G, Adolfsson-Erici M, Odham G, Ekstrand J. 2006. Pharmacokinetics of triclosan following oral ingestion in humans. *J Toxicol Env Heal A* 69:1861-1873.

- Soni MG, Carabin IG, Burdock GA. 2005. Safety assessment of esters of *p*-hydroxybenzoic acid (parabens). *Food Chem Toxicol* 43:985-1015.
- USEPA (US Environmental Protection Agency). 2008. 2006 Inventory Update Reporting: Data Summary. Available: [http://www.epa.gov/iur/pubs/2006\\_data\\_summary.pdf](http://www.epa.gov/iur/pubs/2006_data_summary.pdf) [accessed 19 January 2012].
- USEPA (US Environmental Protection Agency). 2012. Chemical Data Reporting/Inventory Update Reporting. Available: <http://www.epa.gov/iur/> [accessed 24 May 2012].
- Weschler CJ, Nazaroff WW. 2008. Semivolatile organic compounds in indoor environments. *Atmos Environ* 42:9018-9040.
- Wittassek M, Koch HM, Angerer J, Brüning T. 2011. Assessing exposure to phthalates – the human biomonitoring approach. *Mol Nutr Food Res* 55:7-31.
- Yoshida T, Andoh K, Fukuhara M. 2002. Urinary 2,5-dichlorophenol as biological index for *p*-dichlorobenzene exposure in the general population. *Arch Environ Contam Toxicol* 43:481-485.

**Table 1.** Computed arithmetic mean analyte levels in urine ( $\mu\text{g/g}$  creatinine) for US population (aged  $\geq 6$ ).<sup>a</sup>

Analyte	Year	$C_{50}$	$C_{75}$	$C_{90}$	$C_{95}$	GM	GSD	$r^2$	AM
BPA	03-04	2.50	4.29	7.67	11.2	2.43	2.5	1.00	3.7
DCP	05-06	7.32	20.4	89.3	292	6.0	9.2	0.98	71
MBzP	05-06	8.24	15.3	30.2	47.4	7.92	2.9	1.00	14
MEHP	05-06	2.61	5.69	13.7	30.1	2.38	4.3	0.98	6.9
MEHHP	05-06	21.4	46.1	117	235	19.6	4.2	0.99	56
MEOHP	05-06	13.5	28.9	77.7	144	12.4	4.2	0.99	35
MECPP	05-06	32.2	67.5	168	290	30.2	3.8	0.99	75
MEP	05-06	92.3	242	625	1140	90.0	4.6	1.00	288
MiBP	05-06	5.07	8.81	15.2	21.3	5.00	2.4	1.00	7.3
MnBP	05-06	18.3	30.8	50.8	77.8	17.7	2.4	0.99	26
MP	05-06	58.8	221	527	902	63.7	5.2	0.99	247
triclosan	03-04	9.48	43.9	212	368	9.67	9.8	1.00	131

<sup>a</sup> The  $z$ -scores (standard scores) used in the regression were 0.00 for  $C_{50}$ , 0.67 for  $C_{75}$ , 1.28 for  $C_{90}$ , and 1.64 for  $C_{95}$ . The analyte abbreviations refer to the following chemicals —bisphenol A (BPA), 2,5-dichlorophenol (DCP), mono-benzyl phthalate (MBzP), mono-2-ethylhexyl phthalate (MEHP), mono-(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP), mono-(2-ethyl-5-oxohexyl) phthalate (MEOHP), mono-(2-ethyl-5-carboxypentyl) phthalate (MECPP), mono-ethyl phthalate (MEP), mono-isobutyl phthalate (MiBP), mono-*n*-butyl phthalate (MnBP), and methyl paraben (MP). Urinary levels of bisphenol A include both conjugated and unconjugated forms.

**Table 2.** Fractional molar urinary excretion factors ( $f_{ue}$ ) and relevant molecular weights (MW) for converting excretion rates to intake rates.

Parent compound	MW (g/mol)	Analyte	MW (g/mol)	Urinary excretion, $f_{ue}$	Reference for $f_{ue}$
BPA	228	BPA	228	1.0	Dekant and Völkel 2008
BBzP	312	MBzP	256	0.73	Anderson et al. 2001
DEHP	391	MEHP	294	0.059	Koch et al. 2005
		MEHHP	294	0.233	Koch et al. 2005
		MEOHP	292	0.15	Koch et al. 2005
		MECPP	308	0.185	Koch et al. 2005
DEP	222	MEP	194	0.69	Assumed same as DnBP
DiBP	278	MiBP	222	0.69	Assumed same as DnBP
DnBP	278	MnBP	222	0.69	Anderson et al. 2001
<i>p</i> -dichlorobenzene	147	DCP	163	1.0	Yoshida et al. 2002 <sup>a</sup>
methyl paraben	152	MP	152	< 1.0	Upper bound
triclosan	290	triclosan	290	0.57	Sandborgh-Englund et al. 2006

<sup>a</sup> Personal monitoring and urinary excretion data, for ordinary environmental inhalation exposures among 119 adults in Osaka, are consistent with  $f_{ue} = 1.0$ .

**Table 3.** Intake-to-production ratio (IPR) for selected chemicals in the United States.

<b>Chemical</b>	<b>CAS No.</b>	<b><i>I</i> (µg/d per person)<sup>a</sup></b>	<b><i>P</i> (g/d per person)<sup>b</sup></b>	<b>IPR (ppm)</b>
bisphenol A (BPA)	80-05-7	5.6	9.8	0.6
butyl benzyl phthalate (BBzP)	85-68-7	35	0.30	120
di(2-ethylhexyl) phthalate (DEHP)	117-81-7	550	0.82	670
di( <i>n</i> -butyl) phthalate (DnBP)	84-74-2	71	0.094	760
para-dichlorobenzene	106-46-7	97	0.094	1040
di(isobutyl) phthalate (DiBP)	84-69-5	20	0.0030	6800
diethyl phthalate (DEP)	84-66-2	730	0.094	7700
triclosan	3380-34-5	350	0.014-0.044	8,000-24,000
methyl paraben	99-76-3	> 380	< 0.0021	> 180,000

<sup>a</sup> *I* represents the per capita average daily intake of the chemical across the US population.

<sup>b</sup> *P* represents the per capita average daily production plus importation rate of chemical for the US economy.

**Figure Caption**

Fig. 1. Intake-to-production ratio (IPR) estimates for the US population for nine manufactured chemicals. A value of 1 part per million (ppm) indicates that the aggregate intake, summed across the population, is one gram for every million grams manufactured in or imported into the population's economy.




Figure 1