



ENVIRONMENTAL  
HEALTH  
PERSPECTIVES

ehponline.org

Exposure to Tobacco Smoke *in Utero* and  
Subsequent Plasma Lipids, ApoB and CRP Among  
Adult Women in the MoBa Cohort

---

Lea A. Cupul-Uicab, Rolv Skjaerven, Kjell Haug,  
Gregory S. Travlos, Ralph E. Wilson, Merete Eggesbø,  
Jane A. Hoppin, Kristina W. Whitworth, Matthew P. Longnecker

<http://dx.doi.org/10.1289/ehp.1104563>

Online 19 July 2012



**NIEHS**  
National Institute of  
Environmental Health Sciences

National Institutes of Health  
U.S. Department of Health and Human Services

Exposure to Tobacco Smoke *in Utero* and Subsequent Plasma Lipids, ApoB and CRP Among  
Adult Women in the MoBa Cohort

Lea A. Cupul-Uicab<sup>1\*</sup>, Rolv Skjaerven<sup>2,3</sup>, Kjell Haug<sup>2</sup>, Gregory S. Travlos<sup>4</sup>, Ralph E. Wilson<sup>4</sup>,  
Merete Eggesbø<sup>5</sup>, Jane A. Hoppin<sup>1</sup>, Kristina W. Whitworth<sup>1</sup>, Matthew P. Longnecker<sup>1</sup>

<sup>1</sup>Epidemiology Branch, National Institute of Environmental Health Sciences, NIH/DHHS/USA,  
Research Triangle Park, NC, USA

<sup>2</sup>Department of Public Health and Primary Health Care, University of Bergen, Bergen, Norway

<sup>3</sup>Medical Birth Registry of Norway, the Norwegian Institute of Public Health, Bergen, Norway

<sup>4</sup>Cellular and Molecular Pathology Branch, National Institute of Environmental Health Sciences,  
NIH/DHHS/USA, Research Triangle Park, NC, USA

<sup>5</sup>Department of Genes and Environment, Division of Epidemiology, Norwegian Institute of  
Public Health, Oslo, Norway

\*Corresponding author:

Lea A. Cupul-Uicab, ScD.

National Institute of Environmental Health Sciences, MD A3-05, 111 TW Alexander Dr,  
Research Triangle Park, North Carolina 27709, USA; e-mail: [cupuluicabl@niehs.nih.gov](mailto:cupuluicabl@niehs.nih.gov),  
[acuicab@hotmail.com](mailto:acuicab@hotmail.com); tel. 919 541-3630; fax 919 541-2511

Running title: *In Utero* Tobacco Smoke and Dyslipidemia

Key words: clinical chemistry, C-reactive protein, metabolic syndrome, plasma lipids, prenatal exposure delayed effects, smoking, women

Acknowledgments:

We are grateful to all the participating families in Norway who take part in this ongoing cohort study.

Sources of Funding:

This research was supported in part by the Intramural Research Program of the National Institutes of Health (NIH), National Institute of Environmental Health Sciences (NIEHS). The Norwegian Mother and Child Cohort Study is supported by the Norwegian Ministry of Health, contract N01-ES-75558 with the National Institute of Environmental Health Sciences (NIH/NIEHS), NIH/NINDS (grant no.1 U01 NS 047537-01), and the Norwegian Research Council/FUGE (grant no. 151918/S10).

Conflict of interest: None declared

Abbreviations: apoB, apolipoprotein B; BMI, body mass index; CRP, C-reactive protein; CV, coefficient of variation; HDL, high-density lipoprotein cholesterol; LDL, low-density lipoprotein cholesterol; MoBa, The Norwegian Mother and Child Cohort Study; OR, odds ratio

## Abstract

**Background:** Recent findings suggest that maternal smoking during pregnancy may play a role in the development of metabolic alterations in offspring during childhood. However, whether such exposure increases the risk of developing similar metabolic alterations during adulthood is uncertain.

**Objective:** To evaluate the association of *in utero* exposure to maternal tobacco smoke with plasma lipids, apolipoprotein B (apoB), and C-reactive protein (CRP) in adulthood.

**Methods:** The study was based on a subsample of the Norwegian Mother and Child Cohort Study (MoBa), and included 479 pregnant women with a measure of plasma lipids, apoB, and CRP. *In utero* exposure to tobacco smoke (yes vs. no), personal smoking, and other factors were obtained by a self-completed questionnaire at enrollment (~17 weeks of gestation).

**Results:** Women exposed to tobacco smoke *in utero* had higher triglycerides (10.7% higher; 95% confidence interval [CI]: 3.9, 17.9) and lower high-density lipoprotein (HDL) cholesterol (-1.9 mg/dL; 95% CI: -4.3, 0.5) compared with unexposed women, after adjusting for age, physical activity, education, personal smoking, and current body mass index (BMI). Exposed women were also more likely to have triglycerides  $\geq 200$  mg/dL [adjusted odds ratio (aOR)=2.5; 95% CI: 1.3, 5.1] and HDL  $< 50$  mg/dL (aOR=2.3; 95% CI: 1.1, 5.0). Low-density lipoprotein cholesterol, total cholesterol, and apoB were not associated with the exposure. CRP was increased among exposed women, but after adjustment for BMI the association was completely attenuated.

**Conclusions:** In this population, *in utero* exposure to tobacco smoke was associated with high triglycerides and low HDL cholesterol in adulthood, 18 to 44 years after exposure.

## INTRODUCTION

Alterations in fetal programming in response to an adverse fetal environment may contribute to the development of metabolic syndrome (Gluckman and Hanson 2004). Components of the metabolic syndrome, such as elevated triglycerides and reduced high-density lipoprotein cholesterol (HDL) are known risk factors for cardiovascular disease and diabetes (Alberti et al. 2009). Recent findings from a longitudinal study suggest that maternal smoking during pregnancy may play a role in the development of metabolic alterations in offspring during childhood (Huang et al. 2007). However, whether *in utero* exposure to tobacco smoke is associated with metabolic alterations during adulthood is uncertain (Power et al. 2010) and the available epidemiologic data are insufficient. Previous studies, however, have reported an association of *in utero* exposure to tobacco smoke with cholesterol (Jaddoe et al. 2008; Wen et al. 2010) and carotid wall thickening during adulthood (Geerts et al. 2008). To our knowledge, other metabolic risk factors, such as elevated apolipoprotein B (apoB) and C-reactive protein (CRP), have not been studied in relation to *in utero* exposure to tobacco smoke. Elevated triglycerides and reduced HDL (i.e. atherogenic dyslipidemia) are usually accompanied by elevated apoB; CRP is a marker of inflammation that tends to be elevated in people with atherosclerotic conditions and metabolic syndrome (Grundy et al. 2004).

The purpose of the present study was to evaluate the association of *in utero* exposure to maternal tobacco smoke (i.e. 18-44 years earlier for women in the present analysis) with alterations in plasma lipids that are compatible with metabolic syndrome in adulthood. We also evaluated the association of *in utero* exposure to tobacco smoke with low-density lipoprotein cholesterol (LDL), total cholesterol, apoB, and CRP.

## METHODS

This study was based on the Norwegian Mother and Child Cohort Study (MoBa), conducted by the Norwegian Institute of Public Health (Magnus et al. 2006). MoBa is a cohort based on 108,000 pregnancies from 90,700 women enrolled from 1999 to 2008. The majority of all pregnant women in Norway were invited to participate, and 38.5% of invited women participated in the study. Participants were recruited with a mailed invitation before a routine ultrasound examination offered to all pregnant women in Norway at 17-18 weeks of gestation (NIPH (Norwegian Institute of Public Health) 2007)]. The study was approved by the Regional Committee for Medical Research Ethics and the Norwegian Data Inspectorate. Informed consent was provided by each participant. The present study is based on version 4.301 of the quality-assured data files released for research in December 2009. At enrollment, participants were asked to give a plasma specimen (median, 18 weeks of gestation; 25th and 75th percentiles, 17 and 19 weeks) and to complete a questionnaire about demographic characteristics, reproductive health, disease and medication history, lifestyle, and socioeconomic status. The present analysis was based on a sub-sample of 950 women whose plasma specimens were analyzed for lipids, apoB, and CRP for a case-base (Kupper et al. 1975) study on sub-fecundity (Whitworth et al. 2012). For the case-base study, eligibility was restricted to all participants enrolled in 2003 and 2004 who delivered a live born child, provided a plasma specimen, and reported their time-to-pregnancy (n=8,120). Cases (n=400) were selected at random among eligible women who were subfecund (i.e. with a time-to-pregnancy >12 months, n=839); the base sample (n=550) was selected at random from all eligible women regardless of subfecundity status.

### Assessment of *in utero* tobacco smoke

Women's exposure to tobacco smoke *in utero* was ascertained on the baseline questionnaire. Women were asked: "Did your mother smoke when she was pregnant with you?" Those who answered "yes" were classified as having been exposed to tobacco smoke *in utero*; those who responded "no" were considered unexposed. For women who participated with more than one pregnancy in the MoBa cohort, the consistency of answers across pregnancies was verified. In general, if the woman gave two different answers in two consecutive pregnancies (e.g. yes/no; no/yes; no/don't know; yes/don't know), the response was considered inconsistent and the subject was excluded from the analysis. However, if the first answer was "Don't know" and later she gave a different answer (yes or no), we used the latter under the assumption that the woman had asked her mother about her exposure *in utero*. Analyses of data from the MoBa cohort support the validity and reproducibility of self-reported *in utero* exposure to tobacco smoke among MoBa participants (Cupul-Uicab et al. 2011a; Cupul-Uicab et al. 2011c). Women were not asked about cigarette smoke exposure during childhood.

### Plasma measurements

At enrollment, women provided a non-fasting blood specimen collected in ethylenediaminetetraacetic acid (EDTA) tubes; the samples were shipped at ambient temperature from the collection site to Oslo where plasma was extracted and stored at -80° C (the majority of the samples were received the day after collection) (Ronningen et al. 2006). For the present study we used the plasma specimens provided by the women at enrollment. Although some women participated with multiple pregnancies in MoBa, women could contribute with only one pregnancy in the present analysis. Triglycerides, total cholesterol, apoB, CRP, HDL, and LDL

were determined in plasma specimens with an Olympus AU400e Clinical Chemistry Analyzer (Olympus America, Inc., Irvin, TX) using reagents from Beckman Coulter (for triglycerides and cholesterol) and Genzyme Diagnostics (for the others). Quantitative measurement of LDL was done with a direct enzymatic method (N-geneous® LDL-ST cholesterol reagent). CRP was expressed in mg/L and the other clinical chemistries were expressed in mg/dL. The within and between batch coefficients of variation (CV) of the analytical method were < 5% for all clinical chemistries, except for the between batch CV for triglycerides (CV 6.3%).

### Statistical analysis

For the present analysis, women from the case-base study (n=950) were excluded if they had missing data on body mass index (BMI), education, or clinical chemistries (n=8). In addition, women were excluded if, for *in utero* tobacco smoke, there was unknown exposure (n=112) or inconsistent answers (n=7) (see above). After all the exclusions, a total of 823 women were included in the analysis (base sample, n=479).

The associations of *in utero* exposure to tobacco smoke with levels of triglycerides, HDL, LDL, total cholesterol, apoB, and CRP were assessed separately. The main analysis was restricted to women from the base sample (selected without regard to subfecundity status). We also conducted an analysis that included all women from the MoBa sub-sample using weighted linear regression to account for the case-base sampling (Richardson et al. 2007). The weights were the inverse of the sampling probabilities for the original case-base study. Triglycerides and CRP had a skewed distribution and therefore were natural logarithm transformed before modeling; their corresponding coefficients represent the percent difference in geometric mean values for the exposed compared with the unexposed group. For some of the outcomes, we also

assessed whether clinically relevant alterations in levels were associated with *in utero* exposure to tobacco smoke using logistic regression. For this purpose, we used previously recommended cut points to define high triglycerides ( $\geq 200$  mg/dL), low HDL ( $< 50$  mg/dL for females), high total cholesterol ( $\geq 240$  mg/dL), and high CRP ( $\geq 8$  mg/L) (NCEP 2001; Pitiphat et al. 2005).

All models included *in utero* tobacco smoke as the main exposure and were adjusted for woman's age (years), education (high school or less versus more than high school), and physical activity (times a week that the women was engaged in activities such as brisk walking, running/jogging/orienteering, bicycling, gymnastic, aerobics, dancing, skiing, and swimming; categorized as none,  $< 4$  and 4 or more times a week). We use directed acyclic graphs (DAGs) to select this set of *a priori* variables (Greenland et al. 1999). Although the minimal sufficient set of variables to adjust for confounding did not include education, we choose to adjust for participants' education as a proxy indicator of their mother's education and socioeconomic status (see Supplemental Material, Figure S1). Participant's individual annual income was not selected *a priori* because of its correlation with education (Spearman  $r=0.43$ ,  $p < 0.01$ ) and because the exposure and outcomes had stronger correlations with education than with income. Information on the education, socioeconomic status, and lifestyle habits of participants' mothers was not ascertained in MoBa. Additional variables that were associated with at least one of the outcomes in bivariate analyses ( $p \leq 0.20$ ) were assessed as potential confounders (i.e. income, parity, and alcohol consumption (no, yes)) using the change in estimate method, starting with all variables in the models with deletion of one by one in a stepwise manner (Greenland 1989). None of the tested variables caused a change  $\geq 10\%$  in the coefficient for *in utero* smoking, thus we did not adjust for them. Participants' personal smoking, BMI (weight kg/height  $m^2$ ), and birth weight (kg) were potential intermediate variables that were included in selected models. BMI was

entered as a continuous variable in the models because the results were comparable to those obtained using four categories. The effect of adjusting for birth weight was evaluated among the subset of women born in 1967 or later (when the Medical Birth Registry of Norway was established) with available birth weight [88.8% of 752 (89% of 446 for the base sample)]. Multiplicative interactions of *in utero* exposure to tobacco smoke (yes/no) with personal smoking (yes/no), BMI ( $\text{kg}/\text{m}^2$ ), and birth weight (kg) were tested in linear regression models but are not presented because the *p*-values for all interaction terms were  $>0.15$ . We also estimated associations of *in utero* exposure to tobacco smoke (yes/no) with the outcomes in the absence or presence of overweight and obesity ( $\text{BMI} < 25$  and  $\geq 25 \text{ kg}/\text{m}^2$ ) to assess departures from additive effects (Hosmer and Lemeshow 1992). However, numbers were too sparse to conduct similar analyses for smoking and birth weight.

We conducted a number of sensitivity analyses to assess the robustness of our results. We added participant's individual income to models in addition to education because adjusting for both variables might be a more effective means of reducing confounding due to socioeconomic disadvantages throughout the participant's life. We also estimated associations adjusted for age and physical activity only, the minimal sufficient set of covariates based on the DAG. Because lipid levels can vary throughout pregnancy (Vahratian et al. 2010) and plasma specimens were not provided at the same week of gestation for all women, the models were further adjusted for gestational week at blood draw.

In additional sensitivity analysis, multiple imputation by chained equations (MICE) (van Buuren et al. 1999) was performed in the complete sample (i.e. 950 subjects from the original case-base study) to impute values for any variable with missing data. For the imputation procedure we included all clinical chemistries (triglycerides and CRP entered as natural

logarithm), *in utero* exposure to tobacco smoke, all characteristics listed in Table 1, and BMI before pregnancy, weight and gestational age at birth, year of birth, year of enrollment, and age of the participant's mother at delivery. A total of 10 imputed datasets were generated using 20 cycles per imputation (van Buuren et al. 1999), and analyses were repeated using the imputed data. All analyses were done using Stata (Stata Statistical Software, release 10.1; StataCorp, College Station, TX, USA).

## RESULTS

The prevalence of *in utero* exposure to tobacco smoke among women in the base sample was higher (> 35%) among those with less than a college education, lower income, BMI  $\geq 30$ , who smoked more as adults, and women who reported no physical activity (Table 1). Exposure prevalences according to participant characteristics were similar among all 823 women (data not shown). Among women from the base sample, 7.9% had high triglycerides, 6.9% had low HDL, 20.3% had high cholesterol, and 22.1% had high CRP, with similar prevalences among subfecund MoBa participants (see Supplemental Material, Table S1). Compared to subfecund women, those from the base sample tended to be younger, and a higher proportion of them were college educated, had a previous pregnancy, had a BMI <25, and did not smoke (Supplemental Material, Table S1). The range of lipids, apoB, and CRP measured in our sample (Supplemental Material, Table S2) were comparable to reference levels for pregnant women in other populations (Klajnbard et al. 2010; Larsson et al. 2008).

Compared with unexposed women, those exposed to tobacco smoke *in utero* had higher median levels of triglycerides and CRP, and lower mean HDL (Table 2). After adjustment (for

age, physical activity and education) significant associations with triglycerides (13.4% higher; 95% CI: 6.4, 20.9) and HDL (2.6 mg/dL lower; 95% CI: -5.0, -0.2) remained. Associations with all outcomes were similar after additional adjustment for personal smoking, but were attenuated after adjustment for BMI (Table 3). Average values of LDL, total cholesterol, and apoB tended to be higher among women exposed to tobacco smoke *in utero*, but the estimates were imprecise. The positive association of CRP with exposure to tobacco smoke *in utero* was still evident (though non-significant) after adjustment for age, physical activity, and education (16.3% higher in the exposed versus unexposed; 95% CI: -2.6, 39.0), but was close to the null after adjustment for BMI (2.7% higher; 95%CI: -12.7, 20.9) (Table 3). When we analyzed the data from all women using weighted linear regression, *in utero* tobacco smoke remained associated with triglycerides (12.1% higher in the exposed; 95% CI: 5.5, 19.2), HDL cholesterol (2.3 mg/dL lower; 95% CI: -4.5, -0.1), and CRP (14.7% higher; 95% CI: -1.4, 33.5); estimates for LDL cholesterol (3.3 mg/dL higher; 95% CI: -2.5, 9.1), total cholesterol (2.3 mg/dL higher; 95% CI: -4.5, 9.1), and apoB (2.9 mg/dL higher; 95% CI: -1.5, 7.2) were similar to those from the base sample. After adjusting for BMI, the estimates were again attenuated, but were consistent with those from the base sample; *in utero* exposure to tobacco smoke remained associated with triglycerides (9.4% higher in the exposed versus unexposed; 95% CI: 3.1, 16.1) and HDL cholesterol (1.7 mg/dL lower; 95% CI: -3.9, 0.6), but not with CRP (2.1% higher; 95% CI: -10.9, 17.1).

Among 397 women in the base sample with known birth weight (Table 4), triglycerides remained higher and HDL lower among exposed compared with unexposed before and after adjusting for birth weight (in addition to age, education, physical activity, personal smoking, and BMI), and when the analysis was restricted to 381 women born at term (i.e. gestational age  $\geq 37$

weeks). Other lipids, apoB, and CRP were again unrelated to *in utero* exposure to tobacco smoke (Table 4). Associations were similar after adjustment for birth weight among all women (data not shown).

Compared with unexposed women, women exposed to tobacco smoke *in utero* were more likely to have triglycerides  $\geq 200$  mg/dL (adjusted OR [aOR]=2.5; 95% CI: 1.3, 5.1), HDL  $< 50$  mg/dL (aOR=2.3; 95% CI: 1.1, 5.0), and CRP  $\geq 8$  mg/L (aOR=1.6; 95% CI: 1.0, 2.6); the odds of having cholesterol  $\geq 240$  mg/dL was slightly higher (aOR=1.4, 95% CI: 0.9, 2.3) among exposed women (Table 5). After additional adjustment for BMI, the association of *in utero* exposure to tobacco with high triglycerides and low HDL cholesterol remained statistically significant, but that was not the case for CRP. In the analysis that included all women, the associations were consistent with those from the base sample (Table 5).

Compared with women who were unexposed and had a BMI  $< 25$  (kg/m<sup>2</sup>), women exposed to tobacco smoke *in utero* tended to have higher odds of high triglycerides, low HDL, and high total cholesterol regardless of their BMI ( $< 25$  or  $\geq 25$ ), whereas women with BMI  $\geq 25$  tended to have higher odds of high CRP regardless of their exposure; the confidence intervals, however, were wide (Table 6). Among unexposed women, higher BMI ( $\geq 25$  kg/m<sup>2</sup>) was also associated with higher odds of having high triglycerides, low HDL cholesterol, and high total cholesterol.

Estimates from models adjusted for participant's age and physical activity only (the minimum sufficient adjustment set based on a DAG), fell between unadjusted estimates and estimates adjusted for age, education, and physical activity [triglycerides: 15.3% higher in the exposed versus unexposed (95% CI: 8.1, 22.9); HDL cholesterol: 2.9 mg/dL lower (95% CI: -5.2, -0.5); and CRP: 20.8% higher (95% CI: 1.2, 44.1)]. The results from Table 3 and 5 remained

essentially the same after additional adjustment for gestational week at blood draw and income (data not shown). The results obtained from the multiple imputation analyses were comparable to those observed based on complete data (data not shown). Among women selected as cases (i.e. subfecund), *in utero* exposure to tobacco smoke was not associated with any outcome before or after adjusting for BMI; after BMI was added to the models the estimates for triglycerides, HDL cholesterol, and CRP were not in the expected direction (Supplemental Material, Table S3).

## DISCUSSION

In the present study, exposure to tobacco smoke *in utero* was associated with higher triglycerides and lower HDL cholesterol during adulthood in a population of pregnant women. These adverse alterations in plasma lipids are compatible with the metabolic syndrome, and therefore of clinical relevance. The odds of having elevated triglycerides or low HDL cholesterol levels were doubled among women exposed to tobacco smoke *in utero* as compared with unexposed. Plasma LDL, total cholesterol, and apoB were not associated with the exposure in the adjusted analyses. Women exposed to tobacco smoke *in utero* also tended to have higher CRP, though this association was explained by the association of the exposure with BMI.

An adverse lipid profile early in life has been reported among children and adolescents exposed to tobacco smoke *in utero* as well as those exposed to parental smoking during childhood (Huang et al. 2007; Metsios et al. 2011). Among 8,815 men and women (~45 years of age) from the 1958 British birth cohort, *in utero* exposure to tobacco smoke was associated with high triglycerides and low HDL cholesterol, though the latter was limited to women (Power et al. 2010). However, after accounting for life-time covariates and potential mediators (e.g. birth

weight, gestational age, breastfeeding, education, physical activity, personal smoking) simultaneously in the analysis these associations were weaker and no longer statistically significant (Power et al. 2010). Similarly, among 3,824 men and women (23 years of age) from a Brazilian birth cohort, *in utero* exposure to tobacco smoke was associated with low HDL cholesterol in women but not in men; this association was also weakened and no longer statistically significant after accounting for confounders and potential mediators (e.g birth weight, physical activity, personal smoking, BMI, waist circumference) (Horta et al. 2011). In the present study, the positive association between triglycerides and *in utero* exposure to tobacco smoke was consistent and remained after all adjustments; it was also supported by our analysis using the recommended clinical cut-point used to define high triglycerides ( $\geq 200$  mg/dL) (NCEP 2001). As reported in previous studies (Horta et al. 2011; Power et al. 2010), the association between reduced HDL and exposure to tobacco smoke *in utero* may not be independent of adult BMI; however, when using the recommended cut-point for defining low HDL (for women,  $< 50$  mg/dL) (NCEP 2001) the odds were doubled among exposed women compared with unexposed.

Our results differ from two recent studies where *in utero* exposure to tobacco smoke was associated with total cholesterol in adulthood (Jaddoe et al. 2008; Wen et al. 2010). We observed a positive association between *in utero* exposure to tobacco smoke and total cholesterol, but associations were weak and estimates were imprecise. Jaddoe et al. reported that the association of *in utero* smoking with total cholesterol was stronger among participants with moderate overweight compared to those with normal BMI (Jaddoe et al. 2008). In our data there was no significant departure from additive or multiplicative joint effects of BMI and *in utero* exposure to tobacco smoke. In addition, the odds of having high cholesterol among women exposed to tobacco smoke *in utero* were similar when the BMI was  $< 25$  and  $\geq 25$  kg/m<sup>2</sup>.

Whether *in utero* exposure to tobacco smoke (or parental smoking during childhood as a surrogate of *in utero* exposure) is a risk factor for metabolic syndrome during adulthood remains unclear (Hunt et al. 2006; Power et al. 2010). We did not have data on blood pressure, fasting glucose, or central obesity, which are also components of the metabolic syndrome in addition to plasma triglycerides and HDL cholesterol (Alberti et al. 2009). However, a higher prevalence of hypertension and obesity in relation to tobacco smoke *in utero* has been previously reported among 74,023 women enrolled in MoBa, which included some of the participants in the present analysis (Cupul-Uicab et al. 2011b).

In animals, fetal exposure to nicotine at doses relevant to humans (i.e. serum cotinine levels of 136-300 ng/ml as found among moderate to heavy smokers) results in long-term metabolic alterations in adulthood that are consistent with components of the metabolic syndrome in humans (Bruin et al. 2010). In humans however, other less studied constituents of tobacco smoke, such as carbon monoxide, might also contribute to adverse lipid profiles (i.e. carbon monoxide is associated with fetal hypoxia, which can increase oxidative stress and potentially alter lipid metabolism in the fetus) (Chelchowska et al. 2011; Wen et al. 2010).

As in other observational studies, causal associations cannot be established with these data. It is possible that *in utero* exposure to tobacco smoke is acting as a marker of socio-economic disadvantage during childhood, and the latter may be related to a higher risk for adult disease (Donovan and Susser 2011). Lifetime socioeconomic status of the participants was not available in MoBa. As shown in previous studies and in the present study, adjustment for factors such as BMI usually leads to weaker associations; however, the appropriateness of adjustment for risk factors that are affected by the exposure is debatable (Cole and Hernan 2002).

Estimated effects of exposure in conjunction with BMI showed strong associations of *in utero* exposure to tobacco smoke with high triglycerides (aOR 4.7; 95% CI: 1.4, 15.7) and reduced HDL (aOR 4.4; 95% CI: 1.4, 14.4) among women with normal BMI (Table 6). The combined estimate was consistent with additive effects of BMI and exposure on triglycerides and HDL cholesterol, but the statistical power of the study for evaluation of interaction was limited. BMI was associated with high CRP regardless of exposure, and the combined estimate suggests a weak association between *in utero* exposure to tobacco smoke and high CRP among overweight or obese women.

Although exposure data were collected retrospectively, reported exposure to maternal tobacco smoke *in utero* by the adult offspring has shown to be valid (Simard et al. 2008); and previous analysis from MoBa participants also supports the validity and reproducibility of self-reported exposure to tobacco smoke *in utero* (Cupul-Uicab et al. 2011a; Cupul-Uicab et al. 2011c). An analysis of data from multiple populations indicated that birth weight is 149 g lower, on average, in children whose mother's smoked during pregnancy compared with children whose mother's did not smoke (Kramer 1987). The estimated average 181 g reduction in birth weight associated with *in utero* exposure to tobacco smoke in a subset of 11,082 MoBa participants indirectly supports the validity of self-reported *in utero* exposure (Cupul-Uicab et al. 2011c). The reproducibility of self-reported *in utero* exposure to tobacco smoke among MoBa participants who completed questionnaires for multiple pregnancies was high (weighted  $\kappa=0.80$ ) (Cupul-Uicab et al. 2011c). The intensity of *in utero* exposure to tobacco smoke was not ascertained in MoBa; therefore we were unable to assess a dose-response relationship.

The prevalence of *in utero* exposure to tobacco smoke in the present study (28% in the base sample) was similar to that for the MoBa cohort as a whole (~27.8%), but the association

between *in utero* exposure to tobacco smoke and obesity was slightly stronger in the base sample (aOR 2.0; 95% CI: 1.10, 3.77) than in the full MoBa cohort (aOR = 1.53; 95% CI: 1.45, 1.61) (Cupul-Uicab et al. 2011b). As expected, physical activity was associated with higher HDL cholesterol and lower triglycerides, cholesterol, and LDL cholesterol in the present study (data not shown).

Among subfecund women (i.e. with a time to pregnancy > 12months), *in utero* exposure to tobacco smoke was not associated with lipids, apoB, or CRP. Subfecund women are more likely to have conditions associated with lipid alterations (e.g. polycystic ovary syndrome) (Norman et al. 2007) that are probably stronger predictors of the outcomes than *in utero* exposure to tobacco smoke, which may explain the lack of association among this subset of women.

In the present study lipids were measured in non-fasting plasma specimens. Although triglycerides are known to be higher in the postprandial period, a recent prospective study among 26,509 women suggests that non-fasting triglycerides may be a better predictor of cardiovascular events (Bansal et al. 2007). The impact of adjusting for recent food consumption on associations between *in utero* exposure to tobacco smoke and components of the metabolic syndrome was negligible in the 1958 British cohort, a previous study that also measured lipids in non-fasting samples (Power et al. 2010). Because the cutoff points to define high triglycerides are based on fasting levels of triglycerides, potential misclassification of this outcome may be a concern; however, this might not be a concern for the analysis based on the continuous outcome. Furthermore, our results may not be directly comparable to those from fasting subjects or from non-pregnant women, and may not be generalizable to non-pregnant women. Whether or not

metabolic syndrome is a real entity (Kahn et al. 2005), clearly hypertriglyceridemia is an independent risk factor for coronary heart disease (NCEP 2001).

## **CONCLUSION**

Women exposed to tobacco smoke *in utero* were more likely to have high triglycerides and low HDL cholesterol as adults. These adverse alterations in plasma lipids are compatible with metabolic syndrome and may have implications for future cardiovascular disease among the exposed.

## References

- Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. 2009. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 120(16):1640-1645.
- Bansal S, Buring JE, Rifai N, Mora S, Sacks FM, Ridker PM. 2007. Fasting compared with nonfasting triglycerides and risk of cardiovascular events in women. *Jama* 298(3):309-316.
- Bruin JE, Gerstein HC, Holloway AC. 2010. Long-term consequences of fetal and neonatal nicotine exposure: a critical review. *Toxicol Sci* 116(2):364-374.
- Chelchowska M, Ambroszkiewicz J, Gajewska J, Laskowska-Klita T, Leibschang J. 2011. The effect of tobacco smoking during pregnancy on plasma oxidant and antioxidant status in mother and newborn. *Eur J Obstet Gynecol Reprod Biol* 155(2):132-136.
- Cole SR, Hernan MA. 2002. Fallibility in estimating direct effects. *Int J Epidemiol* 31(1):163-165.
- Cupul-Uicab LA, Baird DD, Skjaerven R, Saha-Chaudhuri P, Haug K, Longnecker MP. 2011a. In utero exposure to maternal smoking and women's risk of fetal loss in the Norwegian Mother and Child Cohort (MoBa). *Human reproduction (Oxford, England)* 26(2):458-465.
- Cupul-Uicab LA, Skjaerven R, Haug K, Melve KK, Engel SM, Longnecker MP. 2011b. In Utero Exposure to Maternal Tobacco Smoke and Subsequent Obesity, Hypertension, and Gestational Diabetes Among Women in The MoBa Cohort. *Environ Health Perspect*; <http://dx.doi.org/10.1289/ehp.1103789> [Online 29 Nov 2011].
- Cupul-Uicab LA, Ye X, Skjaerven R, Haug K, Longnecker MP. 2011c. Reproducibility of reported in utero exposure to tobacco smoke. *Annals of epidemiology* 21(1):48-52.
- Donovan SJ, Susser E. 2011. Commentary: Advent of sibling designs. *Int J Epidemiol* 40(2):345-349.

Geerts CC, Bots ML, Grobbee DE, Uiterwaal CS. 2008. Parental smoking and vascular damage in young adult offspring: is early life exposure critical? The atherosclerosis risk in young adults study. *Arterioscler Thromb Vasc Biol* 28(12):2296-2302.

Gluckman PD, Hanson MA. 2004. The developmental origins of the metabolic syndrome. *Trends in endocrinology and metabolism: TEM* 15(4):183-187.

Greenland S. 1989. Modeling and variable selection in epidemiologic analysis. *Am J Public Health* 79(3):340-349.

Greenland S, Pearl J, Robins JM. 1999. Causal diagrams for epidemiologic research. *Epidemiology* 10(1):37-48.

Grundey SM, Brewer HB, Jr., Cleeman JI, Smith SC, Jr., Lenfant C. 2004. Definition of metabolic syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Circulation* 109(3):433-438.

Horta BL, Gigante DP, Nazmi A, Silveira VM, Oliveira I, Victora CG. 2011. Maternal smoking during pregnancy and risk factors for cardiovascular disease in adulthood. *Atherosclerosis* 219(2):815-820.

Hosmer DW, Lemeshow S. 1992. Confidence interval estimation of interaction. *Epidemiology* 3(5):452-456.

Huang RC, Burke V, Newnham JP, Stanley FJ, Kendall GE, Landau LI, et al. 2007. Perinatal and childhood origins of cardiovascular disease. *Int J Obes* 31(2):236-244.

Hunt KJ, Hansis-Diarte A, Shipman K, Korte JE, Fowler SP, Stern MP. 2006. Impact of parental smoking on diabetes, hypertension and the metabolic syndrome in adult men and women in the San Antonio Heart Study. *Diabetologia* 49(10):2291-2298.

Jaddoe VW, de Ridder MA, van den Elzen AP, Hofman A, Uiterwaal CS, Witteman JC. 2008. Maternal smoking in pregnancy is associated with cholesterol development in the offspring: A 27-years follow-up study. *Atherosclerosis* 196(1):42-48.

Kahn R, Buse J, Ferrannini E, Stern M. 2005. The metabolic syndrome: time for a critical appraisal: joint statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care* 28(9):2289-2304.

- Klajnbard A, Szecsi PB, Colov NP, Andersen MR, Jorgensen M, Bjorngaard B, et al. 2010. Laboratory reference intervals during pregnancy, delivery and the early postpartum period. *Clin Chem Lab Med* 48(2):237-248.
- Kramer MS. 1987. Determinants of low birth weight: methodological assessment and meta-analysis. *Bull World Health Organ* 65(5):663-737.
- Kupper LL, McMichael AJ, Spirtas R. 1975. A hybrid epidemiological study design useful in estimating relative risk. *J Am Stat Assoc* 70(351):524-528.
- Larsson A, Palm M, Hansson LO, Axelsson O. 2008. Reference values for clinical chemistry tests during normal pregnancy. *Bjog* 115(7):874-881.
- Magnus P, Irgens LM, Haug K, Nystad W, Skjaerven R, Stoltenberg C. 2006. Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa). *Int J Epidemiol* 35(5):1146-1150.
- Metsios GS, Flouris AD, Angioi M, Koutedakis Y. 2011. Passive smoking and the development of cardiovascular disease in children: a systematic review. *Cardiology research and practice* 2011 (doi:10.4061/2011/587650):6.
- NCEP. 2001. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *Jama* 285(19):2486-2497.
- NIPH (Norwegian Institute of Public Health). 2007. Norwegian Mother and Child Cohort Study (MoBa). Available: [http://www.fhi.no/eway/default.aspx?pid=238&trg=MainArea\\_5811&MainArea\\_5811=5895:0:15,3046:1:0:0:::0:0](http://www.fhi.no/eway/default.aspx?pid=238&trg=MainArea_5811&MainArea_5811=5895:0:15,3046:1:0:0:::0:0) [accessed 6 June 2011].
- Norman RJ, Dewailly D, Legro RS, Hickey TE. 2007. Polycystic ovary syndrome. *Lancet* 370(9588):685-697.
- Pitiphat W, Gillman MW, Joshipura KJ, Williams PL, Douglass CW, Rich-Edwards JW. 2005. Plasma C-reactive protein in early pregnancy and preterm delivery. *Am J Epidemiol* 162(11):1108-1113.
- Power C, Atherton K, Thomas C. 2010. Maternal smoking in pregnancy, adult adiposity and other risk factors for cardiovascular disease. *Atherosclerosis* 211(2):643-648.

Richardson DB, Rzehak P, Klenk J, Weiland SK. 2007. Analyses of case-control data for additional outcomes. *Epidemiology* 18(4):441-445.

Ronningen KS, Paltiel L, Meltzer HM, Nordhagen R, Lie KK, Hovengen R, et al. 2006. The biobank of the Norwegian Mother and Child Cohort Study: a resource for the next 100 years. *Eur J Epidemiol* 21(8):619-625.

Simard JF, Rosner BA, Michels KB. 2008. Exposure to cigarette smoke in utero: comparison of reports from mother and daughter. *Epidemiology* 19(4):628-633.

Vahratian A, Misra VK, Trudeau S, Misra DP. 2010. Prepregnancy body mass index and gestational age-dependent changes in lipid levels during pregnancy. *Obstet Gynecol* 116(1):107-113.

van Buuren S, Boshuizen HC, Knook DL. 1999. Multiple imputation of missing blood pressure covariates in survival analysis. *Statistics in medicine* 18(6):681-694.

Wen X, Triche EW, Hogan JW, Shenassa ED, Buka SL. 2010. Birth weight and adult hypercholesterolemia: subgroups of small-for-gestational-age based on maternal smoking status during pregnancy. *Epidemiology* 21(6):786-790.

Whitworth KW, Haug LS, Baird DD, Becher G, Hoppin JA, Skjaerven R, et al. 2012. Perfluorinated compounds and subfecundity in pregnant women. *Epidemiology* 23(2):257-263.

Table 1. Characteristics of women from the MoBa base sample at enrollment according to self-reported *in utero* exposure to tobacco smoke

Characteristics	Exposed to tobacco smoke <i>in utero</i>		<i>p</i> <sup>a</sup>
	N	%	
All women	479	28.0	
Age (years)			0.07
< 25	43	27.9	
25 - 29	183	34.4	
30 - 34	179	21.8	
≥ 35	74	27.0	
Education			<0.01
< High School	32	37.5	
High School	141	39.0	
College	214	22.9	
> College	92	19.6	
Individual annual income (USD)			0.07
< 30,847	119	36.1	
30,847 - 46,269	200	27.5	
46,270 - 61,693	115	24.3	
> 61,693	45	17.8	
Parity			0.71
0	207	26.1	
1	186	29.0	
≥ 2	86	30.2	
BMI (kg/m <sup>2</sup> )			<0.01
< 25.0	253	22.9	
25.0 - < 30.0	161	28.6	
30.0 +	65	46.2	

Table 1 (cont.)

Characteristics	Exposed to tobacco smoke in utero		<i>p</i> <sup>a</sup>
	N	%	
Smoking (cigarettes/day)			0.02
Non smoker	378	25.7	
< 10	87	33.3	
10 +	14	57.1	
Alcohol drinking			0.77
Non drinker	362	27.3	
Yes	73	31.5	
Missing	44	27.3	
Physical activity (times/week)			0.07
None	99	38.4	
< 4.0	261	26.1	
4.0 - < 7.5	89	24.7	
≥ 7.5	30	20.0	

<sup>a</sup>*p*-values are from Pearson's chi-squared test comparing exposed and unexposed across categories of each variable

Table 2. Plasma lipids, apolipoprotein B, and CRP by *in utero* exposure to tobacco smoke among women from the base sample at enrollment (n = 479)

Plasma levels (mg/dL) <sup>a</sup>	Exposed to tobacco smoke <i>in utero</i>		<i>p</i> <sup>b</sup>
	No (n = 345)	Yes (n = 134)	
Triglycerides	117.2 (47.0)	135.7 (71.0)	<0.01
HDL-C	68.4 ± 11.6	65.3 ± 12.7	0.01
LDL-C	126.5 ± 29.4	130.4 ± 32.4	0.20
Total cholesterol	212.8 ± 33.7	215.2 ± 38.5	0.50
ApoB	99.7 ± 21.7	103.4 ± 24.5	0.11
CRP (mg/L)	4.0 (4.5)	5.0 (5.4)	0.01

Abbreviations: apoB, Apolipoprotein B; CRP, C-reactive protein; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; SD, standard deviation

<sup>a</sup> Plasma levels are presented as means ± standard deviation or geometric means (interquartile range)

<sup>b</sup>*p*-values are from unadjusted linear regressions, except for triglycerides and CRP. Two-sample Wilcoxon rank-sum (Mann-Whitney) test for equality-of-medians was used for triglycerides and CRP

Table 3. Estimated associations of *in utero* exposure to tobacco smoke with lipids, apolipoprotein B, and CRP among adult women from the base sample (n = 479)

Outcomes <sup>a</sup>	Unadjusted		Adjusted							
	$\beta$	(95% CI)	Age, physical activity and education only		Plus personal smoking		Plus BMI		Plus personal smoking and BMI	
	$\beta$	(95% CI)	$\beta$	(95% CI)	$\beta$	(95% CI)	$\beta$	(95% CI)	$\beta$	(95% CI)
Triglycerides	15.8	(8.7, 23.4)	13.4	(6.4, 20.9)	13.4	(6.4, 21.0)	10.6	(3.8, 17.8)	10.7	(3.9, 17.9)
HDL-C	-3.2	(-5.5, -0.8)	-2.6	(-5.0, -0.2)	-2.5	(-4.9, -0.1)	-2.0	(-4.4, 0.4)	-1.9	(-4.3, 0.5)
LDL-C	3.9	(-2.1, 10.0)	3.3	(-2.9, 9.4)	3.2	(-3.0, 9.4)	2.8	(-3.4, 9.1)	2.8	(-3.5, 9.1)
Total cholesterol	2.4	(-4.6, 9.4)	2.1	(-5.1, 9.2)	2.3	(-4.9, 9.4)	1.9	(-5.3, 9.2)	2.1	(-5.2, 9.3)
ApoB	3.7	(-0.8, 8.2)	2.8	(-1.7, 7.4)	2.9	(-1.7, 7.5)	2.2	(-2.4, 6.8)	2.3	(-2.4, 6.9)
CRP	24.6	(4.5, 48.5)	16.3	(-2.6, 39.0)	15.9	(-3.0, 38.5)	2.7	(-12.7, 20.9)	2.9	(-12.6, 21.1)

Abbreviations: ApoB, apolipoprotein B; BMI, body mass index; CRP, C-reactive protein; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol

<sup>a</sup>Associations are expressed as the % difference in the geometric mean for log<sub>n</sub>-transformed outcomes (mg/dL triglycerides and mg/L CRP) or the difference in mean values (mg/dL of HDL-C, LDL-C, total cholesterol, and ApoB) in the exposed compared with the unexposed group.

Table 4. Adjusted<sup>a</sup> coefficients for lipids, apolipoprotein B, and CRP by *in utero* exposure to tobacco smoke among women from the MoBa base sample with available birth weight

Outcomes <sup>c</sup>	All women with available birth weight (n = 397)				Women born at term (n = 381) <sup>b</sup>	
	Before adjusting for birth weight		Adjusted for birth weight <sup>d</sup>		Adjusted for birth weight	
	$\beta$	(95% CI)	$\beta$	(95% CI)	$\beta$	(95% CI)
	<hr/>					
Triglycerides	11.1	(3.6, 19.1)	9.5	(2.1, 17.4)	10.2	(2.6, 18.5)
HDL-C	-3.2	(-5.9, -0.6)	-3.6	(-6.3, -0.9)	-4.2	(-7.0, -1.4)
LDL-C	0.9	(-5.6, 7.5)	0.0	(-6.6, 6.6)	-1.8	(-8.6, 5.1)
Total cholesterol	-0.7	(-8.3, 6.8)	-2.1	(-9.7, 5.5)	-3.6	(-11.5, 4.3)
ApoB	1.0	(-3.8, 5.8)	0.5	(-4.3, 5.4)	-0.2	(-5.2, 4.8)
CRP	5.0	(-12.3, 25.7)	3.4	(-13.9, 24.1)	0.4	(-16.7, 21.0)

Abbreviations: ApoB, apolipoprotein B; CRP, C-reactive protein; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol

<sup>a</sup>All models were adjusted for age, education, physical activity, smoking, and BMI at blood draw

<sup>b</sup>Gestational age  $\geq 37$  weeks

<sup>c</sup>Associations are expressed as the % difference in the geometric mean for  $\log_n$ -transformed outcomes (mg/dL triglycerides and mg/L CRP) or the difference in mean values (mg/dL of HDL-C, LDL-C, total cholesterol, and ApoB) in the exposed compared with the unexposed group.

<sup>d</sup>Additionally adjusted for participant's gestational age at birth

Table 5. Adjusted<sup>a</sup> Odds Ratios (OR) for selected lipids and CRP by *in utero* exposure to tobacco smoke among MoBa participants

Outcomes	Base sample (n = 479)					All women (n = 823) <sup>b</sup>				
	n <sup>c</sup>	Before adjusting		Adjusted for		Before adjusting		Adjusted for		
		for BMI		BMI		for BMI		BMI		
		OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	
Triglycerides $\geq$ 200 mg/dL	38	2.8	(1.4, 5.7)	2.5	(1.3, 5.1)	72	2.5	(1.3, 4.7)	2.2	(1.2, 4.3)
HDL-C < 50 mg/dL	33	2.4	(1.1, 5.1)	2.3	(1.1, 5.0)	61	2.1	(1.1, 4.0)	2.0	(1.0, 4.1)
Total cholesterol $\geq$ 240 mg/dL	97	1.4	(0.9, 2.3)	1.4	(0.8, 2.3)	159	1.4	(0.9, 2.2)	1.4	(0.9, 2.2)
CRP $\geq$ 8 mg/L	106	1.6	(1.0, 2.6)	1.2	(0.7, 2.1)	197	1.6	(1.1, 2.5)	1.2	(0.8, 2.0)

Abbreviations: CRP, C-reactive protein; HDL-C, high density lipoprotein cholesterol

<sup>a</sup>All models were adjusted for age, education, physical activity, and smoking

<sup>b</sup>Logistic regression models weighted for sampling probability

<sup>c</sup>Number of participants classified as having the outcome

Table 6. Adjusted<sup>a</sup> ORs for selected clinical chemistries by *in utero* exposure to tobacco smoke and BMI (<25 and ≥25 kg/m<sup>2</sup>) among adult women

<i>In utero</i> exposure to tobacco smoke (no, yes)/BMI (<25, ≥25)	n	Triglycerides ≥ 200 mg/dL		HDL-C < 50 mg/dL		Total Cholesterol ≥ 240 mg/dL		CRP ≥ 8 mg/L	
		OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Base sample	479								
Unexposed / BMI <25	195	1.0		1.0		1.0		1.0	
Unexposed / BMI ≥25	150	3.5	(1.2, 10.3)	2.5	(0.9, 6.9)	1.5	(0.9, 2.7)	4.0	(2.2, 7.1)
Exposed / BMI <25	58	4.7	(1.4, 15.7)	4.4	(1.4, 14.4)	1.7	(0.8, 3.5)	1.1	(0.4, 2.8)
Exposed / BMI ≥25	76	6.9	(2.3, 20.6)	3.5	(1.2, 10.7)	1.7	(0.9, 3.4)	6.1	(3.2, 12.0)
All women <sup>b</sup>	823								
Unexposed / BMI <25	318	1.0		1.0		1.0		1.0	
Unexposed / BMI ≥25	265	3.1	(1.2, 7.8)	2.5	(1.0, 6.1)	1.5	(0.9, 2.4)	4.0	(2.4, 6.9)
Exposed / BMI <25	98	3.8	(1.4, 10.8)	3.8	(1.3, 11.1)	1.7	(0.9, 3.2)	1.2	(0.5, 2.6)
Exposed / BMI ≥25	142	5.5	(2.1, 14.0)	3.2	(1.2, 8.2)	1.7	(0.9, 3.1)	6.1	(3.3, 11.2)

<sup>a</sup>All models were adjusted for age, education, physical activity, and smoking

<sup>b</sup>Logistic regression models weighted for sampling probability