



ENVIRONMENTAL
HEALTH
PERSPECTIVES

ehponline.org

**Blood Pressure Changes and Chemical Constituents of
Particulate Air Pollution: Results From The Healthy
Volunteer Natural Relocation (HVNR) Study**

**Shaowei Wu, Furong Deng, Jing Huang, Hongyi Wang,
Masayuki Shima, Xin Wang, Yu Qin, Chanjuan Zheng,
Hongying Wei, Yu Hao, Haibo Lv, Xiuling Lu, Xinbiao Guo**

<http://dx.doi.org/10.1289/ehp.1104812>

Online 19 October 2012



NIEHS
National Institute of
Environmental Health Sciences

National Institutes of Health
U.S. Department of Health and Human Services

**Blood Pressure Changes and Chemical Constituents of Particulate Air Pollution:
Results From The Healthy Volunteer Natural Relocation (HVNR) Study**

Shaowei Wu¹, Furong Deng¹, Jing Huang¹, Hongyi Wang², Masayuki Shima³, Xin Wang¹,
Yu Qin¹, Chanjuan Zheng¹, Hongying Wei¹, Yu Hao¹, Haibo Lv¹, Xiuling Lu¹, Xinbiao
Guo¹

¹Department of Occupational and Environmental Health Sciences, Peking University
School of Public Health, Beijing, China

²Department of Cardiology, Peking University People's Hospital, Beijing, China

³Department of Public Health, Hyogo College of Medicine, Hyogo, Japan

Correspondence:

Xinbiao Guo, MD, PhD,

Department of Occupational and Environmental Health Sciences, Peking University School
of Public Health, No. 38 Xueyuan Road, Beijing 100191, China

Phone: 86-10-82801176

Fax: 86-10-62375580

E-mail: guoxb@bjmu.edu.cn

Short running title: Particulate constituents and blood pressure

Key words: air pollution; blood pressure; hypertension; panel study; particulate matter

Acknowledgments

The authors thank Youcheng Liu from University of North Texas Health Science Center for helpful comments on the manuscript.

Sources of Funding

This work was supported by Grants from the National Natural Science Foundation of China

[No.81072267], the National Key Technologies R&D Program of China [No.2006BAI19B06], the National High Technology Research and Development Program of China [No.2012AA062804], and the Academic Award for Excellent Doctoral Candidates of the Ministry of Education of the People's Republic of China (Shaowei Wu).

Competing Financial Interests: None declared.

Abbreviations list

Al, aluminum	PM ₁₀ , particulate matter with aerodynamic diameter≤10 μm
As, arsenic	PP, pulse pressure
Ba, Barium	Sb, antimony
BIT, Beijing Institute of Technology	SBP, systolic blood pressure
BMI, body mass index	Se, selenium
BP, blood pressure	Sn, stannum
Ca, calcium	SO ₄ ²⁻ , sulfate
Cd, cadmium	Sr, strontium
Cl ⁻ , chloride	Ti, titanium
CI, confidence interval	V, vanadium
CO, carbon monoxide	Zn, zinc
Cr, chromium	
Cu, copper	
DBP, diastolic blood pressure	
EC, elemental carbon	
F ⁻ , fluoride	
Fe, iron	
IQR, interquartile range	
K, potassium	
Mg, magnesium	
Mn, manganese	
Mo, molybdenum	
Na, sodium	
Ni, nickel	
NO, nitric oxide	
NO ₂ , nitrogen dioxide	
NO ₃ ⁻ , nitrate	
NO _x , nitrogen oxides	
OC, organic carbon	
Pb, lead	
PM, particulate matter	
PM _{2.5} , particulate matter with aerodynamic diameter≤2.5 μm	
PM _{2.5-10} , particulate matter with aerodynamic diameter between 2.5 and 10 μm	

Abstract

Background: Elevated blood pressure (BP) has been associated with particulate matter (PM) air pollution, but associations with PM chemical constituents are still uncertain.

Objectives: We investigated associations of BP with various chemical constituents of fine particulate (PM_{2.5}) during 460 repeated visits among a panel of 39 university students.

Methods: Resting BP was measured using standardized methods before and after the university students relocated from a suburban campus to an urban campus with different air pollution contents in Beijing. Air pollution data were obtained from central monitors close to student residences. We used mixed-effects models to estimate associations of various PM_{2.5} constituents with systolic BP (SBP), diastolic BP (DBP) and pulse pressure.

Results: An interquartile range increase of 51.2 $\mu\text{g}/\text{m}^3$ in PM_{2.5} was associated with a 1.08-mmHg (95% CI, 0.17 to 1.99) increase in SBP and a 0.96-mmHg (95% CI, 0.31 to 1.61) increase in DBP on the following day. A subset of PM_{2.5} constituents, including carbonaceous fractions (organic carbon and elemental carbon), ions (chloride and fluoride) and metals/metalloid elements (nickel, zinc, magnesium, lead and arsenic), were found to have robust positive associations with different BP variables, though robust negative associations of manganese, chromium and molybdenum with SBP or DBP also were observed.

Conclusions: Our results support relationships between specific PM_{2.5} constituents and BP. These findings have potential implications for the development of pollution abatement strategies that maximize public health benefits.

Introduction

Both elevated blood pressure (BP) and ambient particulate air pollution have been associated with increased cardiovascular morbidity and mortality (Blacher et al. 2000; Brook et al. 2010; Guo et al. 2010; Kan et al. 2007). An elevation in BP may be an important physiological mechanism linking particulate air pollution and adverse cardiovascular outcomes (Brook and Rajagopalan 2009). Among the size fractions of ambient particulate matter (PM), PM with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$) has been associated with adverse cardiovascular effects (Brook et al. 2010; Kan et al. 2007). Controlled human and animal experiments have verified that exposure to $\text{PM}_{2.5}$ is capable of inducing elevated BP (Brook et al. 2009; Chang et al. 2004; Urch et al. 2005; Zanobetti et al. 2004).

Ambient PM is a mixture of various chemical constituents, including carbonaceous fractions (organic carbon/elemental carbon, OC/EC), ions, and transition metals. These constituents may have different effects on cardiovascular responses (Brook et al. 2010). Although several PM chemical constituents (e.g., carbonaceous fractions) have been associated with pro-hypertensive effects in different populations (Mordukhovich et al. 2009; Urch et al. 2005; Zanobetti et al. 2004), evidence for effects of specific PM chemical constituents on BP is still lacking. Specifically, traffic-related PM may play a distinctive role in cardiovascular responses (Auchincloss et al. 2008; Brook et al. 2009; Delfino et al. 2010; Jia et al. 2012; Wu et al. 2010, 2011a, 2011b). Thus, a hypothesis that a group of PM chemical constituents and related sources may confer greater PM cardiovascular toxicity is reasonable based on the existing literature. We conducted the Healthy Volunteer Natural Relocation (HVNR) study to examine the relationship between various $\text{PM}_{2.5}$ chemical constituents and BP changes in a panel of healthy male university students in Beijing, China before and after their relocation from a

suburban campus to an urban campus with different PM air pollution constituents. We hypothesized that this relocation would substantially change the participants' exposures to ambient PM_{2.5} and chemical constituents associated with local pollution sources, and thus facilitate an analysis of relationships between PM_{2.5} chemical constituents and BP in the study population.

Methods

Study Design

Beijing City covers an area of 16,410 km², has nearly 20,000,000 inhabitants, and is about 160 kilometers from the nearest coastline. More than 5,000,000 vehicles are its main source of urban air pollution. Our study participants consisted of a panel of 41 undergraduate male college students from a university in Beijing (Beijing Institute of Technology, BIT). The BIT has two campuses located in different areas of Beijing (Figure 1). Study participants completed their first two years of undergraduate study (from autumn 2008 to summer 2010) at the BIT Liangxiang campus, which is located in a suburban area (Fangshan District), then moved to the BIT main campus, which is located in an urban area (Haidian District) for their next two years of study (from autumn 2010 to summer 2012). The BIT Liangxiang campus is about 2 kilometers from the nearest freeway. There were several active construction sites within 2 kilometers of the campus during the study, in addition to some industrial facilities located within several kilometers of the campus. In contrast, the BIT main campus is located in the Beijing downtown area, along the northwest inner side of the 3rd ring road that circles the city. There were no substantive construction activities or industrial facilities near the main campus during the study. We used the following inclusion criteria to select participants before the study began: male with

a geographical origin other than Beijing, non-obese, no history of smoking and without pulmonary, cardiovascular, and other chronic diseases. We used a self-administered questionnaire to collect personal information, including name, age, and medical history/health status. We scheduled 12 biweekly study visits for each participant over the entire study, including 4 visits during each of the following three time periods: Suburban Period (April 22 to June 20, 2010) at the BIT Liangxiang campus; Urban Period 1 (September 3 to November 8, 2010) and Urban Period 2 (April 10 to June 12, 2011) at the BIT main campus. The study was approved by the Institutional Review Board of Peking University Health Science Center, and informed consent was provided by each participant before the study began.

Blood Pressure Measurement

Study visits were scheduled between 13:00-15:00 on weekdays to minimize diurnal variation of BP. A trained technician in the hospital affiliated with the BIT Liangxiang campus or BIT main campus performed standardized resting BP measurements during each study visit. Participants rested in a sitting position in a quiet room for at least 10 minutes before upper arm BP was measured using an Omron 705IT automated oscillometric monitor (HEM-759-E, Omron Healthcare Co., Ltd. Kyoto, Japan) at least three times with a 1-minute minimum interval between measurements. In most cases the second and third sets of readings were averaged to calculate systolic BP (SBP) and diastolic BP (DBP) (Rioux et al. 2010). However, if the difference between SBP or DBP values of the second and third measurements was >5 mmHg the BP was considered unstable, and another 1 to 3 measurements would be taken until the difference between the last two measurements was ≤ 5 mmHg. Under this condition, all readings (from the second to the last measurement) within a 5-mmHg range of difference were averaged to calculate the final BP values. Pulse pressure (PP) was calculated as the difference between the

average SBP and DBP values. Weight and height were measured during the first study visit in each time period, and body mass index (BMI) was calculated as weight in kilograms divided by height in meters squared. Two volunteers who did not complete study visits after relocation to the urban campus were excluded from data analysis, leaving a total of 39 participants.

Environmental Data

Air pollution and weather data were measured using standard methods and quality controls at a central monitoring site on the BIT Liangxiang campus (Suburban Period) or BIT main campus (Urban Period 1 & 2). The BIT Liangxiang monitoring site was on the roof of a three-story building (about 10 meters high) without any nearby structures that would obstruct air flow, and BIT main campus monitoring site was on the roof of a five-story building (about 15 meters high) located within 200 meters of the 3rd ring road. The instruments and materials used for air monitoring included: SKC sampling systems for PM_{2.5} mass collection on Teflon filters and quartz-fiber filters (SKC Inc., Eighty Four, PA, USA) and a digital dust monitor for real-time PM_{2.5} concentration measurement (LD-3K; Sibata Scientific Technology Inc., Tokyo, Japan); a model T15n enhanced CO measurer for real-time CO concentration measurement (Langan Products Inc., San Francisco, CA, USA); Ogawa passive samplers for NO_x and NO₂ collection on cellulose fiber filters (Ogawa Air Inc., Osaka, Japan); and a HOBO Pro V2 logger for temperature and relative humidity measurements (Onset Corp., Pocasset, MA, USA). Data on PM with aerodynamic diameter $\leq 10 \mu\text{m}$ (PM₁₀) were obtained from the nearest city air monitoring stations (within 5 kilometers of each campus, respectively) under the supervision of Beijing Municipal Environmental Protection Bureau. Concentrations of coarse PM with aerodynamic diameter between 2.5 and 10 μm (PM_{2.5-10}) were calculated as the difference between the measured PM₁₀ and PM_{2.5} concentrations.

Daily PM_{2.5} mass concentrations were determined by standard weighing procedures before and after the sample collection (Wu et al. 2010), and the PM_{2.5} filters were analyzed in the laboratory for the following chemical constituents: OC and EC in quartz-fiber filters by thermo/optical transmission method (Lab. OC/EC analyzer; Sunset Laboratory Inc., Tigard, OR, USA); sulfate, nitrate, chloride and fluoride in Teflon filters by ion chromatography (Model ICS-2000; Dionex Corp., Sunnyvale, CA, USA); aluminum, calcium, sodium, potassium, magnesium, iron and zinc in Teflon filters by inductively coupled plasma atomic emission spectrometry (Model SPS8000; KCHG Co., Ltd., Beijing, China); and strontium, barium, lead, copper, titanium, nickel, molybdenum, cadmium, vanadium, chromium, manganese, arsenic, selenium, stannum and antimony in Teflon filters by inductively coupled plasma mass spectrometry (Model ELAN DRC II; PerkinElmer Inc., Shelton, CT, USA). We also estimated the concentrations of three additional carbonaceous fractions [primary OC (POC), secondary OC (SOC) and particulate organic matter (POM)], as described in detail in Supplemental Material (see Supplemental Material, p.2). PM_{2.5} constituents were classified according to their chemical nature as carbonaceous fractions, negative ions, transition metals, crustal metals and other metals/metalloid elements. NO_x and NO₂ were collected on cellulose fiber filters and concentrations were determined using a spectrophotometer following the manufacturer's specification (Ogawa & Company USA, Inc., Pompano Beach, FL, USA). Nitric oxide (NO) concentrations were calculated as the difference between the NO_x and NO₂ concentrations.

Statistical Analysis

We first compared the mean BP changes between periods by subject using paired T tests, and then used mixed-effects regression models in SAS 9.2 (SAS Institute, Cary, NC, USA) to estimate associations between exposure variables and BP. Environmental data were matched

with BP data for each subject before analysis. The mixed-effects models included a random intercept for each subject to account for within-subject correlations due to repeated measurements. Base models included individual air pollutants or PM_{2.5} constituents, and were adjusted for age, BMI, temperature and relative humidity as continuous variables, plus linear and quadratic terms for temperature and relative humidity (Mordukhovich et al. 2009), and season, month, day-of-week, hour-of-day and study site as binary or categorical variables. In addition, we included a day-of-study variable and a squared day-of-study variable in the models to account for secular trends in associations between air pollution and BP (Penttinen et al. 2001).

We used three kinds of models to investigate the associations between exposure variables and BP after combining the data over the three time periods. First, we modeled individual air pollutants or PM_{2.5} constituents to estimate associations with BP, with adjustment for the potential confounders listed above. Second, we estimated associations of individual PM_{2.5} constituents and BP with adjustment for total PM_{2.5}. Third, we used a constituent residual model analysis to address collinearity between total PM_{2.5} and PM_{2.5} constituents (Wu et al. 2011a). Specifically, we regressed daily concentrations of each PM_{2.5} constituent on total PM_{2.5} concentrations using a separate linear regression model for each time period to generate the constituent residual for each daily concentration value of the constituent. The constituent residual represents the proportion of the constituent that is uncorrelated with total PM_{2.5}, and therefore can be interpreted as a measure of the independent contribution of each constituent to associations with BP.

To estimate the cumulative effects of exposure, we modeled the average concentrations of exposure variables during the 1 to 5 days before the BP measurement (Hoffmann et al. 2012). We reported results using average concentrations during the preceding 1 to 3 days because most

associations with BP were observed with exposures during this time period. Results are expressed as absolute changes (mmHg) with 95% confidence interval(s) (CI) in BP variables associated with interquartile range (IQR) increases in air pollutants and PM_{2.5} constituents. The statistical significant level was defined as $P < 0.05$ (two sided).

Results

The mean age of the eligible study subjects (n=39) was 20.1 years (range 19-22 years), and their mean BMI was 21.2 kg/m² (range 17.2-24.9 kg/m²). Overall, thirty-four subjects completed all 12 biweekly visits, four completed 11 visits, and one completed 8 visits, resulting in a total of 460 visits. SBP and PP increased over the three time periods whereas DBP levels remained relatively stable (Table 1).

Most air pollutants and PM_{2.5} constituents showed substantial variation over the three time periods (Table 2). In particular, concentrations of gaseous air pollutants (e.g., CO, NO_x and NO₂) and levels of several PM_{2.5} carbonaceous fractions related to traffic (OC, EC, POC and POM based on concentrations or proportions of PM_{2.5} mass) were higher during the Urban Periods than in the Suburban Period.

SBP and PP levels during the Urban Periods were significantly higher than during the Suburban Period, and SBP and PP levels during the Urban Period 2 were also significantly higher than those during the Urban Period 1 (Table 3). There was no significant difference in DBP levels between any two periods. IQR increases in major air pollutant concentrations (PM fractions, NO_x and NO₂) during the 1 to 3 days before study visits showed significant positive associations with SBP or DBP, with more consistent associations observed with DBP than SBP (see Supplemental Material, Table S1). An IQR increase (51.2 µg/m³) in PM_{2.5} during the

preceding day was associated with a 1.08 mmHg (95% CI, 0.17 to 1.99) increase in SBP and a 0.96 mmHg (95% CI, 0.31 to 1.61) increase in DBP. There were no significant associations between the major air pollutants and PP (data not shown).

We found significant associations between SBP or DBP and concentrations of several PM_{2.5} constituents during the preceding day, though some associations that were statistically significant based on single-constituent models were not significant after adjustment for PM_{2.5} or based on constituent residual models (Figure 2). However, all three model estimates indicated significant positive associations between SBP and chloride, nickel, and strontium; between DBP and OC, EC, POC, POM, chloride, fluoride, and lead; and between PP and nickel and magnesium. In addition, all three model estimates indicated significant negative associations between SBP and manganese, chromium, and molybdenum, and between DBP and chromium and molybdenum. We also found significant associations between SBP and zinc during the preceding 3 days and between PP and arsenic during the preceding 3 days based on three different models.

Discussion

We evaluated relationships between BP and various air pollutants and PM_{2.5} chemical constituents under natural exposure settings using a panel study design that repeatedly measured resting BP during three time periods with different ambient air pollution exposures. The relocation of the study population from a suburban campus to an urban campus, which was a normal part of their university education, gave us the opportunity to study associations with specific PM_{2.5} constituents that may have distinct cardiovascular effects (Brook et al. 2009). We observed increases in SBP and PP after relocation from the suburban campus to the urban

campus in a megacity with high air pollution levels, and estimated consistent associations between BP measures and a subset of PM_{2.5} chemical constituents.

Major sources of particulate air pollution in Beijing include road dust, motor vehicle exhaust, industry, incineration, and coal burning (Sun et al. 2004). During recent years, the number of motor vehicles in Beijing has increased rapidly and traffic emissions have become a dominant source of ambient air pollution here (Stone 2008; Wang et al. 2009; Zhou et al. 2010). Evidence linking traffic-related air pollution and cardiovascular outcomes has been growing, especially in urban areas where traffic emissions are one of the major pollution sources (Brook et al. 2009; Delfino et al. 2010; Jia et al. 2012; Wu et al. 2010, 2011a, 2011b). OC and EC (or its surrogate black carbon) are two commonly used indicators of traffic emissions (Delfino et al. 2010; Sun et al. 2004; Wang et al. 2009). Short-term exposures to these carbonaceous particles have been associated with pro-hypertensive effects in cardiac patients (Delfino et al. 2010; Zanobetti et al. 2004). In our study, we found that PM_{2.5} measured in the urban area contained larger proportions of carbonaceous fractions than PM_{2.5} measured in the suburban area (Table 2). This suggests a greater contribution of traffic emissions to the particulate air pollution in the urban area. We found robust positive associations between PM_{2.5} carbonaceous fractions and DBP in our study participants. When OC was sub-classified as POC or SOC, the association appeared to be specific to POC (Figure 2). POC is a representative indicator of particulate organics from fossil fuel combustion sources in the context of traffic-related air pollution (Delfino et al. 2010). Therefore, our findings suggest that PM carbonaceous fractions related to traffic may play an important role in pro-hypertensive effects of PM air pollution.

Several PM_{2.5} metal constituents, including nickel, zinc, magnesium, strontium, lead and arsenic, had robust positive associations with SBP, DBP or PP, while manganese, chromium and

molybdenum had robust negative associations with SBP or DBP. Among these constituents, nickel, zinc, manganese, chromium and lead typically come from industrial emissions, including metallurgical processes, but some of these metals (e.g., zinc, nickel and manganese) may also come from traffic emissions (Loranger and Zayed 1995; Sun et al. 2004). Magnesium usually originates from mineral aerosols that may result from resuspended road dust and long-range transported dust (Sun et al. 2004), while arsenic is typically generated from coal burning (Xie et al. 2006). Relationships between BP and these metals/metalloid elements, especially the transition metals (e.g., nickel, zinc, manganese and cadmium), have been well demonstrated in previous toxicological studies *in vivo* (Fiorim et al. 2011; Kasten et al. 1994; Wang et al. 2002; Yanagisawa et al. 2004; Yang et al. 2007). When delivered to the airways, the transition metals could stimulate the production of reactive oxygen species and then induce airway injury and inflammation which are subsequently followed by a series of cardiopulmonary responses (Gonzalez-Flecha 2004). Epidemiologic evidence linking metals/metalloid elements and BP has been growing. For example, chronic exposures to manganese, arsenic and lead have been associated with increased BP or higher hypertension prevalence in populations (Mordukhovich et al. 2012; Navas-Acien et al. 2007). Specifically, we found robust negative associations between manganese, chromium, and molybdenum and SBP or DBP. Environmental manganese is generally considered to lower hypertension risk (Houtman 1996). A previous animal experiment found that infusion of manganese into conscious, restrained rats resulted in a decrease in BP (Kasten et al. 1994), and results from a recent epidemiologic study also found negative associations between chronic manganese exposure and BP in elders (Mordukhovich et al. 2012). However, our findings on the negative associations between BP and chromium and molybdenum are not supported by previous experimental and epidemiological reports, and a causal

explanation for these findings may need further investigation.

We found small but robust positive associations between chloride and SBP and DBP, less consistent associations between fluoride and DBP, and inconsistent associations between sulfate and nitrate and BP. Sulfate and nitrate are typical secondary pollutants that constitute a significant proportion of the PM_{2.5} mass. In urban areas far from coastlines, the major source for airborne chloride may be from burning polyvinylchloride plastic in refuse dumps, and large amounts of pollutants such as fluoride may also be emitted when trash is being smashed or incinerated (Sun et al. 2004). Experimental studies have demonstrated that chloride was critical in the development of hypertension (Kurtz et al. 1987; Kurtz and Morris 1984; Ziomber et al. 2008). That is, dietary intake of several ions, including sodium, potassium, chloride, were able to induce increased BP in rats (Ziomber et al. 2008), whereas sodium loading without chloride failed to increase BP in animals (Kurtz and Morris 1984) or men (Kurtz et al. 1987). Nevertheless, evidence for the relationship between these airborne ions and BP is still rare and requires further investigation.

Associations with the different BP variables differed over the study. SBP and PP levels tended to increase over time, whereas DBP levels did not. Among the BP variables, PP has been regarded as a stronger predictor of adverse cardiac outcomes, especially in hypertensive patients (Blacher et al. 2000). Several studies have investigated associations between air pollution and BP, but only a few of them have examined PP (Auchincloss et al. 2008; Rioux et al. 2010). Our findings also indicated wide variation among individual study participants, with SBP or PP increased >10 mmHg after relocation to the urban campus in some cases. This suggests that susceptibility to effects of air pollutants may vary substantially among individuals in the general population.

The study has several strengths in addition to the natural relocation study design. Participants were young, healthy volunteers who were nonsmokers and free of any cardiovascular compromises. Therefore, confounding by factors such as age, smoking, disease status, medication use, obesity, etc. was unlikely. We conducted the study in spring and autumn seasons to avoid significant climate changes that might confound associations between air pollutants and BP (Adamopoulos et al. 2010). We used constituent residual models to estimate associations with PM_{2.5} constituents, and adjusted for many potential confounders. However, residual or unmeasured confounding cannot be excluded, and generalizability to other populations may be limited.

This study also has several other limitations. First, we used ambient air pollution data from central monitoring sites rather than personal exposure measures (Brook and Rajagopalan 2009). However, all participants lived in school dormitories within 300 meters of the monitoring site for each campus, and they spent most of their time in naturally ventilated buildings near the monitoring site. Second, as in most epidemiologic studies, we are not able to determine whether the observed associations were due to the measured air pollution constituents, or to other factors that varied along with constituent concentrations (e.g., pollution sources, seasons, locations, or even other pollutants that might be related or correlated to the measured constituents). For example, both seasonal and regional variation in estimated effects of air pollution on cardiovascular outcomes have been reported by previous studies (Brook et al. 2009; Choi et al. 2007). Changes in season or location may result in changes in air pollution sources and the constituents of air pollution, which in turn may influence effects of air pollution on cardiovascular system. As a result, we are not able to differentiate effects of season or location from potential effects of air pollution in the current study. Third, other gaseous air pollutants that

may also contribute to the adverse cardiovascular outcomes, e.g., ozone and sulfur dioxide, were not measured in the current study. Finally, there is a potential confounding effect from progression in university education on BP through changing the participants' stress levels across different periods that could not be excluded. However, in view that the estimated air pollution effects were quite strong after adjusting for various factors related to seasonal and regional factors, we believe this kind of effect would not be able to change our findings materially.

In summary, our findings suggest that specific PM_{2.5} chemical constituents may be associated with BP in healthy adults. These findings also suggest potential linkages between pollution sources and PM-related cardiovascular effects. A better understanding of the responsible PM constituents and sources could potentially lead to more targeted and effective regulations (Brook et al. 2010). This is especially important for the most polluted regions and countries around the world. As the largest developing country, China is now facing the worst air pollution problem in the world along with its rapid economic expansion over the past decades (Kan et al. 2012). Our findings thus may have implications for the development of relevant pollution abatement strategies that maximize benefits to the public health.

References

- Adamopoulos D, Vyssoulis G, Karpanou E, Kyvelou SM, Argacha JF, Cokkinos D, et al. 2010. Environmental determinants of blood pressure, arterial stiffness, and central hemodynamics. *J Hypertens* 28:903-909.
- Auchincloss AH, Diez-Roux AV, Dvorchak JT, Brown PL, Barr RH, Davligus ML, et al. 2008. Associations between recent exposure to ambient fine particulate matter and blood pressure in the Multi-Ethnic Study of Atherosclerosis (MESA). *Environ Health Perspect* 116:486-491.
- Blacher J, Staessen JA, Girerd X, Gasowski J, Thijs L, Liu L, et al. 2000. Pulse pressure not mean pressure determines cardiovascular risk in older hypertensive patients. *Arch Intern Med* 160:1085-1089.
- Brook RD, Rajagopalan S. 2009. Particulate matter, air pollution, and blood pressure. *J Am Soc Hypertens* 3:332-350.
- Brook RD, Urch B, Dvorchak JT, Bard RL, Speck M, Keeler G, et al. 2009. Insights into the mechanisms and mediators of the effects of air pollution exposure on blood pressure and vascular function in healthy humans. *Hypertension* 54:659-667.
- Brook RD, Rajagopalan S, Pope CA 3rd, Brook JR, Bhatnagar A, Diez-Roux AV, et al. 2010. Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. *Circulation* 121:2331-2378.
- Chang CC, Hwang JS, Chan CC, Wang PY, Hu TH, Cheng TJ. 2004. Effects of concentrated ambient particles on heart rate, blood pressure, and cardiac contractility in spontaneously hypertensive rats. *Inhal Toxicol* 16:421-429.
- Choi JH, Xu QS, Park SY, Kim JH, Hwang SS, et al. 2007. Seasonal variation of effect of air pollution on blood pressure. *J Epidemiol Commun Health* 61:314-318.
- Delfino RJ, Tjoa T, Gillen DL, Staimer N, Polidori A, Arhami M, et al. 2010. Traffic-related air pollution and blood pressure in elderly subjects with coronary artery disease. *Epidemiology* 21:396-404.
- Fiorim J, Ribeiro Junior RF, Silveira EA, Padilha AS, Vescovi MVA, et al. 2011. Low-level lead exposure increases systolic arterial pressure and endothelium-derived vasodilator factors in rat aortas. *PLoS ONE* 6:e17117.
- Gonzalez-Flecha B. 2004. Oxidant mechanisms in response to ambient air particles. *Mol Aspects Med* 25:169-182.

- Guo Y, Tong S, Zhang Y, Barnett AG, Jia Y, Pan X. 2010. The relationship between particulate air pollution and emergency hospital visits for hypertension in Beijing, China. *Sci Total Environ* 408:4446-4450.
- Hoffmann B, Luttmann-Gibson H, Cohen A, Zanobetti A, de Souza C, Foley C, et al. 2012. Opposing effects of particle pollution, ozone and ambient temperature on arterial blood pressure. *Environ Health Perspect* 120:241-246.
- Houtman JP. 1996. Trace elements and cardiovascular diseases. *J Cardiovasc Risk* 3(1):18-25.
- Jia X, Song X, Shima M, Tamura K, Deng F, Guo X. 2012. Effects of fine particulate on heart rate variability in Beijing: a panel study of healthy elderly subjects. *Int Arch Occup Environ Health* 85:97-107.
- Kan H, Chen R, Tong S. 2012. Ambient air pollution, climate change, and population health in China. *Environ Int* 42:10-19.
- Kan H, London SJ, Chen G, Zhang Y, Song G, Zhao N, et al. 2007. Differentiating the effects of fine and coarse particles on daily mortality in Shanghai, China. *Environ Int* 33:376-384.
- Kasten TP, Settle SL, Misko TP, Currie MG, Nickolsa GA. 1994. Manganese potentiation of nitric oxide-mediated vascular relaxation. *Eur J Pharmacol* 253:35-43.
- Kurtz TW, Al-Bander HA, Morris RC Jr. 1987. "Salt-sensitive" essential hypertension in men: is the sodium ion alone important? *N Engl J Med* 317:1043-1048.
- Kurtz TW, Morris RC Jr. 1984. Dietary chloride and bicarbonate as determinants of desoxycorticosterone hypertension. *J Hypertens (Suppl.2)*:S371-S373.
- Loranger S, Zayed J. 1995. Contribution of methylecyclopentadienyl manganese tricarbonyl (MMT) to atmospheric Mn concentration near expressway: dispersion modeling estimations. *Atmos Environ* 29:591-599.
- Mordukhovich I, Wilker E, Suh H, Wright R, Sparrow D, Vokonas PS, et al. 2009. Black carbon exposure, oxidative stress genes, and blood pressure in a repeated-measures study. *Environ Health Perspect* 117:1767-1772.
- Mordukhovich I, Wright RO, Hu H, Amarasiriwardena C, Baccarelli A, Litonjua A, et al. 2012. Associations of toenail arsenic, cadmium, mercury, manganese and lead with blood pressure in the Normative Aging Study. *Environ Health Perspect* 120:98-104.
- Navas-Acien A, Guallar E, Silbergeld EK, Rothenberg SJ. 2007. Lead exposure and

- cardiovascular disease—a systematic review. *Environ Health Perspect* 115:472-482.
- Penttinen P, Timonen KL, Tiittanen P, Mirme A, Ruuskanen J, Pekkanen J. 2001. Number concentration and size of particles in urban air: effects on spirometric lung function in adult asthmatic participants. *Environ Health Perspect* 109:319-323.
- Rioux CL, Tucker KL, Mwamburi M, Gute DM, Cohen SA, Brugge D. 2010. Residential traffic exposure, pulse pressure, and C-reactive protein: consistency and contrast among exposure characterization methods. *Environ Health Perspect* 118:803-811.
- Stone R. 2008. Beijing's Marathon run to clean foul air nears finish line. *Science* 321:636-637.
- Sun Y, Zhuang G, Wang Y, Han L, Guo J, Dan M, et al. 2004. The air-borne particulate pollution in Beijing—concentration, composition, distribution and sources. *Atmos Environ* 38:5991-6004.
- Urch B, Silverman F, Corey P, Brook JR, Lukic KZ, Rajagopalan S, et al. 2005. Acute blood pressure responses in healthy adults during controlled air pollution exposures. *Environ Health Perspect* 113:1052-1055.
- Wang S, Paek D, Kim R, Cha B. 2002. Variation of systolic blood pressure in rats exposed to cadmium and nickel. *Environ Res A* 88:116-119.
- Wang X, Westerdahl D, Chen LC, Wu Y, Hao J, Pan X, et al. 2009. Evaluating the air quality impacts of the 2008 Beijing Olympic Games: On-road emission factors and black carbon profiles. *Atmos Environ* 43:4535-4543.
- Wu S, Deng F, Niu J, Huang Q, Liu Y, Guo X. 2010. Association of heart rate variability in taxi drivers with marked changes of particulate air pollution in Beijing in 2008. *Environ Health Perspect* 118:87-91.
- Wu S, Deng F, Niu J, Huang Q, Liu Y, Guo X. 2011a. Exposures to PM_{2.5} components and heart rate variability in taxi drivers around the Beijing 2008 Olympic Games. *Sci Total Environ* 409:2478-2485.
- Wu S, Deng F, Niu J, Huang Q, Liu Y, Guo X. 2011b. The relationship between traffic-related air pollutants and cardiac autonomic function in a panel of healthy adults: a further analysis with existing data. *Inhal Toxicol* 23: 289-303.
- Xie R, Seip HM, Wibetoe G, Nori S, McLeod CW. 2006. Heavy coal combustion as the dominant source of particulate pollution in Taiyuan, China, corroborated by high concentrations of arsenic and selenium in PM₁₀. *Sci Total Environ* 370:409-415.

- Yanagisawa H, Sato M, Nodera M, Wada O. 2004. Excessive zinc intake elevates systemic blood pressure levels in normotensive rats-potential role of superoxide-induced oxidative stress. *J Hypertens* 22:543-550.
- Yang HT, Chou HJ, Han BC, Huang SY. 2007. Lifelong inorganic arsenic compounds consumption affected blood pressure in rats. *Food Chem Toxicol* 45:2479-2487.
- Zanobetti A, Canner MJ, Stone PH, Schwartz J, Sher D, Eagan-Bengston E, et al. 2004. Ambient pollution and blood pressure in cardiac rehabilitation patients. *Circulation* 110:2184-2189.
- Zhou Y, Wu Y, Yang L, Fu L, He K, Wang S, et al. 2010. The impact of transportation control measures on emission reductions during the 2008 Olympic Games in Beijing, China. *Atmos Environ* 44:285-293.
- Ziomber A, Machnik A, Dahlmann A, Dietsch P, Beck FX, Wagner H, et al. 2008. Sodium-, potassium-, chloride-, and bicarbonate-related effects on blood pressure and electrolyte homeostasis in deoxycorticosterone acetate-treated rats. *Am J Physiol Renal Physiol* 295:F1752-F1763.

Tables

Table 1. Descriptive statistics on blood pressure by time period

Variable	Period	N	Mean±SD	Range	<i>P</i> -value ^a
SBP, mmHg	Suburban Period	153	117.0±10.1	91-146	0.010
	Urban Period 1	156	119.1±11.4	90-154	
	Urban Period 2	151	120.9±12.1	89-156	
	Total	460	119.0±11.3	89-156	
DBP, mmHg	Suburban Period	153	65.8±7.3	41-91	0.847
	Urban Period 1	156	66.2±6.9	54-94	
	Urban Period 2	151	65.9±6.7	54-89	
	Total	460	66.0±7.0	41-94	
PP, mmHg	Suburban Period	153	51.2±8.3	29-75	<0.001
	Urban Period 1	156	52.9±7.9	32-71	
	Urban Period 2	151	55.0±9.3	32-73	
	Total	460	53.0±8.7	29-75	

^a *P*-value for analysis of variance for repeated measurements.

Table 2. Descriptive statistics on environmental variables over the study

Variable	Suburban Period		Urban Period 1		Urban Period 2		IQR ^c
	Mean±SD ^a	% of PM _{2.5} ^b	Mean±SD	% of PM _{2.5}	Mean±SD	% of PM _{2.5}	
PM ₁₀ , µg/m ³	135.4±64.1		111.3±75.0		129.4±87.2		66.0
PM _{2.5-10} , µg/m ³	56.1±47.5		36.3±29.3		69.5±62.8		42.3
PM _{2.5} , µg/m ³	82.0±46.6		78.1±72.5		59.9±40.3		51.2
Carbonaceous fractions							
OC, µg/m ³	10.2±5.5	15.5	12.6±9.0	20.6	10.6±3.9	24.0	3.7
EC, µg/m ³	2.0±1.2	2.88	3.3±2.6	4.65	1.7±0.9	3.64	1.5
POC, µg/m ³	4.4±2.3	6.26	8.2±6.3	11.4	6.0±3.1	13.2	4.1
SOC, µg/m ³	5.9±4.9	9.30	4.4±3.3	9.20	4.6±2.4	10.8	1.9
POM, µg/m ³	16.4±8.8	24.9	20.1±14.4	32.9	16.9±6.2	38.4	5.9
Ions							
SO ₄ ²⁻ , µg/m ³	16.8±15.0	18.4	12.0±14.2	13.0	8.9±9.0	13.1	10.0
NO ₃ ⁻ , µg/m ³	2.9±2.8	2.98	3.5±4.7	3.21	2.1±2.8	2.57	2.6
Cl ⁻ , µg/m ³	1.0±1.1	1.18	1.5±1.8	1.73	0.9±1.2	1.40	1.2
F ⁻ , ng/m ³	31.9±32.3	0.052	57.9±48.2	0.107	51.7±47.1	0.099	57.0
Transition metals							
Fe, µg/m ³	0.8±0.4	1.21	0.7±0.4	1.16	0.8±0.9	1.30	0.5
Zn, µg/m ³	0.4±0.3	0.52	0.4±0.4	0.50	0.4±0.3	0.62	0.3
Mn, ng/m ³	65.9±45.0	0.116	60.4±30.6	0.110	48.3±26.9	0.096	35.5
Ti, ng/m ³	48.5±33.3	0.071	38.0±24.2	0.068	52.6±79.8	0.093	29.5
Cu, ng/m ³	30.1±24.3	0.038	37.5±35.4	0.049	26.3±22.9	0.042	26.9
Cr, ng/m ³	18.5±22.5	0.036	8.9±5.6	0.019	8.5±6.7	0.021	5.1
Ni, ng/m ³	3.7±2.2	0.006	3.4±3.5	0.006	3.3±2.5	0.007	2.1
Cd, ng/m ³	2.6±2.8	0.003	3.2±3.4	0.004	2.4±2.7	0.004	2.5
V, ng/m ³	3.1±1.8	0.004	1.6±1.1	0.003	1.7±1.9	0.003	1.3
Mo, ng/m ³	1.9±2.1	0.004	1.3±0.8	0.002	0.8±0.7	0.002	0.9

Table 2. (continued)

Variable	Suburban Period		Urban Period 1		Urban Period 2		IQR ^c
	Mean±SD ^a	% of PM _{2.5} ^b	Mean±SD	% of PM _{2.5}	Mean±SD	% of PM _{2.5}	
Crustal metals							
K, µg/m ³	1.1±0.9	1.32	1.4±0.9	2.59	1.0±0.9	1.71	1.0
Ca, µg/m ³	1.1±0.8	1.64	0.8±0.4	1.61	1.0±0.7	1.79	0.7
Al, µg/m ³	1.0±0.8	1.46	0.5±0.2	0.97	0.8±1.6	1.23	0.5
Na, µg/m ³	0.6±0.6	0.82	0.8±0.5	1.60	0.6±0.4	1.46	0.4
Mg, ng/m ³	292.4±249.1	0.43	194.2±98.9	0.37	318.4±341.6	0.57	148.3
Ba, ng/m ³	14.0±8.2	0.021	17.2±10.0	0.033	14.6±12.4	0.027	11.8
Sr, ng/m ³	9.5±6.4	0.015	6.0±3.1	0.011	6.6±5.0	0.012	4.8
Other metals/metalloid elements							
Pb, ng/m ³	127.7±89.6	0.158	125.0±122.5	0.153	119.2±111.7	0.181	127.7
As, ng/m ³	16.0±19.9	0.018	17.1±24.0	0.018	32.6±35.2	0.058	19.5
Sn, ng/m ³	7.3±5.7	0.009	9.2±9.5	0.012	6.8±5.8	0.011	6.8
Sb, ng/m ³	6.6±5.6	0.008	8.6±7.6	0.011	6.0±4.9	0.010	6.1
Se, ng/m ³	7.3±5.9	0.009	5.4±4.6	0.008	3.9±3.4	0.006	4.4
CO, ppm	0.90±0.33		1.67±0.60		1.47±0.45		0.74
NO _x , ppb	41.6±18.1		70.7±39.4		48.3±17.7		35.1
NO ₂ , ppb	24.1±8.0		38.0±14.8		31.7±11.8		14.2
NO, ppb	17.4±12.4		32.8±29.3		16.5±10.9		26.8
Temperature, °C	23.8±5.3		17.3±6.5		22.4±4.5		-
Relative humidity, %	47.5±15.9		55.7±15.7		37.9±14.5		-

^a Data are presented as daily mean levels.

^b Calculated as daily mean percents of PM_{2.5} mass.

^c Interquartile range values of air pollutant concentrations during the preceding 1 day before the BP measurement.

Table 3. Mean BP changes (mmHg) between periods by subject

Period pair	Variable	N (subjects) ^a	Mean±SD ^b	Range	<i>P</i> -value ^c
Urban Period 1-Suburban Period	SBP	36	2.2±5.6	-11.8-12.8	0.024
	DBP	36	0.4±3.3	-7.3-9.9	0.472
	PP	36	1.8±4.7	-11.2-12.7	0.026
Urban Period 2-Suburban Period	SBP	34	3.5±5.3	-8.2-15.9	0.001
	DBP	34	-0.1±2.1	-3.9-4.6	0.711
	PP	34	3.7±5.1	-6.0-13.7	<0.001
Urban Period 2-Urban Period 1	SBP	37	1.7±4.6	-9.8-13.2	0.035
	DBP	37	-0.4±3.3	-8.1-6.4	0.291
	PP	37	2.1±3.8	-9.4-9.1	0.001

^a Each comparison is restricted to the subjects who had completed all the 8 study visits over the two periods being compared.

^b Presented as the mean difference between the two periods (the latter period of the pair is used as the reference).

^c *P*-value for paired T test.

Figure Legends

Figure 1. Map of the two BIT campuses involved in the study.

Figure 2. BP changes associated with IQR increases in PM_{2.5} constituents at concentration during the preceding day before the BP measurement. Estimates are adjusted for age, BMI, season, month, day-of-study, squared day-of-study, day-of-week, hour-of-day, study site, and temperature and relative humidity in linear and quadratic terms. The vertical lines represent the 95% CIs. For constituent-PM_{2.5} joint models, we used the main effect estimates of PM_{2.5} constituents for result presentation. (A) SBP; (B) DBP; (C) PP.

Diamonds: single-constituent model; triangles: constituent-PM_{2.5} joint model; squares: single-constituent residual model.




Figure 1
180x132mm (96 x 96 DPI)

