

**ehp**

**ENVIRONMENTAL  
HEALTH  
PERSPECTIVES**

ehponline.org

**Neighborhood Effects on Heat Deaths:  
Social and Environmental Predictors of Vulnerability in  
Maricopa County, Arizona**

**Sharon L. Harlan, Juan H. Declet-Barreto,  
William L. Stefanov, Diana B. Petitti**

**<http://dx.doi.org/10.1289/ehp.1104625>**

**Online 16 November 2012**



**NIEHS**  
National Institute of  
Environmental Health Sciences

National Institutes of Health  
U.S. Department of Health and Human Services

## Neighborhood Effects on Heat Deaths:

Social and Environmental Predictors of Vulnerability in Maricopa County, Arizona

Sharon L. Harlan<sup>1</sup>, Juan H. Declet-Barreto<sup>1</sup>, William L. Stefanov<sup>2</sup>, Diana B. Petitti<sup>3</sup>

<sup>1</sup> School of Human Evolution and Social Change, Arizona State University, Tempe, Arizona, USA

<sup>2</sup> Science Applications Research and Development, Jacobs/ESCG, NASA Space Center, Houston, Texas, USA

<sup>3</sup> Department of Biomedical Informatics, Arizona State University, Scottsdale, Arizona, USA

**Address correspondence:** S. Harlan, School of Human Evolution and Social Change, Arizona State University, Tempe, AZ 85284-2402. Telephone: 480-727-6780. Fax: 480-965-7671. E-mail: [sharon.harlan@asu.edu](mailto:sharon.harlan@asu.edu)

**Running title:** Neighborhood Effects on Heat Deaths

**Key words:** climate, GIS, heat mortality, neighborhoods, remote sensing, vulnerability

**Abbreviations:** AC: air conditioning; GIS: Geographic Information System; HVI: heat vulnerability index; LISA: Local Indicator of Spatial Association

**Acknowledgements:** This research was supported by the National Science Foundation (Grant No. GEO-0816168). We thank Sarah Santana, Vjollca Berisha, and Mare Schumacher of the Maricopa County Department of Public Health; William Johnson, Wade Bannister, and Miwa Edge at Arizona State University's Center for Health Information Research; research assistant Shuo Yang; Scott Long, Edward Hackett, and our co-investigators on the Urban Vulnerability to Climate Change project.

The authors declare they have no competing financial interests.

**Abstract**

**Background:** Most heat-related deaths occur in cities and future trends in global climate change and urbanization may amplify this trend. Understanding how neighborhoods affect heat mortality fills an important gap between studies of individual susceptibility to heat and broadly comparative studies of temperature-mortality relationships in cities.

**Objectives:** We estimated neighborhood effects of population characteristics and the built and natural environments on deaths due to heat exposure in Maricopa County, Arizona (2000-2008).

**Methods:** We used U.S. Census data and remotely sensed vegetation and land surface temperature to construct indicators of neighborhood vulnerability and a Geographic Information System to map vulnerability and residential addresses of people who died from heat exposure in 2,081 census block groups. Binary logistic regression and spatial analysis were used to associate deaths with neighborhoods.

**Results:** Neighborhood scores on three factors – Socioeconomic Vulnerability, Elderly/Isolation, and Unvegetated Area – varied widely throughout the study area. The preferred model (based on fit and parsimony) for predicting the odds of one or more deaths from heat exposure within a census block group included the first two factors and surface temperature in residential neighborhoods, holding constant population size. Spatial analysis identified clusters of neighborhoods with the highest heat vulnerability scores. A large proportion of deaths occurred among people, including homeless persons, who lived in the inner cores of the largest cities and an industrial corridor.

**Conclusions:** Place-based indicators of vulnerability complement analyses of person-level heat risk factors. Surface temperature might be used in Maricopa County to identify the most

heat-vulnerable neighborhoods but more attention to the socio-ecological complexities of climate adaptation is needed.

## Introduction

Deaths caused by extremely hot weather are a major public health concern and temperature-mortality relationships in many cities have been intensively studied over the past decade (McMichael et al. 2008). Global climate models predict higher temperatures and more frequent, longer, and intense extreme heat events over most of the world, and this has greatly increased estimates of future deaths due to heat (IPCC 2007). Most heat deaths occur in cities (Kovats and Koppe 2005) and this trend is likely to continue for socio-environmental reasons: (1) cities in many types of climate regimes experience heat waves (Anderson and Bell 2011), (2) cities, especially in developing nations of Asia and Africa, are growing rapidly (UNFPA 2007), (3) large numbers of vulnerable populations, such as the poor, homeless, and elderly, reside in cities and (4) cities are warmer than surrounding rural areas due to urban heat islands (Heisler and Brazel 2010).

Many studies on urban heat-related mortality examine individual-specific risk factors. Commonly identified physiological risks, summarized by Balbus and Malina (2009), are advanced or young ages, underlying diseases, disability, and pregnancy. Cardiovascular disease and several other illnesses are risk factors for heat-related death (McMichael et al. 2006). Deaths from heat exposure also occur among people who lack access to cool environments or are physically active in hot weather (Kilbourne et al. 1982). Living in poverty is a key individual risk factor for deaths related to heat because it decreases the odds of access to medical care and protective resources (Balbus and Malina 2009).

Neighborhoods are “ecological units nested within successively larger communities” (Sampson et al. 2002:445) and neighborhood effects on human development and life cycle events, individual behavior, social outcomes, and health risks have long been studied in the

social and health sciences (Sampson et al. 2002; Pickett and Pearl 2001; Brooks-Gunn et al. 1997). Many indicators of poor health, such as low birth weight (Debbink and Badder 2011), obesity (Mujahid et al. 2008), and coronary heart disease (Diez-Roux et al. 2001) are spatially clustered within neighborhoods. Socioeconomic context is important because many poor neighborhoods lack institutional capacities for education, health care, and employment (Wilson 1987) and have poor quality housing. Many minorities live in low-income areas with high levels of social isolation and concentrated disadvantage (Sampson 2009). Lochner et al. (2002) found that economic deprivation and low levels of social capital defined as trust, social ties, and reciprocity were associated with higher all-cause mortality in Chicago neighborhoods.

There are numerous suspected pathways through which living in a poor neighborhood could lead to higher risks of heat deaths. An emergent literature on community and environmental heat risk factors has examined differences in socioeconomic and social network variables (Klinenberg 2002), the built environment (Kovats and Koppe 2005; O'Neill et al. 2005), and the biophysical environment (Harlan et al. 2006; Ruddell et al. 2010). Studies in St. Louis (Smoyer 1998), Chicago (Klinenberg 2002; Johnson et al. 2012), and Philadelphia (Johnson et al. 2009) have investigated differences in deaths by neighborhood during heat waves. Although such studies of neighborhood-specific deaths are rare, investigations of fine-scale variation in social and environmental neighborhood contexts within cities may fill an important gap between studies of individual susceptibility to heat and broadly comparative studies of temperature-mortality relationships in cities across different climate regimes and geographic locations, such as McMichael et al. (2008).

In this study we investigated neighborhood effects on heat exposure deaths in Maricopa County, Arizona over a nine-year period (2000-2008). Extremely high temperatures occur almost daily for six months a year in this desert climate. The decedents were identified by the county health department using a surveillance system designed specifically to detect deaths caused by or related to environmental (weather-related) heat. Our question was, what characteristics of urban neighborhoods affect the risk of residents dying from extreme heat?

We traced neighborhood effects through two pathways that are connected to individual health outcomes. Neighborhood *compositional effects* are aggregated population characteristics, especially those related to socioeconomic status and age, which increase the risks of heat mortality. Neighborhood *contextual effects* are characteristics of the built and natural environments in which people perform their daily activities. Contextual influences on urban heat mortality, such as land cover and microclimates, have been studied much less than population characteristics but they may be important markers for places that are vulnerable to heat.

In metropolitan regions, there is a strongly increasing temperature gradient from rural areas to central cities known as the urban heat island effect (Heisler and Brazel 2010). Diurnal temperature cycles vary substantially among neighborhoods and differences in neighborhood land cover characteristics – vegetation, impervious surface, bare soil, water – and building structures are drivers of local temperatures (Buyantuyev and Wu 2010). More vegetation lowers local temperature because green areas – trees, other plants, and lawns in yards and parks – cool the air and ground surfaces through shading and evapotranspiration. Studies have documented neighborhood variations in land cover and/or temperature within

Phoenix (Jenerette et al. 2011; Buyantuyev and Wu 2010), Detroit (Zhang et al. 2011), Philadelphia (Johnson et al. 2009), and Baltimore (Huang et al. 2011).

Amount of vegetative cover and land surface temperature, e.g. the temperature one would sense by touching the land surface, are biophysical properties of the outdoor neighborhood environment that are associated with human heat stress (Harlan et al. 2006) and heat-related deaths (Johnson and Wilson 2009; Johnson et al. 2012) in specific geographic contexts. Remotely sensed data from satellites provides synoptic spatial coverage of vegetation and land surface temperature for entire neighborhoods but the temporal coverage of a given scene is limited to a single point in time, much like a snapshot from a hand-held camera. On the other hand, air temperature—measured approximately 2 meters above the land surface—is monitored continuously at weather stations but their proximity to residential neighborhoods is highly variable. Heat mortality studies that use air temperature measurements from a single station or an average temperature of several stations ignore spatial variability.

Our study used two approaches to identify heat-vulnerable neighborhoods. First, we quantified neighborhood thermal properties with spatially continuous remotely sensed metrics to provide measures of difference in neighborhood environmental contexts that cannot be obtained from spatially discontinuous air temperature measurements. Second, we used Reid and colleague's (2009) cumulative Heat Vulnerability Index (HVI) developed for census tracts in the U.S. and subsequently applied to predicting morbidity and mortality in some zip codes in five states (Reid et al. 2012). We re-calculated and mapped Reid's index at the finer spatial resolution of census block groups in the urbanized area of Maricopa County. HVI is an index of vulnerability that sums aggregated neighborhood population

characteristics, prevalence of AC, and vegetation in order to *cumulate* the known risk factors for heat mortality. The rationale for mapping vulnerability is to locate spatially the distributions of social disadvantages and environmental hazards so that effective adaptive strategies can be developed for populations that are most likely to be exposed to them (Cutter and Finch 2007). Mapping may help policymakers identify areas of cities that are at high risk for heat-related mortality, in need of preventive actions during hot weather, or high priorities for environmental modifications.

One of our original contributions is to perform analyses that evaluate the relationship between the HVI and heat-associated mortality in Maricopa County residential neighborhoods. Another is to assess differences among models that use neighborhood HVI or land surface temperature (hereafter referred to as surface temperature) to predict the odds of heat deaths. The sparse literature leads us to expect that neighborhood effects of social variables, green spaces, and temperature are all positively related to heat mortality.

## **Methods**

***Study area.*** The population center of Maricopa County is the Phoenix metropolitan area, which comprises more than 20 contiguous municipalities and three tribal communities. Phoenix is the largest city but several other medium and large cities are also located within our study area, numbering 3.8 million people in 2010.

Chronically hot weather causes many heat fatalities in the desert southwest (mean daily high temperature in Phoenix = 39.4°C during May-September, 2000-2008). Arizona has led the nation in deaths from heat exposure (CDC 2006) and there is a strong positive temperature-mortality relationship in Maricopa County (Yip et al. 2008). In some Phoenix

neighborhoods, a heat stress index was above NOAA's danger threshold for 18% of all hours during summer 2003 (Harlan et al. 2006).

There is no single agreed upon definition of neighborhood, despite widespread interest in neighborhood effects (Sampson 2002). Reid et al. (2009), Smoyer (1998), and Johnson et al. (2009) used census tract boundaries in their studies of intra-urban heat vulnerability. We used block group boundaries from the 2000 U.S. Census to define neighborhoods for our study because they are subdivisions of larger census tracts and therefore more socially and ecologically homogeneous than tracts (see also Johnson et al. 2012). We included 2,081 block groups in our analysis after eliminating 19 with fewer than 10 residents, 11 with missing data on any variable in the analysis, and two containing only institutionalized populations in a state hospital or juvenile corrections facility.

***Neighborhood composition.*** We considered variables from the 2000 U.S. Census used by Reid et al. (2009) to derive a national HVI based on literature showing strong relationships with individual heat mortality, specifically, percentages of populations that were ethnic minorities, living below the poverty line, had less than a high school education, elderly (65 or older), living alone, and that were both elderly and living alone. In addition, our analysis included percentage of Latino immigrants, defined as foreign-born Spanish-only speakers, because they were 10.4% of Maricopa County's population in 2000 and may be vulnerable to environmental health problems, in part due to conditions in their neighborhoods (Grineski et al. 2007). We did not use diabetes prevalence when constructing our block-group-level HVI because diabetes data were only available at the state level.

*Neighborhood context.*

We defined neighborhood contextual effects related to heat vulnerability as the cooling capacities of indoor (prevalence of AC) and outdoor (vegetation and surface temperature) environments. These measures represent average differences in neighborhood environments to which people are exposed that may influence exposure to heat, but do not provide information on individual-level exposure to heat.

The Maricopa County Tax Assessor records whether residential parcels have central AC or an evaporative cooler, an apparatus that cools home interiors by evaporating water. Evaporative coolers are used in some older and low-income homes in Arizona instead of AC but they are not effective when the drier air of June and July transitions to the higher humidity of the August monsoon in central Arizona (Kalkstein and Kalkstein 2004). We spatially joined the Assessor's 2010 parcel registry using a Geographic Information System (GIS) to calculate the percentage of single-family houses in each census block group that did not have central AC or an evaporative cooler (these are not reported separately in the data source). The available data do not account for houses with window units only or for AC/cooler availability in apartment complexes.

We calculated surface temperature and the Normalized Difference Vegetation Index (NDVI; Tucker 1979)—a numerical measure of green vegetation abundance at the pixel scale based on the light reflectance properties of vegetation—for each pixel in the study area from a Landsat scene (30 meter resolution) acquired on 24 July 2000 at 1055 hours local time. NDVI and surface temperature pixel scores were aggregated to block group boundaries to measure mean outdoor thermal properties of neighborhoods. We also used NDVI and surface temperature standard deviations (SD) because the spatial variability of vegetation

and temperatures within neighborhoods are important properties of neighborhoods' thermal profiles (Jenerette et al. 2007).

We chose July measures because half the heat exposure deaths occurred during this month in each year of the study. In the Phoenix area, air and surface temperatures differ widely and consistently across neighborhoods and differences are greater during the summer and even greater during heat waves (Harlan 2006; Ruddell 2010; Jenerette 2011).

***Heat deaths.*** The Maricopa County Department of Public Health (MCDPH) has a surveillance system specifically designed to identify heat-caused and heat-related deaths associated with weather (MCDPH 2011). Their system collects information from several sources: the Office of Medical Examiner case list, which identifies suspected heat-associated deaths referred to the Medical Examiner; the Arizona Department of Health Services, which identifies deaths with environmental heat mentioned on death certificates; local hospitals, which identify suspected heat-associated morbidity/mortality cases; and media reports of heat deaths. A daily multi-causes of death search of the electronic death certificate database identifies conditions associated with environmental heat: International Classification of Disease, Tenth Revision codes (WHO 2007) (X30-exposure to excessive natural heat, T67.X-effects of heat and light, P81.0-environmental hyperthermia of newborn) and key phrases in the text fields for causes and underlying causes of death (heat exposure, environ, exhaustion, sun, heat stress, heat stroke, hyperthermia). The surveillance system designated 455 deaths caused by or related to environmental heat exposure that occurred between 2000 and 2008 (annual mean=50.6; SD=11.6; range=25 to 85). The age distribution of the decedents was: 0-19 (5.1%); 20-39 (13.0%); 40-64 (45.4%); 65-74 (14.3%); 75+ (22.1%).

We geocoded 278 residential street addresses of these decedents and calculated the incidence of heat deaths for each block group. Other residential addresses were either outside Maricopa County or unknown. The MCDPH considers unknown street addresses in the county to be a marker of homelessness. We were able to geocode a place of injury address from the death certificate for 50 heat decedents with unknown residential street addresses in Maricopa County (presumed homeless). These deaths were analyzed separately.

*Analysis.* Means, standard deviations, and Pearson correlations for variables related to heat vulnerability are reported in Table 1. Variables were coded so higher scores denote higher vulnerability. We performed a Principal Components Analysis on the correlation matrix for the HVI variables using varimax rotation and standard statistical criteria, which yielded three principal factors (Eigenvalues>1.5; total explained common variance=79.8%). Standardized scores for each factor (mean=0; SD=1) were assigned to block groups. Following Reid et al. (2009), each factor score was also divided into 6 equal increments of +/- 1.0 SD and assigned integer values 1 ( $\geq 2$  SD below mean) to 6 ( $\geq 2$  SD above mean). HVI for block groups were calculated by summing the integer scores for all three factors (possible total scores of 3-18, actual scores ranged from 5-16, median=10; mean=10.6; SD=1.6). We mapped the index onto the study area and then overlaid point estimates of the residential addresses of decedents at the centroid of block groups.

We estimated binary logistic regressions (SAS version 9.2) to predict the odds that a block group was home to one or more decedents who died from heat exposure (1=yes; 0=no). There were zero deaths in 88.6% (1,843) of the block groups (204=1 death; 31=2 deaths; 3=3 deaths). Due to small numbers of a rare event (Cromley and McLafferty 2002), we did not attempt to model mortality rates. There were too few block groups with more than one death to

explain variation in number of deaths. Population size in 2000 (best available estimate for 2000-2008) was included in each model to control for the number of people at risk in each neighborhood.

To evaluate neighborhood effects on the odds of heat deaths, we tested eight models with different combinations of covariates: the HVI additive index score, each individual HVI factor score from the PCA, all three HVI factor scores, percent of homes with no AC/coolers, land surface temperature, and a combination of all HVI factor scores and surface temperature. We used the Bayesian Information Criterion (BIC) to make pairwise comparisons between the binomial regression models. The BIC assesses model fit penalized for the number of estimated parameters. For comparisons of models with different covariates, the more parsimonious model with the smaller BIC is preferred (Raftery 1995:134). Raftery's (1995:139) widely cited criteria define "very strong" evidence for a preferred model as BIC difference  $>10$  (analogous to  $p \leq 0.01$ ), "strong" evidence = 6-10 (analogous to  $p \leq 0.05$ ), "positive" evidence = 2-6, and "weak" evidence = 0-2.

We conducted sensitivity analysis by comparing HVI predictions of heat deaths with predictions of an alternative indicator of social vulnerability, which has been used to identify neighborhoods that are intergenerational "poverty traps." The index of concentrated disadvantage (Sampson 2009) is comprised of six neighborhood-level census variables: percentages of individuals that are ethnic minorities, unemployed, under age 18, and percentages of households below poverty, receiving welfare, female-headed. HVI and concentrated disadvantage tap a similar underlying construct of neighborhood social disadvantage but they have only ethnicity and poverty variables in common. The other four variables in the concentrated disadvantage index are not known heat risk factors.

In order to assess whether the July 24 2000 Landsat scene represented a stable pattern of variability in regional temperatures, we also estimated models with Landsat-based surface temperatures recorded on two alternative dates.

Finally, we used spatial statistics to measure clustering of neighborhoods with similar HVI index scores. We calculated the Local Indicator of Spatial Association (LISA, Anselin 1995) to measure spatial autocorrelation of the index. LISA scores for each block group represent the locally-disaggregated component of the global Moran's I (de Smith et al. 2007).

## **Results**

Results of the Principal Components Analysis are reported in Table 2, which shows the HVI variables loadings on three principal factors: (1) Socioeconomic Vulnerability, (2) Elderly/Isolation, and (3) Unvegetated Area. In contrast with Reid et al. (2009), who identified AC as a separate principal factor for their national HVI, AC/cooler loaded with socioeconomic variables in our analysis. In addition, Unvegetated Area was an independent factor in Maricopa County, but was a component of the social/environmental factor in the national HVI. Another difference was that the variable age 65 or older loaded with diabetes prevalence to form an elderly/diabetes factor in Reid's national analysis, whereas in our study, advanced age was a component of the Elderly/Isolation factor, along with living alone and the interaction between age and living alone. Higher percentages of elderly residents and individuals living alone tended to occur in the same neighborhoods. (We did not include diabetes prevalence in our factor analysis.) Otherwise, except for differences noted above, the Maricopa County HVI factor structure was similar to the factor structure for the national HVI derived by Reid et al (2009).

The map of metropolitan Phoenix (Figure 1) illustrates HVI integer scores for block groups from lowest to highest vulnerability. Neighborhoods in the inner cores of the two largest cities (Phoenix and Mesa) and a corridor in the northwestern suburbs (Glendale to Sun City) had the highest scores. Lowest scores were in urban fringe neighborhoods to the east and west of major municipal centers. The distribution of heat deaths (Figure 1) shows that residential neighborhoods of decedents who died from heat exposure were located throughout the metropolitan area, but block groups with two or three deaths were more common in higher vulnerability areas.

Results are reported in Table 3 for four binary logistic regression models predicting the odds of heat exposure deaths for neighborhood residents. The predictors were HVI integer score of block group (Model 1), three normalized factor scores (Model 2), surface temperature mean and SD (Model 3), and two factors plus surface temperature (Model 4). P-values < 0.001 for population size in each model (not shown). Hosmer-Lemeshow (Hosmer et al.1991) chi-square statistics showed that goodness-of-fit for each model was adequate (p-values $\geq$ 0.46).

In Models 1-3, each social and biophysical neighborhood indicator was associated with the odds of one or more deaths from heat exposure among census block residents (p-values< 0.05) (Table 3). In Model 3, a 1° C increase in mean surface temperature was associated with a 32% increase in the odds of a death from heat exposure, after adjusting for surface temperature SD and population size.

Using the BIC and Raftery's (1995) statistical criteria for goodness-of-fit model comparison, there is positive evidence that Model 2 (HVI factor scores) is preferred over Model 1 (HVI additive index score). There is very strong evidence that Model 2 is preferred over Model

3 (only surface temperature) and positive evidence that Model 4 (Socioeconomic Vulnerability and Elderly/Isolation factors and surface temperature) is preferred over Model 2.

The other four HVI models we tested used each factor score and the percent no AC/cooler variable one at a time (results not shown). Using Raftery's (1995) criteria for comparing BICs, there is strong to very strong support for preferring Models 1, 2, and 4 over models with only single factors or no AC/cooler; strong to very strong support for preferring Model 3 over models with only Elderly/Isolation, Unvegetated Area, or no AC/cooler; and positive support for preferring a model with the Socioeconomic Vulnerability factor over Model 3.

The model with the index of concentrated disadvantage had a poor goodness-of-fit statistic (Hosmer-Lemshow p-value =0.098, results not shown). We substituted regional surface temperatures from Landsat scenes on two other days into Models 3 and 4 (See Supplemental Material, Table S1). Estimates based on regional temperatures on June 6, 2000 were nearly identical to those for July 24, 2000 but estimates based on data for October 25, 1990 were attenuated, consistent with smaller differences in vegetation among neighborhoods due to lower heat stress in autumn compared with the summer months. (Almost no heat exposure heat deaths occur in October.)

We observed a moderate degree of overall spatial clustering of vulnerability scores among neighborhoods (Global Moran's I for HVI =0.37; p-value=0.05). LISA analysis (Figure 2), identified neighborhood clusters with similar or dissimilar scores (p-value<= 0.05). High/high (red) clusters in Figure 2 represent neighborhoods with extremely high vulnerability scores next to others with extremely high scores. One-quarter of heat decedents lived in high/high clusters. Only 9.6% of heat decedents lived in clusters with low/low

vulnerability scores (blue) or dissimilar scores (e.g., purple areas indicating low vulnerability neighborhoods next to high). Sixty percent of the 50 heat deaths among homeless people occurred in high/high vulnerability clusters (see Supplemental Material, Figure S1).

## **Discussion**

We evaluated socio-environmental influences on heat-related deaths in this study. Mapping and statistical analyses were used to identify ‘hot spots’ of vulnerable places where socioeconomic disadvantages, unvegetated landscapes, and high temperatures were co-located with residences of heat decedents. Our analysis extended methods used in a small number of previous studies that identified neighborhoods at risk for heat-related deaths.

Patterns in Maricopa County were generally consistent with Reid and colleagues’ (2009) national analysis of a factor structure for a heat vulnerability index. Two exceptions were the configurations of factor loadings for AC and vegetation variables in our study that may be particular to desert cities. Only 13% of single-family homes in the Phoenix area lacked central AC or an evaporative cooler, which is much lower than a cross-section of 14 other U.S. cities (Jannsen et al. 2002). The desert southwest has an exceptionally hot climate and only the poorest households in Phoenix go without AC or a cooler. In our study, AC had a strong negative correlation with neighborhood poverty, low educational attainment, proportions of ethnic minorities and Latino immigrants (Table 1). All these variables loaded on the same factor, suggesting that AC in single-family homes is related to Socioeconomic Vulnerability to heat in Maricopa County (Table 2).

Percent Unvegetated Area was a separate factor in our analysis and had a weak but significant positive association with the odds of at least one heat death in a census block.

These findings indicate that vegetation may be a protective factor against human heat stress in all neighborhoods in desert cities. Our study confirms the importance of understanding regional variations in heat vulnerability indicators.

The preferred model for predicting residents' odds of dying from heat exposure in Maricopa County used Socioeconomic Vulnerability and Elderly/Isolation factors and mean surface temperature adjusted for population size and surface temperature SD in residential neighborhoods. The HVI model was preferred over a model using an index of neighborhood concentrated disadvantage, which included two social variables that are specifically related to heat risk and four variables that are not. Measurement errors in AC, land cover, and temperature may have attenuated the relationships between some vulnerability indicators and heat deaths. Given strong interest in adapting local built environments to climate change, it is important to continue developing more refined measures of environmental variables.

We used decedents' residential addresses as the indicator of vulnerable places because: (1) they represent the social and ecological contexts of many daily activity spaces, (2) they are consistent with other studies of neighborhood effects on social and health outcomes, and (3) the place of injury address on death certificates is not consistently coded for heat deaths. Mapping heat deaths onto the HVI derived for our study population showed that one-quarter of decedents lived in a relatively small number of highly vulnerable neighborhood clusters that were mainly in central city areas (red areas in Figure 2). Living conditions could have contributed to these deaths but people might have died in a different place, such as work or other activity sites. More research is needed on the circumstances of heat deaths.

We can say, however, that a majority of heat deaths among the homeless were reported in high vulnerability areas. Homeless people live in areas near downtown Phoenix and along industrial and transportation corridors that extend east and northwest from the central city where shelters, services, and hospitals are located (Sanchez 2011).

Heat mortality research in urban neighborhoods should be replicated in other areas with different urban forms and climates. More research is also needed to understand processes underlying the spatial distribution of heat-vulnerable places and their relationships to deaths, especially deaths from other causes that may be related to hot weather, such as cardiovascular disease (McMichael et al. 2006). We did not include a neighborhood indicator of general population health or specific disease incidence in our analysis but if such an indicator were available, it might be an important vulnerability marker. In addition, the deaths we analyzed were identified by the county's heat death surveillance system, which could introduce unknown sources of variation into how causes of deaths were identified. Weather and other circumstances beyond the scope of this study could affect the temporal and spatial distributions of deaths over nine years.

This type of research provides a foundation of knowledge for local interventions. We compared several ways that models could be constructed using census data and remotely sensed data available for the U.S. Surface temperature may be a reasonable substitute for indices that rely on statistical computations with many variables if the intended purpose of the model is to prevent heat deaths. Although Model 4, which included several population characteristics and temperature, was preferred for predicting the odds of heat deaths, Model 3, which only included temperature variables, also had an adequate goodness-of-fit statistic. Temperature difference is a simple way to communicate relative risks to residents of

vulnerable neighborhoods. Satellite data relevant to neighborhood-scale investigations has excellent spatial coverage but it is collected discontinuously. Augmentation with a network of weather stations to measure air temperatures could also be useful.

There are many ways in which health interventions could be targeted to heat-vulnerable neighborhoods: extreme weather warning systems, preventive actions for the homeless, improved emergency response, or heat island mitigation. Although AC has been widely associated with lower morbidity and mortality rates from heat-related illnesses, it is not a panacea for adapting to rising urban temperatures and heat waves. Vulnerable people and places do not have equal access to AC in Maricopa County, in other cities (Janssen et al. 2002; O'Neill et al. 2005), or in developing countries with climates hotter than Phoenix (Sivak 2009). Looking toward the mid-to-late 21<sup>st</sup> century, the southwestern U.S. is expected to experience continuing drought that may dramatically reduce the amount of water available to irrigate urban vegetation for cooling outdoor temperatures and to produce energy for powering AC (Gober 2010). Over 90% of Arizona's energy is supplied by coal, natural gas, and nuclear power plants that use large amounts of water in the production and delivery of electricity (Pasqualetti and Kelley 2007). The complete array of responses needed to reduce heat risks for people in vulnerable places is complex and public health officials should cooperate with other stakeholders to coordinate a broad range of policies.

## **Conclusions**

We estimated neighborhood effects of population characteristics and features of the built and natural environments on deaths due to heat exposure in Maricopa County, Arizona (2000-2008). Spatial patterns showed substantial variability among neighborhoods in vulnerability to heat, odds of heat deaths occurring, and locations of vulnerable

neighborhood clusters. Many inner-city neighborhoods had higher vulnerability scores and more deaths, whereas higher neighborhood income and education, younger white populations, greener landscapes, AC, and cooler microclimates in suburban neighborhoods were associated with reduced heat vulnerability and fewer deaths. Heat deaths of homeless persons were reported primarily in the inner-city. Many decedents, however, lived in neighborhoods with lower vulnerability scores and, therefore, place-based indicators of vulnerability are complements and not substitutes for person-level risk variables. Surface temperature might be used as a single indicator in Maricopa County to identify the most heat-vulnerable neighborhoods. However, more attention to the socio-ecological complexities of climate mitigation and adaptation is a high public health priority. There are major local challenges ahead in preventing heat deaths under global regimes of climate change and urbanization.

## References

- Anderson GB, Bell ML. 2011. Heat waves in the United States: mortality risk during heat waves and effect modification by heat wave characteristics in 43 U.S. communities. *Environ Health Perspect* 119:210-218.
- Anselin, L 1995. Local indicators of spatial association—LISA. *Geogr Anal*, 27:93-115.
- Balbus JM, Malina C. 2009. Identifying vulnerable subpopulations for climate change health effects in the United States. *J Occup and Environ Med* 51:33-37.
- Brooks-Gunn J, Duncan GJ, Klebanov PK, Sealand N. 1993. Do neighborhoods influence child and adolescent development? *AJS* 99:353-395.
- Buyantuyev A, Wu J. 2010. Urban heat islands and landscape heterogeneity: linking spatiotemporal variations in surface temperatures to land-cover and socioeconomic patterns. *Landsc Ecol* 25:17-33.
- Centers for Disease Control (CDC). 2006. Heat-related deaths – United States, 1999-2003. *MMWR Morb Mortal Wkly Rep* 55:796-798.
- Cromley EK, McLafferty SL. 2002. *GIS and Public Health*. New York: Guilford Press.
- Cutter SL, Finch C. 2008. Temporal and spatial changes in social vulnerability to natural hazards. *Proc Natl Acad Sci USA* 105:2301-2306.
- Debbink MP, Bader, MDM. 2011. Racial segregation and low birth weight in Michigan's metropolitan areas. *Am J Public Health* 101:1714-1720.
- de Smith MJ, Goodchild MF, Longley PA. 2006. *Geospatial Analysis*. Troubador Publishing Ltd.
- Diez-Roux AV. 2001. Investigating neighborhood and area effects on health. *Am J Public Health* 91:1783-1789.
- Gober P. 2010. Desert urbanization and the challenges of water sustainability. *Current Opinion in Environmental Sustainability* 2:1-7.
- Grineski S, Bolin B, Boone C. 2007. Criteria air pollution and marginalized populations: environmental inequity in metropolitan Phoenix, Arizona. *Soc Sci Q* 88:535-554.
- Harlan SL, Brazel A, Prashad L, Stefanov WL, Larsen L. 2006. Neighborhood microclimates and vulnerability to heat stress. *Soc Sci Med* 63:2847-2863.
- Heisler GM, Brazel AJ. 2010. The urban physical environment: temperature and urban heat islands. In: *Urban Ecosystem Ecology* (Aitkenhead-Peterson J, Volder A, eds.). American

- Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Agronomy Monograph 55, 29-55.
- Hosmer DW, Taber S, Lemeshow S. 1991. The importance of assessing the fit of logistic regression models: a case study. *Am J Public Health* 81:1630-1635.
- Huang G, Zhou W, Cadenasso ML. 2011. Is everyone hot in the city? Spatial pattern of land surface temperatures, land cover and neighborhood socioeconomic characteristics in Baltimore, MD. *J Environ Manage* 92:1753-1759.
- Intergovernmental Panel on Climate Change (IPCC). 2007. Synthesis Report: Contribution of Working Groups I, II, and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.
- Janssen NAH, Schwartz J, Zanobetti A, Suh HH. 2002. Air conditioning and source-specific particles as modifiers of the effect of PM10 on hospital admissions for Heart and Lung Disease. *Environ Health Perspect* 110:43-49.
- Jenerette GD, Harlan SL, Brazel A, Jones N, Larsen L, and Stefanov WL. 2007. Regional relationships between vegetation, surface temperature, and human settlement in a rapidly urbanizing ecosystem. *Landsc Ecol* 22:353-365.
- Jenerette GD, Harlan SL, Stefanov WL, and Martin CA. 2011. Ecosystem services and urban heat riskcape moderation: water, green spaces, and social inequality in Phoenix, USA. *Ecol Appl* 21:2637-2651.
- Johnson DP, Stanforth A, Lulla V, Luber G. 2012. Developing an applied extreme heat vulnerability index utilizing socioeconomic and environmental data. *Appl Geogr* 35: 23-31.
- Johnson DP, Wilson JS. 2009. The socio-spatial dynamics of extreme urban heat events: the case of heat-related deaths in Philadelphia. *Appl Geogr* 29: 419-434.
- Johnson DP, Wilson JS, Luber GC. 2009. Socioeconomic indicators of heat-related health risk supplemented with remotely sensed data. *Int J Health Geogr* 8:57-70. doi:10.1186/1476-072X-8-57
- Kalkstein AJ, Kalkstein LS. 2004. The development of an evaporative cooler warning system for Phoenix, Arizona. Presented to the NOAA/NWS Office, Phoenix, Arizona. Available: [http://www.udel.edu/SynClim/Evap\\_cooler.pdf](http://www.udel.edu/SynClim/Evap_cooler.pdf). [accessed 11 October 2012].
- Kilbourne EM, Choi K, Jones S, Thacker SB. 1982. Risk factors for heatstroke: a case control study. *JAMA* 247:3332-3336.

- Klinenberg E. 2002. *Heat Wave: A Social Autopsy of Disaster in Chicago*. Chicago: University of Chicago Press.
- Kovats RS, Koppe C. 2005. Heat waves: past and future impacts on health. In: *Integration of Public Health with Adaptation to Climate Change* (Ebi KL, Smith JB, Burton I, eds.). New York: Taylor and Francis.
- Lochner KA, Kawachi I, Brennan RT, Buka SL. 2002. Social capital and neighborhood mortality rates in Chicago. *Soc Sci Med* 56:1797-1805.
- Maricopa County Department of Public Health. 2011 (December). Heat-Associated Deaths in Maricopa County, AZ: Report for 2010. Available: <http://www.maricopa.gov/publichealth/Services/EPI/pdf/heat/2010annualreport.pdf> [accessed 16 February 2010].
- Mujahid MS, Diez Roux AV, Shen, M, Gowda, D, Sanchez B, Shea S et al. 2008. Relation between neighborhood environments and obesity in multi-ethnic study of atherosclerosis. *Am J Public Health* 167:1349-1357.
- McMichael AJ, Wilkinson P, Kovats RD, Pattenden S, Hajat S, Armstrong B, et al. 2008. International study of temperature, heat and urban mortality: the 'ISOTHURM' project. *Int JEpidemiol* 37:1121-1131.
- McMichael AJ, Woodruff RE, Hales S. 2006. Climate change and human health: present and future risks. *Lancet* 367:859-869.
- O'Neill MS, Zanobetti A, Schwartz J. 2005. Disparities by race in heat-related mortality in four U.S. cities: the role of air conditioning prevalence. *J Urban Health* 82:191-197.
- Pasqualetti M, Kelley S. 2007. The water costs of electricity in Arizona. Arizona Water Institute. Available: <http://www.circleofblue.org/waternews/wp-content/uploads/2010/08/AZ-Solar-Water-Fact-Sheet.pdf>. [accessed 12 October 2012].
- Pickett KE, Pearl M. 2001. Multilevel analyses of neighborhood socioeconomic context and health outcomes: a critical review. *J Epidemiol Community Health* 55: 111-122.
- Raftery AE. 1995. Bayesian model selection in social research. 1995. *Sociol Methodol* 25:111-163.
- Reid CE, Mann JK, Alfasso R, English PB, King GC, Lincoln RA, Margolis HG, Rubado DJ, Sabato JE, West NL, Woods B, Navarro KM, Balmes JR. 2012. Evaluation of a heat

- vulnerability index on abnormally hot days: an environmental public health tracking study. *Environ Health Perspect* 120:715-720.
- Reid CE, O'Neill MS, Gronlund CJ, Brines SJ, Brown DG, Diez-Roux AV, Schwartz J. 2009. Mapping community determinants of heat vulnerability. *Environ Health Perspect* 117:1730-1736.
- Ruddell DM, Harlan SL, Grossman-Clarke S, Buyanteyev A. 2010. Risk and exposure to extreme heat in microclimates of Phoenix, AZ. In: *Geospatial Contributions to Urban Hazard and Disaster Analysis*. (Showalter P, Lu Y, eds). Springer-Verlag.
- Sanchez C. 2011. Tricks of the shade: heat related coping strategies of urban homeless persons in Phoenix, Arizona. Master's Thesis, Arizona State University.
- Sampson RJ. 2009. Racial stratification and the durable tangle of neighborhood inequality. *Ann Am Acad Pol and Soc Sci* 621:260-280.
- Sampson RJ, Moronoff, JD, Gannon-Rowley T. 2002. Assessing 'neighborhood effects': social processes and new directions in research. *Annu Rev Sociol* 28:443-478.
- Sivak M. 2009. Potential energy demand for cooling in the 50 largest metropolitan areas of the world: implications for developing countries. *Energy Policy* 37:1382-1384.
- Smoyer KE. 1998. Putting risk in its place: methodological considerations for investigating extreme event health risk. *Soc Sci Med* 47:1809-1824.
- Tucker CJ. 1979. Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sens Environ* 8:127-150.
- United Nations Population Fund (UNFPA). State of the World Population 2007: Unleashing the Potential of Urban Growth. Available: <http://www.unfpa.org/swp/2007/english/introduction.html> [accessed 8 September 2011].
- WHO (World Health Organization). 2007. International Classification of Diseases. 10th Revision. Available: <http://apps.who.int/classifications/icd10/browse/2010/en> [accessed 5 November 2012].
- Wilson WJ. 1987. *The Truly Disadvantaged: The Inner City, the Underclass, and Public Policy*. Chicago: University of Chicago Press.
- Yip F, Flanders W, Wolkin A, Engelthaler D, Humble W, Antonio N, Lewis, L et al. 2008. The impact of excess heat events in Maricopa County, Arizona: 2000-2005. *Int J Biometeorol* 52:765-772.

Zhang K, Oswald EM, Brown DG, Brines SJ, Gronlund CJ, White-Newsome, JL, Rood, RB, O'Neill MS. 2011. Geospatial exploration of spatial variation of summertime temperatures in the Detroit metropolitan region. *Environ Res* 111:1046-1053.

Table 1. Means, standard deviations, and Pearson's correlations for variables in the 2000 U.S. Census Maricopa County block groups (n=2081)

| Variables in Analysis                      | Ethnic Minority | Latino Immigrant | Below Poverty | No H.S. Diploma | Age 65 or Older | Age 65 x Alone | Living Alone | No AC/Cooler | Unvegetated Area (mean) | Unvegetated Area (SD) | Land Surface Temp (mean) | Land Surface Temp (SD) |
|--|-----------------|------------------|---------------|-----------------|-----------------|----------------|--------------|--------------|-------------------------|-----------------------|--------------------------|------------------------|
| Mean                                       | 0.34            | 0.11             | 0.12          | 0.20            | 0.13            | 0.08           | 0.24         | 0.13         | -0.21                   | -0.11                 | 54.28                    | 2.14                   |
| Std. Dev.                                  | 0.26            | 0.12             | 0.13          | 0.19            | 0.17            | 0.09           | 0.14         | 0.22         | 0.07                    | 0.06                  | 1.94                     | 1.19                   |
| Ethnic Minority                            | 1.00            |                  |               |                 |                 |                |              |              |                         |                       |                          |                        |
| Latino Immigrant                           | 0.81**          | 1.00             |               |                 |                 |                |              |              |                         |                       |                          |                        |
| Below Poverty                              | 0.74**          | 0.71**           | 1.00          |                 |                 |                |              |              |                         |                       |                          |                        |
| No H.S. Diploma                            | 0.84**          | 0.80**           | 0.73**        | 1.00            |                 |                |              |              |                         |                       |                          |                        |
| Age 65 or Older                            | -0.40**         | -0.25**          | -0.18**       | -0.14**         | 1.00            |                |              |              |                         |                       |                          |                        |
| Age 65 x Alone                             | -0.26**         | -0.14**          | -0.03         | -0.02           | 0.88**          | 1.00           |              |              |                         |                       |                          |                        |
| Living Alone                               | -0.16**         | -0.09**          | 0.10**        | -0.60**         | 0.45**          | 0.63**         | 1.00         |              |                         |                       |                          |                        |
| No AC/Cooler                               | 0.67*           | 0.64**           | 0.57**        | 0.68**          | -0.16**         | -0.04          | -0.06        | 1.00         |                         |                       |                          |                        |
| Unvegetated Area (mean) 07/24/2000         | 0.16**          | 0.17**           | 0.22**        | 0.20**          | 0.00            | 0.02           | 0.04         | 0.15**       | 1.00                    |                       |                          |                        |
| Unvegetated Area (SD) 07/24/2000           | 0.10**          | 0.13**           | 0.11**        | 0.10**          | -0.14**         | -0.08**        | -0.06**      | 0.16**       | 0.69**                  | 1.00                  |                          |                        |
| Land Surface Temperature (mean) 07/24/2000 | 0.35**          | 0.32**           | 0.33**        | 0.37**          | -0.11**         | -0.06**        | -0.08**      | 0.32**       | 0.78**                  | 0.67**                | 1.00                     |                        |
| Land Surface Temperature (SD) 07/24/2000   | -0.08**         | -0.13**          | -0.07**       | -0.06**         | 0.10**          | 0.04           | 0.00         | -0.18*       | -0.47**                 | -0.86**               | -0.54**                  | 1.00                   |

\* p<=0.05; \*\*p<=0.01

Table 2. Principal Components Analysis of heat vulnerability variables in the 2000 U.S. Census Maricopa County block groups (n=2081)

| Variables in<br>Principal Components<br>Analysis | Factor Loadings <sup>a</sup>                 |                                 |                                  |
|--|--|---------------------------------|----------------------------------|
|  | Factor 1:<br>Socio-economic<br>Vulnerability | Factor 2:<br>Elderly/ Isolation | Factor 3:<br>Unvegetated<br>Area |
| Ethnic Minority                                  | 0.91   | -0.25                           | 0.04                             |
| Latino Immigrant                                 | 0.90   | -0.11                           | 0.06                             |
| Below Poverty Line                               | 0.86   | 0.04                            | 0.09                             |
| No H.S. Diploma                                  | 0.92   | 0.03                            | 0.06                             |
| No Central AC/Cooler                             | 0.79   | -0.03                           | 0.09                             |
| Age 65 or older                                  | -0.19  | 0.88                            | -0.04                            |
| Age 65 or older living alone                     | -0.03  | 0.96                            | -0.02                            |
| Living Alone                                     | 0.01   | 0.77                            | 0.01                             |
| Unvegetated Area (mean)                          | 0.14   | 0.06                            | 0.91                             |
| Unvegetated Area (SD)                            | 0.05   | -0.10                           | 0.92                             |

a. Factor extraction was performed using varimax rotation so that the factors are uncorrelated with each other. The numbers in the columns are factor loadings that represent correlations between the variables and factors and also the weights of each variable on the factors.

Table 3. Odds ratios (95% CI) from binary logistic regression models of at least one heat-associated death in the 2000 U.S. Census Maricopa County block group of residence, 2000-2008<sup>a,b</sup> (n=2,081)

| Variables in Models                        | Model 1<br>Heat Vulnerability Index<br>(HVI) Integer Scores |                   | Model 2<br>HVI Factor Scores<br>1-3 |              | Model 3<br>Land Surface Temperature<br>07/24/2000 |              | Model 4<br>HVI Factor Scores 1 and 2<br>and Land Surface<br>Temperature |              |
|--|---|-------------------|-------------------------------------|--------------|---|--------------|---|--------------|
|  | Odds ratio  | (CI) <sup>c</sup> | Odds ratio                          | (CI)         | Odds ratio  | (CI)         | Odds ratio  | (CI)         |
| HVI (integer scores)                       | 1.34***   | (1.23, 1.45)      |                                     |              |   |              |   |              |
| Socioeconomic Vulnerability (HVI Factor 1) |   |                   | 1.50***                             | (1.33, 1.70) |   |              | 1.34***   | (1.17, 1.53) |
| Elderly / Isolation (HVI Factor 2)         |   |                   | 1.38***                             | (1.22, 1.56) |   |              | 1.39***   | (1.23, 1.56) |
| Unvegetated Area (HVI Factor 3)            |   |                   | 1.19*                               | (1.02, 1.39) |   |              |   |              |
| Land Surface Temperature (mean)            |   |                   |                                     |              | 1.32***   | (1.20, 1.45) | 1.23***   | (1.11, 1.36) |
| Land Surface Temperature (SD)              |   |                   |                                     |              | 1.16*   | (1.01, 1.34) | 1.09  | (0.94, 1.27) |
| -2 log L                                   | 1406.64   |                   | 1385.18                             |              | 1412.40   |              | 1372.73   |              |
| Hosmer-Lemeshow p-value                    | 0.46  |                   | 0.99                                |              | 0.74  |              | 0.58  |              |
| BIC <sup>d</sup>                           | 1429.56   |                   | 1423.38                             |              | 1442.96   |              | 1418.57   |              |

a. Dependent variable: at least one decedent who died from heat exposure lived in the census block group (1=yes; 0=no).  
 b. Intercept and population size of census block groups in each model; p<0.001 (not shown).  
 c. CI: confidence interval.  
 d. BIC = -2logL+Np\*Ln(n) where Np=number of parameters and n = 2081.

\* p<=0.05; \*\* p<=0.01; \*\*\* p<=0.001

## Figure Legends


### Figure 1.

Heat Vulnerability Index scores (using a method modified from Reid et al. 2009) mapped for 2,081 census block groups in Maricopa County, AZ. Higher scores represent higher vulnerability. Number of heat-associated deaths (2000-2008) is denoted by size of green circles in block groups of decedents' residences. The map inset in the lower right corner outlines in red the urbanized area of Maricopa County that is shown in the larger map. The county, which also contains a much larger area of uninhabited desert and sparse settlement, is outlined in blue. The urbanized area in the larger map covers all the cities and all but one of the major towns in the county. Only four residences of people who died from heat exposure were located outside the urbanized area (see four green circles in map inset).

### Figure 2.


Univariate analysis of the Local Indicator of Spatial Association (LISA) identified clusters of census block groups in Maricopa County, AZ with similar or dissimilar Heat Vulnerability Index scores ( $p\text{-value} \leq .05$ ). High/high (red) areas in the map are clusters of neighboring census block groups with uniformly high vulnerability scores; low/low (blue) areas are clusters with low vulnerability scores; low/high (purple) areas represent a census block group with a low vulnerability score neighbored by high vulnerability block groups; high/low (pink) areas represent a census block group with a high vulnerability score neighbored by low vulnerability block groups. Entries in the legend also show the percentages of 2000-2008 heat-related decedents who were residents in each type of cluster. On the map, number of heat-associated deaths (2000-2008) is denoted by size of green circles in block groups of decedents' residences.

Figure 1



279x215mm (200 x 200 DPI)

Figure 2



279x215mm (200 x 200 DPI)