

ehp

**ENVIRONMENTAL
HEALTH
PERSPECTIVES**

ehponline.org

Monitoring Intracellular Redox Changes in Ozone-Exposed Airway Epithelial Cells

**Eugene A. Gibbs-Flournoy, Steven O. Simmons,
Philip A. Bromberg, Tobias P. Dick, and
James M. Samet**

<http://dx.doi.org/10.1289/ehp.1206039>

Online 18 December 2012



NIEHS

**National Institute of
Environmental Health Sciences**

**National Institutes of Health
U.S. Department of Health and Human Services**

Monitoring Intracellular Redox Changes in Ozone-Exposed Airway Epithelial Cells

Eugene A. Gibbs-Flournoy¹, Steven O. Simmons², Philip A. Bromberg³, Tobias P. Dick⁴, and James M. Samet⁵

¹Curriculum in Toxicology, University of North Carolina, Chapel Hill, North Carolina, USA;

²Integrated Systems Toxicology Division, National Health and Environmental Effects Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina, USA;

³Center for Environmental Medicine, Asthma and Lung Biology, University of North Carolina, Chapel Hill, North Carolina, USA; ⁴Division of Redox Regulation, DKFZ-ZMBH Alliance, German Cancer Research Center (DKFZ), Heidelberg, Germany;

⁵Environmental Public Health Division, NHEERL, U.S. EPA, Chapel Hill, North Carolina, USA.

Corresponding author contact information

James M. Samet, 104 Mason Farm Rd., EPA Human Studies Facility, Chapel Hill NC 27599-7310 USA. Telephone: (919) 966-0665. Fax: (919) 962-6271. E-mail: samet.james@epa.gov

EPA Disclaimer

The research described in this article has been reviewed by the National Health and Environmental Effects Research Laboratory, U.S. EPA, and approved for publication. The

contents of this article should not be construed to represent Agency policy nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

Keywords: glutathione, human airway epithelial cells, imaging, intracellular, oxidative stress, ozone, NADPH, redox, roGFP.

Short Title: Imaging Redox Changes Using O₃ as a Model Oxidant

Abbreviations:	2-AAPA	2-acetylamino-3-[4-(2-acetylamino-2-carboxyethylsulfanylthiocarbonylamino)phenylthiocarbamoylsulfanyl]propionic acid
	DTT	Dithiothreitol
	E _{GSH}	Glutathione Redox Potential
	GPx	Glutathione Peroxidases
	Grx1	Glutaredoxin 1
	GSH	Reduced Glutathione
	GSSG	Oxidized Glutathione
	LS+G	Locke Solution with Glucose
	LS-G	Locke Solution without Glucose
	PPP	Pentose Phosphate Pathway

Acknowledgements: The authors gratefully appreciate the encouragement and support of Dr. Wan-Yun Cheng, Mr. Robert Silbajoris, and Ms. Lisa Dailey.

CFI Statement: None of the authors have any actual or potential competing financial interests.

Abstract

Background: The toxicity of many xenobiotic compounds is believed to involve oxidative injury to cells. Direct assessment of mechanistic events involved in xenobiotic-induced oxidative stress is not easily achievable. Development of genetically-encoded probes designed for monitoring intracellular redox changes represents a methodological advance with potential applications in toxicological studies.

Objective: Test the utility of redox-sensitive GFP (roGFP)-based redox sensors for monitoring real-time intracellular redox changes induced by xenobiotics in toxicological studies.

Methods: roGFP2, a reporter of the glutathione redox potential (E_{GSH}), was used to monitor E_{GSH} in cultured human airway epithelial cells (BEAS 2B) undergoing exposure to 0.15 – 1.0 ppm O_3 . Cells were imaged in real-time using a custom-built ozone exposure system coupled to a confocal microscope.

Results: O_3 exposure induced a dose- and time- dependent increase of the cytosolic glutathione redox potential. Additional experiments confirmed that roGFP2 is not directly oxidized but properly equilibrates with the glutathione redox couple: Inhibition of endogenous Grx1 disrupted roGFP2 responses to O_3 and a Grx1-roGFP2 fusion protein responded more rapidly to O_3 exposure. Selenite-induced upregulation of GPx expression enhanced roGFP2 responsiveness to O_3 , suggesting that (hydro)peroxides are intermediates linking O_3 exposure to glutathione oxidation.

Conclusion: Exposure to O_3 induces a profound increase in the cytosolic E_{GSH} of airway epithelial cells that is indicative of an oxidant-dependent impairment of glutathione redox homeostasis. These studies demonstrate the utility of using genetically-encoded redox reporters

in making reliable assessments of cells undergoing exposure to xenobiotics with strong oxidizing properties.

Background

The intracellular redox environment is a highly dynamic setting governed by the formation and degradation of various reactive species of oxygen and nitrogen. Under normal physiological conditions, the cytosol, nucleus, and mitochondrial matrix space maintain homeostatic conditions in favor of a highly reducing environment (Cannon and Remington 2008). Intracellular reducing conditions are largely maintained by millimolar concentrations of reduced glutathione and its accessory enzymes that together comprise the glutathione system (Anderson 1998). Ultimately, maintenance of the intracellular glutathione redox potential (E_{GSH}) comes from the metabolism of glucose, as glutathione is reduced by glutathione reductase using NADPH produced by the pentose phosphate pathway (Wamelink et al. 2008).

A number of pathophysiological states are associated with changes in the intracellular glutathione redox potential (Dubinina and Dadali 2010; Ma 2010; Yang and Omaye 2009). Such “oxidative stress” is commonly cited as a mechanistic feature of the toxicity of numerous xenobiotic compounds linked to adverse health outcomes (Bargagli et al. 2009; Ciencewicki et al. 2008; Kohen and Nyska 2002). For instance, the health effects of the potent ambient air pollutant ozone (O_3) are understood to be mediated through an oxidative stress mechanism involving the oxidation of cellular biomolecules (Ballatori et al. 2009; Kelly et al. 1995; Mudway and Kelly 2000). In the lung, ozone exposure causes decrements in pulmonary function and induces inflammatory responses derived from the bronchial epithelium, a major target of ozone exposure (Ballinger et al. 2005; Ciencewicki et al. 2008; Kelly et al. 1995; Mudway and Kelly 2000; Pryor 1992; Pryor et al. 1995; Song et al. 2010). Due to its high reactivity, O_3 interacts with cellular and extracellular biomolecules resulting in multiple types of oxidative damage to lipids, proteins, and nucleic acids (Kelly et al. 1995; Laumbach 2010; Mudway and

Kelly 2000; Srebot et al. 2009; Van der Vliet et al. 1995; Yang and Omaye 2009). While numerous studies have established oxidant damage of biomolecules as a result of O₃ exposure, direct measures of ozone-mediated 'oxidative stress' have been difficult to achieve, yet alteration of a defined intracellular redox couple like glutathione would represent an important early indicator of the oxidative effects of O₃ exposure.

Recent methodological advances have been made it possible to focus studies of pro-oxidative changes to specific redox couples within defined subcellular compartments (Meyer and Dick 2010), potentially affording greater specificity in mechanistic investigations of the oxidative effects of xenobiotic exposures. A new generation of genetically-encoded fluorophores permits direct assessment of the oxidative effects of xenobiotic compounds in relation to the glutathione redox pair (GSSG/GSH) with unprecedented spatial and temporal resolution (Cheng et al. 2012; Dooley et al. 2004; Meyer and Dick 2010). roGFP2 acts as a reporter of intracellular glutathione redox potential by equilibrating with the GSH/GSSG redox pair (Gutscher et al. 2008; Meyer and Dick 2010; Morgan et al. 2011). In short, in a reaction that depends on catalysis by glutaredoxins, roGFP2 responds to oxidation of reduced glutathione (GSH) to its oxidized form, GSSG, via the internal formation of a disulfide bond (Gutscher et al. 2008; Meyer 2008; Meyer and Dick 2010) (Figure 1). The formation of the disulfide bond alters the spectral characteristics of the GFP fluorophore causing the intensity of the emitted green fluorescence (~520 nm) induced by excitation at 488 nm to decrease, while causing the emitted fluorescence after excitation at 405 nm to increase, thus making this sensor a ratiometric probe. Further efforts to improve the responsiveness of roGFP2 have led to the conjugation of pathway-specific enzymes to create a chimeric fusion of proteins operating as redox relays. In particular, the conjugation of

glutaredoxin 1 (Grx1) to roGFP2 has been shown to enhance the kinetics of the roGFP2 response to the oxidation of glutathione (Gutscher et al. 2008).

In the present study, we used live-cell microscopy to monitor the cytosolic E_{GSH} of airway epithelial cells undergoing exposure to ozone in real-time. Here, we report an approach to validate the use of roGFP2-based redox sensors in toxicological studies of xenobiotics with strong oxidizing properties.

Methods

Materials and Reagents

Tissue culture media and supplements were purchased from Lonza (Walkersville, MD). Wilco Wells glass-bottom culture dishes were obtained from Ted Pella (Redding, CA) and Warner Instruments (Hamden, CT). Fugene 6 transfection reagent was acquired from Roche Applied Science (Indianapolis, IN). Kits to measure intracellular glutathione and NADPH were bought from Promega (Madison, WI) and AbCam (Cambridge, MA), respectively. Laboratory reagents and chemicals including Hydrogen peroxide, Dithiothreitol (DTT), 2-acetylamino-3-[4-(2-acetylamino-2-carboxyethylsulfanylthiocarbonylamino)phenylthiocarbamoylsulfanyl]propionic acid (2-AAPA), Buthionine sulfoximine (BSO), and Sodium selenite were obtained from Sigma-Aldrich (St. Louis, MO). Basic laboratory supplies were purchased from Fisher Scientific (Raleigh, NC).

Cell Culture

Transformed human airway epithelial cells (BEAS-2B, subclone S6; (Reddel et al. 1988)) were cultured as previously described (Tal et al. 2010) and maintained in serum-free keratinocyte growth medium, KGM (Lonza, Walkersville, MD). The cells were incubated in a humidified incubator at 37°C in 5% CO₂. For most live-cell exposures, BEAS-2B cells were plated in 35mm Wilco Wells glass-bottom dishes with a 12 mm #1.5 glass aperture (Ted Pella, Redding, CA).

Genetically Encoded Redox Sensors

Plasmid for the redox-sensitive green fluorescent protein 2, roGFP2, was the generous gift of Dr. S. J. Remington (University of Oregon, Eugene, OR). Plasmid for the hydrogen peroxide sensor, HyPer, was purchased from Evrogen (Axxora, Farmingdale, NY). Cytosolic and mitochondrially targeted versions of roGFP2 and HyPer were placed in lentiviral vectors as described previously (Cheng et al. 2012).

Plasmid Transfection and Lentiviral Transduction

One to two days prior to exposure, BEAS-2B cells were transiently transfected with 1 – 2 ug of plasmid DNA encoding Grx1-roGFP2, HyPer, or roGFP2 using 3 – 5 ul of Fugene 6 transfection reagent for each 35mm culture dish. Stable expression of a specific genetically encoded fluorescent reporter was performed via lentiviral transduction. In short, a lentivirus encoding roGFP2 or HyPer (specifically targeted to the cytosol or mitochondria) was incubated for 4 hrs (37°C / 5% CO₂) with wild-type BEAS-2B cells using a multiplicity of infection (MOI) of 5 – 10 in a single well of a 6-well dish. The viral particles were then removed and fresh KGM was placed on the cells, and the BEAS-2B cells were then allowed to grow to confluency. Upon confluency, the cells were expanded to T75 dishes where they were propagated for multiple

passages. For some experiments, stably transduced cells were sorted for optimal fluorescence expression at the UNC Core Flow Cytometry Facility.

Exposure Conditions

Newly transfected or stably transduced BEAS-2B cells expressing the fluorescent reporter of interest were cultured as described above. Prior to exposure, subconfluent cells were equilibrated in Locke solution (Taylor-Clark and Udem 2010) for 2 hrs. at 37°C in 5% CO₂. For the studies herein, two versions of Locke Solution were derived by adding or excluding 1 mg/ml dl-glucose. Irrespective of the type of Locke solution used to equilibrate cells, all live-cell O₃ exposures were performed using 0.5 ml of Locke solution without glucose. For all imaging experiments, cells were exposed in a custom-built stage-top exposure system maintained at 37°C with 1.5 L/min of 5% CO₂/balance air at a relative humidity \geq 95% (see Supplemental Material, Figure S1). In some experiments, cells were pretreated with 100 μ M 2-AAPA, a dithiocarbamate inhibitor of glutaredoxins, during the 2 hr buffer equilibration period. Similarly, pretreatment of cells with 1 μ M sodium selenite for 24 – 48 hrs prior to O₃ exposure was carried out to induce the overexpression of glutathione peroxidases (GPx).

For each experiment, cells were exposed to control air (5% CO₂/Balance air) or ozone concentrations ranging from 0.15 – 1.0 ppm. The entire exposure period typically consisted of 3 component intervals collectively lasting up to an hour in length. They included: 1) an initial untreated baseline period of 5 min; 2) an exposure period of up to 45 min; and 3) a 10 min control exposure period in which cells were oxidized by 0.1 to 1.0 mM hydrogen peroxide for 5 min and then reduced by 10 mM DTT for an additional 5 min. During these exposures, the ozone concentration in the exposure chamber was monitored in real-time using a Dasibi Model 1003-

AH Ozone Analyzer sampling at a flow of 2.0 L/min. Ozone exposures for non-imaging assays were performed using exposure chambers operated by the U.S. Environmental Protection Agency's (EPA) Environmental Public Health Division.

Imaging Analysis

All live-cell experiments done in real-time were conducted using a Nikon Eclipse C1si spectral confocal imaging system equipped with an Eclipse Ti microscope, Perfect Focus System, and 404 nm, 488 nm, 561 nm, and 633 nm primary laser lines (Nikon Instruments Corporation, Melville, NY). Images were acquired using a 60X Plan Apo lens. For experiments involving the genetically encoded fluorescent reporters, roGFP2 and HyPer, green fluorescence was observed via the use of independent excitations at 404 and 488 nm while emitted light was collected for each using a 525/30 nm band-pass filter (Chroma, Bellows Falls, VT). Results were calculated as ratios of the emissions excited by the 488 nm and 404 nm lasers scanned sequentially at a frequency of 1 min. All imaging data were acquired using the Nikon EZ-C1 software.

Measurement of Intracellular NADPH

After cells were equilibrated in fresh KGM or Locke solution (+/- glucose) for 2 hrs, the intracellular levels of total NADPH were assessed using an AbCam NADP/NADPH assay kit. The manufacturer's instructions provided with the kit were used to carry out this assay. Following the equilibration period, cells were immediately placed on ice and washed with cold 1X PBS just prior to the initial lysis step. Absorbance was read at 450 nm using a PolarStar Optima microplate reader (BMG Labtech, Durham, NC).

Statistical Analysis

All imaging data were quantified using NIS-Elements AR software (Nikon, Melville, NY). For each experiment, the responses of 5–10 cells were collected as regions of interest and then averaged to derive an overall response. Data are expressed as the mean of at least three repeated experiments. Pairwise comparisons of control and treatment groups were performed using ANOVA and linear regression with $p < 0.05$ taken as being statistically significant.

Results

Ozone exposure induces an increase in the cytosolic glutathione redox potential

The presence of glucose in the exposure media is known to shorten the half-life of O_3 (Taylor-Clark and Udem 2010). Therefore, in these experiments, cells were first equilibrated in Locke solution containing glucose (LS+G) for 2 hrs prior, and then switched to Locke solution without glucose (LS-G) for the exposure. Exposure of BEAS-2B cells expressing cytosolic roGFP2 to 0.15 - 0.50 ppm O_3 resulted in a dose- and time- dependent probe response, reflecting an increase in the cytosolic glutathione redox potential (Figure 2; see Supplemental Material, Figure S5). Increasing O_3 concentration hastened the onset while elevating the magnitude of the oxidative response reported by roGFP2 (Figure 2). Addition of 0.1 mM H_2O_2 at the end of each O_3 exposure produced a maximal response which was fully reversible with the addition of 10 mM DTT.

Glucose deprivation potentiates the elevation of E_{GSH} induced by ozone exposure

On an individual basis, BEAS 2B cells equilibrated in the presence of glucose (LS+G) were observed to respond variably to a given concentration of O_3 , suggesting substantial heterogeneity of redox homeostasis within the cellular population under the given conditions. Figure 3A shows the individual responses of seven BEAS-2B cells in the same field of view being exposed to 0.5 ppm O_3 , with some cells responding with strongly increasing E_{GSH} , while others responded only minimally. Furthermore, some cells exhibited an intermediate response followed by a recovery of E_{GSH} despite continued O_3 exposure. Given the importance of pentose phosphate pathway (PPP)-generated NADPH in maintaining intracellular reduced glutathione levels, we hypothesized that glucose deprivation would sensitize the cells to a subsequent O_3 exposure. As shown in Figure 3B, depriving the cells of glucose for 2 hr prior to exposure homogenized the magnitude, time of onset, and rate of response of the cells to O_3 . Glucose status did not affect the probe response to addition of H_2O_2 and DTT (Figures 3 A and B). As expected, glucose deprivation led to decreased cellular NADPH levels (see Supplemental Material, Figure S2).

Validation of glutathione-dependent roGFP2 responses to ozone exposure

Given the extreme reactivity of O_3 with biomolecules, we considered the possibility that the spectral changes of the probe interpreted as changes in E_{GSH} are the result of a direct oxidation of roGFP2 by O_3 itself or by an O_3 -generated secondary oxidant. We therefore undertook a series of experiments to determine whether O_3 -induced changes in the roGFP2 fluorescence intensity ratio involve components of the glutathione system through which roGFP2 has been demonstrated to respond (Figure 1). We first aimed to confirm that O_3 exposure leads to increased levels of GSSG. To this end we assessed the extent of intracellular glutathione oxidation in control and

ozone treated cells. We observed up to 3-fold increases in GSSG following 1 ppm O₃ exposure in LS-G as compared to air controls (see Supplemental Material, Figure S3) which agrees with previous reports of the effect of O₃ exposure on intracellular glutathione (Chalfant and Bernd 2011; Todokoro et al. 2004).

Next, we asked whether the roGFP2 response to O₃ is influenced by glutaredoxin (Grx) activity. Grx is essential to mediate roGFP2 oxidation by GSSG (Figure 1), but should play no role if roGFP2 is directly oxidized by O₃. On the one hand, we compared the response of roGFP2 (which interacts with endogenous Grx) with that of Grx1-roGFP2, a translational fusion of glutaredoxin-1 and roGFP2. The fusion of these components is known to kinetically improve the equilibration between roGFP2 and GSSG in a highly specific manner (Gutscher et al. 2008; Meyer and Dick 2010). We found that the O₃-induced increase of E_{GSH} in BEAS cells expressing Grx1-roGFP2 occurred earlier and at a faster rate relative to that reported by cells expressing unlinked roGFP2, thus indicating a GSSG/Grx-specific response (Figure 4A). In contrast, we investigated the effect of glutaredoxin inhibition by pretreating cells with the dithiocarbamate derivative 2-AAPA (Sadhu et al. 2012). The inhibitor completely blocked roGFP2 responses to ozone and H₂O₂ (which also act indirectly through glutathione oxidation and reduction, respectively) (Figure 4B), thus confirming the role of glutaredoxin as the catalyst necessary for roGFP2 responsiveness to O₃. Together these observations confirm that O₃ is not simply sensed by the probe through direct oxidation, but rather by its specific effects on the glutathione redox couple.

Having confirmed that O₃ induces formation of GSSG which is then detected by the roGFP2 probe, we asked whether glutathione peroxidases (GPx), major generators of GSSG, are involved in the ozone response. Here we investigated the role of glutathione peroxidase activity in O₃-

induced roGFP2 redox changes by pretreating BEAS cells with 1 μ M sodium selenite for up to 48 hrs prior to O₃ exposure. Previous studies have used selenium supplementation as an effective means of increasing glutathione peroxidase expression, a finding that we also observed in preliminary studies with BEAS cells (see Supplemental Material, Figure S4) (Helmy et al. 2000; Holben and Smith 1999; Leist et al. 1996). Se-induced overexpression of GPx was found to accelerate roGFP2 oxidation during a 0.5 ppm O₃ exposure (Figure 4C), thus suggesting that O₃ gives rise to peroxides, which are then converted by GPx to GSSG, which is in turn reported by roGFP2 through the intervention of Grx.

Investigating the role of secondary products in ozone-induced redox changes

Since the data shown in Figure 4C suggested the involvement of peroxides in the O₃ response, we asked whether there is a specific role for hydrogen peroxide. For these experiments we examined generation of H₂O₂ as a consequence of O₃ exposure using the cytosolic-targeted H₂O₂-sensor, HyPer. O₃ caused a relatively modest increase in the HyPer response during the exposure period (Figure 5). However, the observed HyPer response did not precede nor match the magnitude of the roGFP2 response, making it unlikely that the observed increase in E_{GSH} is primarily caused by H₂O₂ generation.

Mitochondrial oxidant production, frequently associated with increased oxidation of mitochondrial glutathione, has been implicated as a contributing factor in the cellular response to xenobiotics (Cheng et al. 2010; Cheng et al. 2012; Hanson et al. 2004). Therefore, we next used mitochondrially-targeted roGFP2 (roGFP2-m) to assess the impact of ozone exposure on the mitochondrial glutathione redox potential of BEAS-2B cells. As shown in Figure 6, exposure to 1 ppm ozone, twice the amount used for cytosolic assessments, induced an increase in

mitochondrial E_{GSH} . However, relative to the cytosolic roGFP2 response, the increase in mitochondrial E_{GSH} occurred at a slower rate and achieved a lower magnitude. This suggests that mitochondrial oxidants are not the primary source of the oxidants that lead to increased cytosolic E_{GSH} .

Discussion

“Oxidative stress” is a frequently cited mechanistic component of the adverse health effects induced by numerous xenobiotic compounds (Bargagli et al. 2009; Chung and Marwick 2010; Ciencewicki et al. 2008; Jones 2008; Kohen and Nyska 2002; MacNee 2001; Ward 2010; Yang and Omaye 2009). However, the term “oxidative stress” is a very broad concept and the detection of early and specific indices of oxidant stress has proven to be methodologically difficult. The advent of genetically-encoded fluorescent reporters that are sensitive to their redox environment has enabled real-time imaging-based assessments of oxidant outcomes in living cells with unprecedented spatial and temporal resolution. In this study, we validated the use of one such reporter, roGFP2, for the specific assessment of xenobiotic-induced changes in the glutathione redox potential using ozone as a model toxicant and BEAS-2B cells as a model of the human bronchial epithelium.

The pro-oxidative change in E_{GSH} observed in this study represents an early event in the oxidant injury caused by O_3 . Ozone is a potent oxidant gas that has the potential to interact directly with virtually any cellular component, potentially including fluorescent reporter molecules such as roGFP2. Thus, in interpreting the probe response observed in O_3 -exposed BEAS-2B cells, we

had to consider the possibility that O₃ could be bypassing the glutathione system through which roGFP2 sensors normally respond (Gutscher et al. 2008; Meyer and Dick 2010).

Our findings strongly suggest that even in the presence of a strong oxidant like O₃, roGFP2 is oxidized only indirectly through its known coupling to the glutathione system. This conclusion is supported by several observations: First, glucose deprivation increased O₃-mediated roGFP2 oxidation, consistent with the requirement for NADPH in robustly maintaining E_{GSH}, the lack of glucose preventing regeneration of reducing equivalents throughout the exposure period. NADPH levels were approximately 70% lower in cells equilibrated in the absence of glucose, which appears to be sufficient to sensitize cells uniformly. In addition, it is important to bear in mind that there are other cellular processes that draw on the NADPH pool, and the continued lack of glucose largely prevents active regeneration of reducing equivalents throughout the exposure period.

Second, we confirmed the role of glutaredoxin in mediating the roGFP2 response to O₃. Grx1 is required to transfer oxidative equivalents from the glutathione pool to roGFP2. In previous studies using Grx1-roGFP2, the chimeric linkage of glutaredoxin-1 to roGFP2 enhanced responses to physiological oxidants such as H₂O₂ (Gutscher et al. 2008; Meyer and Dick 2010). Importantly, Grx1-roGFP2 also accelerated the roGFP2 response to O₃, where as inhibition of endogenous Grx with 2-AAPA prevented roGFP2 oxidation in the presence of O₃. While early reports describe 2-AAPA as being an inhibitor of glutathione reductase, a more recent study reports that this dithiocarbamate derivative acts as a direct inhibitor of glutaredoxins as well (Sadhu et al. 2012). Thus, the finding that 2-AAPA-treated cells failed to respond to O₃ supports the involvement of Grx and is consistent with the claim that 2-AAPA is an inhibitor of glutaredoxins. Importantly, the fact that use of this inhibitor was effective at disconnecting the

glutathione pool from the redox reporter argues against a non-specific interaction between O₃, or a secondary oxidant, and the roGFP2 sensor.

Lastly, findings from experiments examining the effect of overexpression of glutathione peroxidases (GPx) using prolonged selenite supplementations are consistent with an upstream role of GPx in the oxidative pathway leading to O₃-induced roGFP2 oxidation. GPx couple the reduction of (hydro)peroxides to the generation of GSSG (Arthur 2000; Mates and Sanchez-Jimenez 1999; Meyer and Dick 2010). In our system, the increased expression of GPx enhanced roGFP2 oxidation, suggesting that O₃ exposure generates (hydro)peroxides, which then drive the formation of GSSG. In fact, ozone has been shown to produce many types of lipid hydroperoxides upon exposure (Mudway and Kelly 2000; Yang and Omaye 2009). Moreover, cells exposed to O₃ may have an increased H₂O₂ burden as well. Overall, the results from GPx overexpressing cells suggest that the increased activity of these enzymes leads to an enhanced catalytic destruction of peroxides with concomitant GSH oxidation, leading to the roGFP2 response to ozone.

Using the H₂O₂ probe HyPer, our initial assessments suggested slightly elevated H₂O₂ production following ozone exposure of BEAS cells. The quantitative interpretation of HyPer responses is however difficult. It is not clear to what extent the OxyR domain of HyPer may be outcompeted by endogenous peroxidases. In addition, HyPer is highly pH sensitive, as much as the cpYFP module on which it is based (Schwarzlander et al. 2012). Thus, an O₃-induced intracellular acidification could dampen the HyPer response to H₂O₂. Nevertheless, following O₃ exposure, HyPer responded to exogenously applied H₂O₂ and DTT as expected, which demonstrates the general functionality of the probe. Taken together, the delayed time of onset, slow rates of response, and the relatively low magnitude of the HyPer responses, may suggest

that H_2O_2 production is not a major factor in the total O_3 -induced E_{GSH} effects. Likewise, the measurements in mitochondrially-targeted roGFP do not support a mitochondrial source for the O_3 -induced increase in cytosolic E_{GSH} . These results suggest that the relevant oxidant species, potentially a hydroperoxide, is primarily generated in the cytosol, or within the outer membranes, and the mitochondrial E_{GSH} response would be expected to lag behind that of the cytosol. Additionally, differences in peroxidase composition and activity may contribute to the lag in the O_3 -induced mitochondrial E_{GSH} response relative to the cytosol.

The studies presented herein cannot completely exclude a partial contribution of direct interactions between O_3 , or its secondary byproducts, with the thiols of the roGFP2 sensor. In addition, because the roGFP2 fluorescence ratio reflects E_{GSH} , which is a function of both the GSSG:GSH ratio and the total glutathione concentration, it is possible that O_3 -induced electrophilic attack mediates the changes reported by roGFP2 by consuming reduced GSH. It should, however, be noted that the E_{GSH} in the cytosol or mitochondrial matrix is much more sensitive to an increase in GSSG than to depletion of GSH. This is because the E_{GSH} in the cytosol (around -320 mV, or even lower, in mitochondria) represents nanomolar GSSG in a millimolar pool of GSH. To deflect the roGFP2 signal from -320 mV to about -260 mV, only requires the concentration of GSSG to increase from 200 nM to 20 μM (in a 10 mM total glutathione pool). To achieve the same magnitude of E_{GSH} response by depleting GSH exclusively would require a loss of 90% of GSH (e.g. from 10 mM to 1 mM) (Meyer and Dick, 2010). Such a massive depletion of GSH by ppm concentrations of ozone (which could generate only limited amounts of electrophiles) seems stoichiometrically unlikely. If one also considers the relative kinetics of GSSG generation and consumption (GSSG generation by GPx exhibits second order rate constants in the range of $1 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$), it appears reasonable to suggest that

the ozone effect reported in our study is primarily due to GSSG generation, and that the contribution of glutathionylated electrophile(s) formation is minor.

Thus, while there may be several different O₃-induced processes that together drive glutathione oxidation, including lipid peroxidation, H₂O₂ generation and others, the available evidence strongly suggests that the cytosolic roGFP2 responses to O₃ exposure are appropriately reporting the glutathione redox potential. Taken together, these results demonstrate that roGFP2-based sensors can be used to monitor shifts in glutathione redox homeostasis in O₃-exposed cells. Furthermore, the experimental approach utilized may be used for the validation of “oxidant stress” induced by other reactive xenobiotics in living cells.

Conclusion

These studies demonstrate the utility of using genetically-encoded fluorescent reporters in making reliable assessments of cells undergoing exposure to xenobiotics with strong oxidizing properties.

References

- Anderson ME. 1998. Glutathione: An overview of biosynthesis and modulation. *Chem Biol Interact* 111-112:1-14.
- Arthur JR. 2000. The glutathione peroxidases. *Cell Mol Life Sci* 57:1825-1835.
- Ballatori N, Krance SM, Notenboom S, Shi S, Tieu K, Hammond CL. 2009. Glutathione dysregulation and the etiology and progression of human diseases. *Biol Chem* 390:191-214.
- Ballinger CA, Cueto R, Squadrito G, Coffin JF, Velsor LW, Pryor WA, et al. 2005. Antioxidant-mediated augmentation of ozone-induced membrane oxidation. *Free Radic Biol Med* 38:515-526.
- Bargagli E, Olivieri C, Bennett D, Prasse A, Muller-Quernheim J, Rottoli P. 2009. Oxidative stress in the pathogenesis of diffuse lung diseases: A review. *Respir Med* 103:1245-1256.
- Cannon MB, Remington SJ. 2008. Redox-sensitive green fluorescent protein: Probes for dynamic intracellular redox responses. A review. *Methods Mol Biol* 476:51-65.
- Chalfant M, Bernd K. 2011. Detecting ozone-induced changes in cellular redox balance via gsh/gssg-glo™ assay. [Internet] Available from: http://www.promega.com/resources/articles/pubhub/tpub_059-detecting-ozone-induced-changes-in-cellular-redox-balance-via-gsh-gssg-glo-assay/ [accessed 14 December 2012].
- Cheng W-Y, Tong H, Miller EW, Chang CJ, Remington J, Zucker RM, et al. 2010. An integrated imaging approach to the study of oxidative stress generation by mitochondrial dysfunction in living cells. *Environ Health Perspect* 118:902-908.
- Cheng W-Y, Currier J, Bromberg PA, Silbajoris R, Simmons SO, Samet JM. 2012. Linking oxidative events to inflammatory and adaptive gene expression induced by exposure to an organic particulate matter component. *Environ Health Perspect* 120:267-274.
- Chung KF, Marwick JA. 2010. Molecular mechanisms of oxidative stress in airways and lungs with reference to asthma and chronic obstructive pulmonary disease. *Ann N Y Acad Sci* 1203:85-91.
- Ciencewicki J, Trivedi S, Kleeberger SR. 2008. Oxidants and the pathogenesis of lung diseases. *J Allergy Clin Immunol* 122:456-468.

- Dooley CT, Dore TM, Hanson GT, Jackson WC, Remington SJ, Tsien RY. 2004. Imaging dynamic redox changes in mammalian cells with green fluorescent protein indicators. *J Biol Chem* 279:22284-22293.
- Dubinina EE, Dadali VA. 2010. Role of 4-hydroxy- trans-2-nonenal in cell functions. *Biochemistry (MOSC)* 75:1069-1087.
- Gutscher M, Pauleau A-L, Marty L, Brach T, Wabnitz GH, Samstag Y, et al. 2008. Real-time imaging of the intracellular glutathione redox potential. *Nat Methods* 5:553(557).
- Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, et al. 2004. Investigating mitochondrial redox potential with redox-sensitive green fluorescent protein indicators. *J Biol Chem* 279:13044-13053.
- Helmy MH, Ismail SS, Fayed H, El-Bassiouni EA. 2000. Effect of selenium supplementation on the activities of glutathione metabolizing enzymes in human hepatoma hep g2 cell line. *Toxicology* 144:57-61.
- Holben DH, Smith AM. 1999. The diverse role of selenium within selenoproteins: A review. *J Am Diet Assoc* 99:836-843.
- Jones DP. 2008. Radical-free biology of oxidative stress. *Am J Physiol Cell Physiol* 295:C849-C868.
- Kelly FJ, Mudway I, Krishna MT, Holgate ST. 1995. The free radical basis of air pollution: Focus on ozone. *Respir Med* 89:647-656.
- Kohen R, Nyska A. 2002. Invited review: Oxidation of biological systems: Oxidative stress phenomena, antioxidants, redox reactions, and methods for their quantification. *Toxicol Pathol* 30:620-650.
- Laumbach RJ. Outdoor air pollutants and patient health. *Am Fam Physician* 81:175-180.
- Leist M, Raab B, Maurer S, Rösick U, Brigelius-Flohé R. 1996. Conventional cell culture media do not adequately supply cells with antioxidants and thus facilitate peroxide-induced genotoxicity. *Free Radic Biol Med* 21:297-306.
- Ma Q. 2010. Transcriptional responses to oxidative stress: Pathological and toxicological implications. *Pharmacol Ther* 125:376-393.
- MacNee W. 2001. Oxidative stress and lung inflammation in airways disease. *Eur J Pharmacol* 429:195-207.

- Mates JM, Sanchez-Jimenez F. 1999. Antioxidant enzymes and their implications in pathophysiologic processes. *Front Biosci* 4:D339-345.
- Meyer AJ. 2008. The integration of glutathione homeostasis and redox signaling. *J Plant Physiol* 165:1390-1403.
- Meyer AJ, Dick TP. 2010. Fluorescent protein-based redox probes. *Antioxid Redox Signal* 13:621-650.
- Morgan B, Sobotta MC, Dick TP. 2011. Measuring egsh and h2o2 with rogfp2-based redox probes. *Free Radic Biol Med* 51:1943-1951.
- Mudway IS, Kelly FJ. 2000. Ozone and the lung: A sensitive issue. *Mol Aspects Med* 21:1-48.
- Pryor WA. 1992. How far does ozone penetrate into the pulmonary air/tissue boundary before it reacts? *Free Radic Biol Med* 12:83-88.
- Pryor WA, Squadrito GL, Friedman M. 1995. A new mechanism for the toxicity of ozone. *Toxicol Lett* 82-83:287-293.
- Reddel RR, Ke Y, Gerwin BI, McMenamin MG, Lechner JF, Su RT, et al. 1988. Transformation of human bronchial epithelial cells by infection with sv40 or adenovirus-12 sv40 hybrid virus, or transfection via strontium phosphate coprecipitation with a plasmid containing sv40 early region genes. *Cancer Res* 48:1904-1909.
- Sadhu SS, Callegari E, Zhao Y, Guan X, Seefeldt T. 2012. Evaluation of a dithiocarbamate derivative as an inhibitor of human glutaredoxin-1. *J Enzyme Inhib Med Chem*.
- Schwarzlander M, Murphy MP, Duchon MR, Logan DC, Fricker MD, Halestrap AP, et al. 2012. Mitochondrial 'flashes': A radical concept rephined. *Trends Cell Biol*.
- Song H, Tan W, Zhang X. 2010. Ozone induces inflammation in bronchial epithelial cells. *J Asthma* 48:79-83.
- Srebot V, Gianicolo EA, Rainaldi G, Trivella MG, Sicari R. 2009. Ozone and cardiovascular injury. *Cardiovasc Ultrasound* 7:30.
- Tal TL, Simmons SO, Silbajoris R, Dailey L, Cho S-H, Ramabhadran R, et al. 2010. Differential transcriptional regulation of il-8 expression by human airway epithelial cells exposed to diesel exhaust particles. *Toxicol Appl Pharmacol* 243:46-54.
- Taylor-Clark TE, Undem BJ. 2010. Ozone activates airway nerves via the selective stimulation of trpa1 ion channels. *J Physiol* 588:423-433.

- Todokoro M, Mochizuki H, Tokuyama K, Utsugi M, Dobashi K, Mori M, et al. 2004. Effect of ozone exposure on intracellular glutathione redox state in cultured human airway epithelial cells. *Inflammation* 105-114.
- Van der Vliet A, O'Neil CA, Eiserich JP, Cross CE. 1995. Oxidative damage to extracellular fluids by ozone and possible protective effects of thiols. *Arch Biochem Biophys* 321:43-50.
- Wamelink M, Struys E, Jakobs C. 2008. The biochemistry, metabolism and inherited defects of the pentose phosphate pathway: A review. *J Inherit Metab Dis* 31:703-717.
- Ward PA. 2010. Oxidative stress: Acute and progressive lung injury. *Ann N Y Acad Sci* 1203:53-59.
- Yang W, Omaye ST. 2009. Air pollutants, oxidative stress and human health. *Mutat Res* 674:45-54.

Figure Legends

Figure 1. roGFP2 interactions with the glutathione system (adapted from Meyer and Dick, 2010). Glutathione peroxidases (GPx) oxidize GSH to GSSG in response to peroxides, including H_2O_2 and lipid hydroperoxides (LOOH), thus increasing the glutathione redox potential (E_{GSH}). In response to the increase in GSSG, one of the engineered vicinal cysteines of roGFP2 becomes S-glutathionylated by glutaredoxin (Grx). Glutathionylation in turn causes disulfide bond formation and alteration of the spectral properties of the GFP fluorophore. In the reductive pathway, Grx catalyzes the reduction of roGFP2 disulfide bonds through deglutathionylation as GSSG levels decrease and normal levels of GSH are reestablished by glutathione reductase (GR), at the expense of NADPH, causing a renormalization of E_{GSH} . Glucose and the pentose-phosphate pathway (PPP) create NADPH, which is used by GR to reduce GSSG to GSH.

Figure 2. Exposure to O_3 induces a dose- and time-dependent increase in the cytosolic glutathione redox potential in airway epithelial cells. BEAS-2B cells expressing cytosolic roGFP2 were exposed to clean air for 5 min followed by a 0 (Air Control), 0.15, 0.25 or 0.50 ppm O_3 exposure for 35 min in a stage-top exposure system maintained at 37 °C, >90% relative humidity, and 5 % CO_2 . Shown are the ratiometric values (404/488) calculated from the fluorescence intensity emitted at 510 nm induced by sequential excitation at 404 and 488 nm, and plotted relative to the 5 min baseline. Addition of 0.1 mM H_2O_2 at the end of the O_3 exposure produced a maximal response which was fully reversible with the addition of 10 mM DTT. The data shown were derived from 3 or more separate experiments monitoring 7 or more cells in real-time throughout the exposure period.

Figure 3. Glucose deprivation sensitizes cells to O_3 -induced roGFP2 oxidation. Shown are the responses of 7 BEAS-2B cells equilibrated in Locke Solution containing 1 mg/ml glucose

(A), or 0 mg/ml (B). Cells were exposed to 0.5 ppm O₃. Addition of 0.1 mM H₂O₂ at the end of the O₃ exposure produced a maximal response which was fully reversible with the addition of 10 mM DTT.

Figure 4. Manipulation of the glutathione system modulates roGFP2 responses to ozone. A.

Changes in cytosolic glutathione redox potential induced by 0.5 ppm O₃ exposure as reported by BEAS-2B cells expressing either roGFP2 or Grx1-roGFP2. **B.** BEAS-2B cells were pretreated with 100 μM 2-AAPA, a glutaredoxin inhibitor, prior to exposure to 0.5 ppm O₃. The responses shown are the normalized 404/488 ratios plotted relative to their established baseline. **C.** BEAS-2B cells were pretreated with 1 μM sodium selenite for 48 hours prior to 0.5 ppm O₃ exposure. To facilitate comparison of the responses in A and C, the normalized ratios were plotted as a percentage of the signals obtained at maximal oxidizing and reducing conditions achieved using 1 mM H₂O₂ and 10 mM DTT, respectively. Other experimental conditions were as described for Figure 2. Values shown are mean ± SE (n ≥ 3).

Figure 5. Comparison between roGFP2 and HyPer responses to O₃.

BEAS-2B cells expressing either roGFP2 or the H₂O₂ sensor HyPer, were exposed to 0.5 ppm O₃ as described for Figure 2. To facilitate comparison, the normalized ratios were plotted as a percentage of the signals at maximal oxidation and reduction achieved using 1 mM H₂O₂ and 10 mM DTT, respectively. Values shown are mean ± SE (n ≥ 3).

Figure 6. O₃-induced E_{GSH} changes affects the cytosol more rapidly than the mitochondrial matrix.

BEAS-2B cells expressing roGFP2 targeted to either the cytosol or mitochondria were exposed to either 0.5 ppm (roGFP2-Cyto) or 1.0 ppm (roGFP2-Mito). For direct comparison, the normalized ratios were plotted as a percentage of the maximal oxidation achieved using 1 mM

H₂O₂ and 10 mM DTT. Other experimental conditions were as described in Figure 2. Values shown are mean \pm SE (n \geq 3).




Figure 1
18x11mm (600 x 600 DPI)




Figure 2
81x49mm (300 x 300 DPI)




Figure 3
78x98mm (600 x 600 DPI)




Figure 4
119x188mm (600 x 600 DPI)




Figure 5
75x38mm (600 x 600 DPI)




Figure 6
81x42mm (600 x 600 DPI)