

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **A Time-Stratified Case-Crossover Study of Ambient Ozone Exposure and Emergency Department Visits for Specific Respiratory Diagnoses in California (2005-2008)**

Brian J. Malig, Dharshani L. Pearson, Yun Brenda Chang, Rachel Broadwin, Rupa Basu, Rochelle S. Green, and Bart Ostro

#### **Table of Contents**

**Table S1.** Geographic indicators and exposure variable means by climate zone for eligible<sup>a</sup> respiratory EDVs in California, 2005-2008.

**Table S2.** Demographics by climate zone for eligible respiratory EDVs in California, 2005-2008.

**Table S3.** Demographic comparisons of eligible respiratory EDVs for those with ozone exposures assigned compared to those missing available co-pollutant exposure data with 20km in California, 2005-2008. NH = non-Hispanic; k = km;

**Table S4.** Excess risks (95% CI) per 10ppb ozone for respiratory outcomes in one- and two-pollutant analyses restricted to the population where another pollutant metric was available for full year and warm season (May-October). Reported risks [(OR-1)\*100] are pooled estimates using random effects meta-analysis from climate zone-specific estimates obtained using conditional logistic regression comparing exposures on visit days with others of the same day of

the week within the same month, adjusting for apparent temperature (lag<sub>0</sub> and lag<sub>1-3</sub>) and county influenza visits.

**Table S5.** Climate zone-level effect estimates and standard errors for warm season (May-October) respiratory EDV types in California, 2005-2008, using the best fitting lag.

**Table S6.** Meta-regression coefficients for warm season (May- October) respiratory EDV types and climate zone-level variables in California, 2005-2008.

**Figure S1.** Excess risks (95% CI) per 10ppb ozone (lag 03) for COPD limited to persons 50 years of age and above, in one- and two-pollutant analyses restricted to the population where another pollutant metric was available for (A) full year, (B) warm season (May-October). O<sub>3</sub>=no subset, O<sub>3</sub> (subset w/NO<sub>2</sub>) = models restricted to population with nitrogen dioxide exposures available; O<sub>3</sub> (adj. for NO<sub>2</sub>) = models with same restricted population but also adjusted for nitrogen dioxide. O<sub>3</sub> (subset w/CO) = models restricted to population with carbon monoxide exposures available; O<sub>3</sub> (adj. for CO) = models with same restricted population but also adjusted for nitrogen dioxide. O<sub>3</sub> (subset w/SO<sub>2</sub>) = models restricted to population with carbon monoxide exposures available; O<sub>3</sub> (adj. for SO<sub>2</sub>) = models with same restricted population but also adjusted for sulfur dioxide. Reported risks [(OR-1)\*100] are pooled estimates using random effects meta-analysis from climate zone-specific estimates obtained using conditional logistic regression comparing exposures on visit days with others of the same day of the week within the same month, adjusting for apparent temperature (lag<sub>0</sub> and lag<sub>13</sub>) and county influenza visits.