



Estimating Pesticide Exposure from Dietary Intake and Organic Food Choices: The Multi-Ethnic Study of Atherosclerosis (MESA)

Cynthia L. Curl, Shirley A.A. Beresford, Richard A. Fenske, Annette L. Fitzpatrick, Chensheng Lu, Jennifer A. Nettleton, and Joel D. Kaufman

<http://dx.doi.org/10.1289/ehp.1408197>

Received: 30 January 2014

Accepted: 23 January 2015

Advance Publication: 5 February 2015

This article will be available in its final, 508-conformant form 2–4 months after Advance Publication. If you require assistance accessing this article before then, please contact [Dorothy L. Ritter](#), *EHP* Web Editor. *EHP* will provide an accessible version within 3 working days of request.

Estimating Pesticide Exposure from Dietary Intake and Organic Food Choices: The Multi-Ethnic Study of Atherosclerosis (MESA)

Cynthia L. Curl,¹ Shirley A.A. Beresford,² Richard A. Fenske,¹ Annette L. Fitzpatrick,²
Chensheng Lu,³ Jennifer A. Nettleton,⁴ and Joel D. Kaufman^{1,2,5}

¹Department of Environmental and Occupational Health Sciences, University of Washington, Seattle, Washington, USA; ²Department of Epidemiology, University of Washington, Seattle, Washington, USA; ³Department of Environmental Health, Harvard University, Boston, Massachusetts, USA; ⁴Department of Epidemiology, Human Genetics, and Environmental Sciences, University of Texas Health Science Center, Houston, Texas, USA; ⁵Department of Medicine, University of Washington, Seattle, Washington, USA

Address correspondence to Cynthia L. Curl, 1910 University Drive, Boise, ID 83725 USA.

Telephone: 208.426.3924. E-mail: cynthiacurl@boisestate.edu

Running title: Estimating dietary OP exposure in MESA

Acknowledgments: This research was supported by the United States Environmental Protection Agency's Science to Achieve Results (STAR) Fellowship Program (FP917129) and by contracts N01-HC-9519 through N01-HC-95165 and N01-HC-95169 from the National Heart, Lung and Blood Institute. It was also supported by the National Institute of Environmental Health Sciences through grants K24ES013195 and P30ES07033, and by internal funding from the University of Washington's Department of Environmental and Occupational Health Sciences. This publication has not been formally reviewed by the EPA. The views expressed in this document are solely

those of the University of Washington and the EPA does not endorse any products or commercial services mentioned in this publication.

Competing financial interests: The authors declare they have no actual or potential competing financial interests.

Abstract

Background: Organophosphate pesticide (OP) exposure to the US population is dominated by dietary intake. The magnitude of exposure from diet depends partly upon personal decisions such as which foods to eat and whether to choose organic food. Most studies of OP exposure rely on urinary biomarkers, which are limited by short half-lives and often lack specificity to parent compounds. A reliable means of estimating long-term dietary exposure to individual OPs is needed to assess the potential relationship with adverse health effects.

Objectives: We assessed long-term dietary exposure to 14 OPs among 4,466 participants in the Multi-Ethnic Study of Atherosclerosis, and examined the influence of organic produce consumption on this exposure.

Methods: Individual-level exposure was estimated by combining information on typical intake of specific food items with average OP residue levels on those items. In an analysis restricted to a subset of participants who reported rarely or never eating organic produce (“conventional consumers”), we assessed urinary dialkylphosphate (DAP) levels across tertiles of estimated exposure (n=480). In a second analysis, we compared DAP levels across subgroups with differing self-reported organic produce consumption habits (n=240).

Results: Among conventional consumers, increasing tertile of estimated dietary OP exposure was associated with higher DAP concentrations ($p < 0.05$). DAP concentrations were also significantly lower in groups reporting more frequent consumption of organic produce ($p < 0.02$).

Conclusions: Long-term dietary exposure to OPs were estimated from dietary intake data, and estimates were consistent with DAP measurements. More frequent consumption of organic produce was associated with lower DAPs.

Introduction

Organophosphate pesticides (OPs) have been the most commonly used insecticides in the US for more than three decades. Following the passage of the Food Quality Protection Act of 1996, which required food tolerance decisions to consider cumulative and aggregate risk (FQPA 1996), the US Environmental Protection Agency (EPA) conducted chemical-specific risk reassessments of all OPs. These reassessments resulted in substantial reductions in OP use, including the elimination of many agricultural and nearly all residential uses of OPs (Clune et al. 2012). Despite these reductions, OPs remain the primary form of insect control in American agriculture, with over 33 million pounds applied in 2007 (Grube et al. 2011). According to data from the National Health and Nutrition Examination Survey from 2003-2004, OP exposure is prevalent; metabolites of OPs were detected in the urine of more than 75% of the US population (Barr et al. 2011).

The EPA's 2006 Cumulative Risk Assessment for OPs determined that the primary route of exposure in the general US population is through diet (USEPA 2006). Studies show that consumption of an organic diet—consisting of food grown without the use of most synthetic pesticides, including OPs—can lead to a substantial and immediate reduction in OP exposure, with metabolite levels dropping below limits of detection immediately after the introduction of organic diets (Lu et al. 2006; Lu et al. 2008). Concentrations of urinary OP metabolites in children consuming organic diets are consistently below limits of detection (Curl et al. 2003; Lu et al. 2006; Lu et al. 2008).

Many studies of OP exposure employ urinary biomarkers to estimate dose. However, OP biomarkers have significant limitations as exposure assessment tools. OP metabolites have short

half-lives, only representing exposures over approximately two days prior to sample collection (Garfitt et al. 2002; Griffin et al. 1999; Kwong 2002), and within-individual measurements are highly variable (Attfield et al. 2014; Griffith et al. 2011; Kissel et al. 2005). Further, OP metabolites can be found, preformed, in food items and in the environment (Lu et al. 2005; Quirós-Alcalá L et al. 2012; Zhang X et al. 2008). If these metabolites are excreted unchanged, as has been shown in experimental studies (Forsberg et al. 2011; Timchalk et al. 2007), exposures based on urinary biomarker levels may be overestimated. Dialkylphosphate (DAP) metabolites are common byproducts of the metabolism of most OPs, and are frequently used as OP biomarkers as these six compounds represent combined exposure to at least 28 OP pesticides (Bravo et al. 2004). Since individual OPs can vary in toxicity by as much as 6,000-fold (USEPA 2006), this lack of specificity limits the utility of DAPs in risk assessment. For all of these reasons, the oft-used urinary biomarkers do not provide a gold standard for OP exposure assessment, particularly for estimation of long-term exposure. A better measure is one that would accurately quantify exposure to specific parent compounds of known toxicity and would reflect typical, rather than acute, exposures.

We assessed long-term dietary OP exposure in a cohort of 4,466 participants by combining self-reported information on typical dietary intake with average residue levels in those items from a national database. We further assessed the relationship between these estimates and urinary DAP concentrations in a subset of participants with conventional diets (n=480). This analysis of inter-method comparability was intended as a check on the face validity of our estimates. In a second subset of participants (n=240), we investigated the association between self-reported organic produce consumption habits and urinary DAP levels.

Methods

Study population

The Multi-Ethnic Study of Atherosclerosis (MESA) was initiated in 1999 to investigate the progression of subclinical cardiovascular disease among 6,814 participants from six metropolitan areas: Baltimore, Maryland; Chicago, Illinois; Forsyth County, North Carolina; Los Angeles County, California; New York, New York; and St. Paul, Minnesota (Bild et al. 2002).

Participants were recruited using random-digit dialing and mailed brochures, and were aged 45 to 84 years at enrollment with an approximately equal gender ratio. The MESA cohort is 39% Caucasian, 28% African American, 22% Hispanic, and 12% Chinese-American, and all participants were free of clinical cardiovascular disease at recruitment. MESA was approved by the institutional review boards at the universities where they were recruited, and all subjects gave written informed consent.

MESA Food Frequency Questionnaire (FFQ)

Most data collection in MESA is structured around a series of clinical exams, scheduled at approximately two-year intervals. The analysis presented here employs data collected at the most recent exam, “Exam 5”, which spanned April 2010 through February 2012. All participants attending this exam were asked to complete a modified Block-style 120-item FFQ, in which they were asked about their “usual” consumption frequency and serving size of specific foods and beverages “over the past year”. Characteristic of the Block FFQ designs, serving sizes were quantified as small, medium or large. The MESA FFQ was developed for a multi-ethnic population, and the validity of this tool has been demonstrated previously (Nettleton et al. 2009).

The MESA FFQ includes 20 line items pertaining specifically to fruits and vegetables, with each line item referring to between one and six individual foods. This study included any fruit or vegetable for which there was both intake data available from the FFQ and pesticide residue data from the USDA Pesticide Data Program (PDP, see below). The following foods were included in this analysis: apples, apple juice, asparagus, blueberries, broccoli, cantaloupe, grapes, green beans, collard greens, lettuce, mangoes, nectarines, oranges, peaches, pears, spinach, strawberries, summer squash, sweet potatoes and tomatoes. Fruit and vegetable components of mixed dishes were not considered.

Data from the Nutrition Data System for Research (NDS-R database, Nutrition Coordinating Center, Minneapolis, MN, USA) was used to estimate the relative frequency of consumption of each food within a line item. For example, if a participant reports eating one medium serving per day of “Apples, applesauce and pears”, they were assumed to consume 0.86 servings of apples, 0.08 servings of applesauce, and 0.06 servings of pears each day. Gram weights per serving were imputed according to survey data from the National Health and Nutrition Examination Survey (NHANES) (USDA 2006).

In addition to asking about food and beverage intake, participants were also asked about organic produce consumption. Specifically, participants were asked how often the fruit and vegetables they ate were organically grown, defined as “[having] a ‘USDA Organic’ label, purchased locally from an ‘organic farm’, or grown without pesticides in a home garden.” Options were “Seldom or Never,” “Sometimes,” and “Often or Always.” This study included all participants who completed an Exam 5 MESA FFQ.

The United States Department of Agriculture (USDA) Pesticide Data Program (PDP)

Comprehensive information on pesticide residues in food at “point-of-sale” locations (e.g., grocery stores) is provided by USDA through their Pesticide Data Program (USDA 2014). Since 1991, the PDP has repeatedly tested over 95 commodities, including fruits, vegetables and juices, for residues of more than 450 pesticides, including all OPs registered in the US or for which there are import tolerances. Each year, nearly 2 million analyses are conducted through the PDP, and the results are publically available (USDA 2014). PDP samples are selected without regard to country of origin, variety or organic labeling, though relatively few samples included in the PDP database are organic (e.g., in 2010, fewer than 3% of all included samples carried an organic label) (USDA 2012). Therefore, we treated the PDP data as if it represented conventionally grown produce.

Not all commodities are monitored in every year. To capture a more complete set of food items to which OPs are applied, we combined PDP data from 2008-2010. We included all OP pesticides that were detected at least once in a fruit or vegetable that was also listed on the MESA FFQ. This resulted in inclusion of the following 14 OPs: azinphosmethyl, chlorpyrifos, diazinon, dichlorovos, dimethoate, malathion, methidathion, omethoate, oxydemeton methyl, phosmet, acephate, bensulide, ethoprop and methamidophos.

Dietary exposure assessment: Food Consumption – Chemical Residue (FCCR) approach

After identifying the specific pesticides and food items to be included in this analysis, we calculated the average concentration of each OP measured in each food item. PDP samples with values below detection limits were set to zero. To calculate dietary OP exposure, we combined individual-level intake data from FFQs and average residue data from the PDP, in a food

consumption-chemical residue (FCCR) approach (MacIntosh et al. 2001). These estimates are calculated exclusively based on food intake information and do not incorporate information on self-reported organic consumption habits.

Individual-level exposures were calculated in two ways. First, we calculated exposure in units of methamidophos equivalents, to provide a metric that can inform risk assessment. We multiplied each individual's typical intake of each food item by the average residue of each OP measured on that food. The result was then multiplied by the relative toxicity of that particular OP as compared to an index chemical (methamidophos), using a Relative Potency Factor approach (USEPA 2006). This generated exposures that were then summed across pesticides and food items, to yield an estimate of total daily exposure for each participant. This value was then divided by the participant's body weight. This calculation is shown below:

$$Exposure \left(\frac{ng \text{ methamidophos equivalents}}{kg \text{ body weight} * day} \right) = \frac{\left\{ average \text{ daily intake } \left(\frac{g \text{ food}}{day} \right) * concentration \left(\frac{ng \text{ OP}}{g \text{ food}} \right) * toxicity \text{ (unitless)} \right\}}{body \text{ weight (kg)}}$$

These “methamidophos-equivalent” exposures are most useful for understanding toxicity and predicting risk, but are not directly comparable to results from urinary biomarker analyses, as the molar quantities excreted are independent of toxicity-weighting and are not affected by body weight. For comparison with measurements of urinary DAPs, we calculated individual-level exposure in units of nmols/day. Here, we converted average OP residue levels in each food item to their molar equivalents, and multiplied that quantity by each individual's reported typical intake of each food item:

$$Exposure \left(\frac{nmols \text{ OPs}}{day} \right) = \left\{ average \text{ daily intake } \left(\frac{g \text{ food}}{day} \right) * concentration \left(\frac{ng \text{ OP}}{g \text{ food}} \right) * molecular \text{ weight } \left(\frac{nmol \text{ OP}}{ng \text{ OP}} \right) \right\}$$

We then summed across food items and OPs. For this analysis, we excluded four OPs that do not metabolize to form DAPs: acephate, bensulide, ethoprop and methamidophos.

Evaluation of FCCR-based estimates

We did not hypothesize a strong correlation at the individual level between the FCCR-based exposure estimates and urinary biomarker measurements, due to the temporal mismatch between the exposures these measures represent (short- vs long-term). Instead, we hypothesized that individuals with higher estimated exposures would, in aggregate, have higher DAP concentrations in any given spot urine sample than those with lower estimated exposures.

Testing this hypothesis evaluates the inter-method comparability of the FCCR-based exposure estimate compared to the DAP measurements, and thus provides a check on the face validity of our estimates. Because consumption of organic food has been shown to reduce OP exposure (Lu et al. 2006; Lu et al. 2008), we evaluated the relationship between the FCCR-based exposure estimates (in units of nmol OPs/day) and urinary DAP concentrations exclusively in those participants who reported that they rarely or never consumed organic produce, termed “conventional consumers.”

After calculating FCCR-based exposure estimates for each MESA participant, we evaluated the range of resulting exposure estimates among the conventional consumers. We then determined the boundaries of three tertiles of exposure – high, medium and low – each designed to include an equal number of MESA participants. Each participant was then assigned to the appropriate tertile based on their estimated exposure. We then analyzed urinary DAP concentrations in urine samples collected from two subsets of conventional consumers, each of which included 240 participants:

1) *Random sample*. This comparison included three groups of 80 participants who were randomly selected from among each of the three tertiles of exposure.

2) *Demographically-matched sample*. This comparison included three groups of 80 participants who were selected from each tertile of exposure estimates. These groups were intentionally selected have similar distributions of gender, race/ethnicity, age, income and education.

Evaluation of the impact of organic consumption habits

We were also interested in understanding the influence of self-reported organic produce consumption habits on urinary DAP levels. For this analysis, we selected 80 sets of three participants. Each set contained one participant from each category of self-reported organic produce consumption: “rarely or never”; “sometimes”; or “often or always”. These sets were matched on estimated exposure, such that the standard deviation of the exposure estimates was less than 0.5 nmol OP/day within a given set. By matching on estimated exposure (by definition, a weighted metric of produce intake), we ensure that any differences in DAP concentrations are based exclusively on differences in organic consumption habits rather than differences in produce intake. We then compared urinary DAP levels across the resulting three groups. The selected sets were constructed separately from the subsets of 240 participants described in the previous section.

Urinary dialkylphosphate (DAP) biomarker analysis

Spot urine samples were collected from all MESA participants attending Exam 5 upon arrival at the clinic. Samples were frozen in multiple aliquots, shipped frozen and stored at -80°C at a central laboratory at the University of Vermont.

For this study, aliquots (1.0 mL) of selected samples were shipped frozen to the Exposure Biology Laboratory at Harvard School of Public Health. These samples were analyzed for four DAP metabolites (dimethylphosphate [DMP], dimethylthiophosphate [DMTP], diethylphosphate [DEP], and diethylthiophosphate [DETP]), using the method described by DeAlwis et al. (2009), which involves automated solid phase extraction, on-support derivatization and isotope dilution-GC/MS. Two DAPs (dimethyldithiophosphate [DMDTP] and diethyldithiophosphate [DEDTP]) were not measured, due to analytical expense and the fact that these compounds are typically found in relatively low concentrations compared to the other DAPs (Barr et al. 2011). Quality control was assessed with standards, blanks, and spiked samples. Concentrations in laboratory blanks were all below limits of detection, and since average matrix spike recoveries were high (ranging from 92% [DMP] to 105% [DETP]), samples were not recovery-adjusted.

Analysis occurred in two batches, and detection limits varied between batches. The detection limits were as follows: DMP, 0.5 and 1.0 ng/mL; DEP, 0.5 ng/mL in both sets; DMTP, 2.0 and 0.5 ng/mL; and DETP, 0.5 and 1.0 ng/mL. To avoid batch-related biases, samples of a given metabolite were censored at the higher of the two batch's detection limit. All results below the higher detection limits were assigned a value of that detection limit divided by the square root of two, a commonly applied method for substitution of censored data when the data are not expected to be normally distributed (Hornung and Reed 1990). This method is consistent with treatment of values below the limit of detection used in the CDC National Report on Human Exposure to Environmental Chemicals (CDC 2009).

Urinary creatinine concentration was measured via a colorimetric thin film methodology using the Vitros 950IRC instrument (Johnson & Johnson Clinical Diagnostics Inc, Rochester, NY).

Urinary metabolite concentrations were creatinine adjusted to account for dilution.

Statistical analysis

Differences in participant demographic and socioeconomic characteristics were examined across comparison groups using chi-square tests. Urinary DAP concentrations were compared across groups using generalized linear regression models, with DAP concentration as the dependent variable and comparison group as the independent variable. Post-hoc Tukey HSD tests were conducted to evaluate pairwise differences in mean DAP levels across tertiles. All analyses were conducted using SAS 9.3 (Cary, NC).

Results

A total of 4,466 MESA participants attended clinic Exam 5, completed the MESA FFQ and provided relevant demographic information (Table 1). Consistent with the MESA study design, this was a diverse group; 41% of these participants were Caucasian, 12% were Chinese-American, 26% were African-American, and 22% were Hispanic. The gender ratio was fairly equal, with slightly more women (53%) than men (47%). MESA is a cohort of older adults, with just over a third of this group aged 55-64, another third aged 65-74 and the remainder over 75 years of age. Among the cohort as a whole, participants who reported more frequent consumption of organic produce also reported eating more produce overall. Median produce consumption among individuals who reported that they “often or always” ate organic produce was 3.7 servings/day, compared to 3.0 servings/day among those who “sometimes” ate organic produce and 2.2 servings/day among those who “rarely or never” did so ($p < 0.0001$).

We estimated FCCR-based exposure estimates in units of mg methamidophos equivalents per kg body weight per day (mg/kg-day) for all of these participants (Table 2). These values were lognormally-distributed, with a median exposure of 2.8 ng/kg-day and an interquartile range of 1.4 to 5.2 ng/kg-day.

FCCR-based exposure estimates for comparison with urinary DAPs

FCCR-based exposure estimates were also calculated in units of nmol/day for all study participants, and the distribution of these estimates is presented in Figure 1. This distribution is highly skewed, with a range of 0 - 49.3 nmols/day, a median of 3.8 nmols/day and an interquartile range of 1.7 – 6.9 nmols/day.

In the analysis of the relationship between FCCR-based exposure estimates and urinary metabolite levels, we considered only those participants who reported that they rarely or never ate either organic fruit or organic vegetables (n=2,670, 60%). Among these participants, the lowest tertile of exposure estimates ranged from 0 – 1.8 nmol OPs/day; the middle tertile ranged from 1.8 – 4.7 nmol OPs/day; and the highest tertile ranged from 4.7– 49.3 nmol OPs/day (Table 3).

The first comparison of the FCCR-based exposure estimates and urinary DAP levels included three groups of 80 conventional consumers who were randomly selected from each tertile of estimated exposure to OPs. One individual who was randomly selected from Tertile 2 was excluded from all analyses, due to an implausibly high DAP result (33,145 nmol DAPs/g creatinine). As shown in Table 3, the lowest tertile of exposure included more men, African-American and Hispanic participants, younger individuals, and those with less education than did the higher tertiles, though this difference was only statistically significant for gender (p = 0.02).

Table 4 shows the distributions of exposure predictions in each group included in the urinary DAP comparisons. By design, the three groups compared in the random sample have distinctly different magnitudes of estimated exposure to OPs, and urinary DAP concentrations were found to be significantly different across groups (medians: 56, 79, and 104 nmol DAP/g creatinine, $p < 0.04$; Table 5 and Figure 2a). Post-hoc pairwise comparisons of the mean DAP levels in these three groups showed the highest and lowest tertile to be significantly different from one another; the middle group was not significantly different from the highest or lowest tertile.

The second analysis included a different three groups of 80 conventional consumers who were selected from each tertile of exposure estimates in order to provide each group with similar frequency distributions of gender, race/ethnicity, age, income and education. The resulting groups were all 55-56% women, 55-56% Caucasian, 15-16% Chinese, 14% African-American, 14-15% Hispanic, and similarly matched in age group, income and education (Table 3). This demographically-matched analysis yielded the same result as the random selection analysis: urinary DAP concentrations were found to be significantly different across the three tertiles of estimated OP exposure (medians: 63, 70 and 110 nmol DAP/g creatinine, $p < 0.03$; Table 5 and Figure 2b). As with the random sample selection, post-hoc pairwise comparisons of the mean DAP levels in these three groups showed the highest and lowest tertile to be significantly different from one another, but the middle group was not significantly different from the highest or lowest tertile.

Evaluation of the impact of organic consumption habits

We conducted a separate analysis of the association between urinary DAP concentration and organic consumption habits among three groups matched on FCCR-based exposure estimates but

differing by self-reported frequency of organic produce consumption (“rarely or never,” “sometimes,” and “often or always”). Among participants included in this analysis, the median FCCR-based exposure estimate was 9.0 nmol OPs/day (interquartile range 7.0 – 11.4 nmol OPs/day, Table 4). Because participants in each group were constrained to essentially match on fruit and vegetable intake (which is notably high among individuals who choose to consume organic food), the characteristics of each group in this analysis are more similar than might otherwise be expected. Participants in this comparison were significantly more likely to be women ($p < 0.001$), to have a Bachelor’s degree or higher ($p < 0.01$), and to have an income greater than \$30,000/year ($p = 0.02$), and were somewhat more likely to be Caucasian ($p = 0.06$) than the rest of the cohort (Table 3).

We observed significant differences in urinary DAP concentrations based on self-reported frequency of organic produce consumption ($p < 0.02$; Table 5 and Figure 3). The median DAP concentrations among individuals who rarely or never consumed organic produce was 163 nmol DAP/g creatinine. Among those who sometimes consumed organic produce, the median was 121 nmol DAP/g creatinine, and among individuals who often or always ate organic produce, the median was 106 nmol DAP/g creatinine.

Discussion

This study provides estimates of long-term dietary OP exposure in a population for which information on organic food consumption is also available. The estimates were consistent with the results of urinary DAP biomonitoring, increasing our confidence in this methodology. DAP biomarkers are imperfect measures of long-term exposure, due to their short half-lives, lack of specificity to parent compounds, and potential to represent exposure to preformed metabolites

(Sudakin and Stone 2011). Despite these limitations, numerous studies have successfully used DAPs to identify risk factors for OP exposure, including proximity to farmland (Lu et al. 2000), agricultural season and timing of pesticide applications (Koch et al. 2002), living with pesticide applicators (Loewenherz et al. 1997), or consuming conventional diets (Curl et al. 2003). In this study, we used these DAP biomarkers in a novel way: to assess the face validity of our proposed exposure assessment method, which suffers from none of the aforementioned limitations of the DAPs. We found that low DAP levels were measured when exposure estimates were low and higher DAP levels were measured when exposure estimates were higher.

In addition to estimating dietary OP exposure, we also observed a significant relationship between increasing consumption of organic produce and lower DAP levels among individuals who were matched on FCCR-based exposure (essentially, a weighted metrics of produce intake). This finding is consistent with previous studies showing that consumption of organic food measurably reduces OP exposure (Lu et al. 2006; Lu et al. 2008). We suggest that future studies of the effect of dietary OP intake on health include organic food consumption as a potential effect modifier in epidemiological models. This is perhaps increasingly important, as the prevalence of organic food consumption in the US is on the rise. Several studies over the past decade have reported that 40-50% of individuals and households purchase organic food at least occasionally (Bellows et al. 2010; Onyango et al. 2007; Smith et al. 2009; Zepeda and Li 2007; Zhang F. et al. 2008), consistent with the consumption frequency reported in MESA participants in this study (40%).

While this is the first study of its kind to include information on organic food consumption habits, it is not the first to use a “food consumption – chemical residue” approach to estimate

dietary exposure. An earlier Danish study evaluated the potential cumulative effects of exposure to OP and carbamate pesticides by combining dietary intake data from a nationwide food consumption survey with residue data from a Danish pesticide residue monitoring program (Jensen et al. 2003). However, this study did not include any comparisons of the estimated exposures with biological monitoring data. Macintosh and colleagues estimated exposures to 11 contaminants – including three OPs – in 120,000 US adults enrolled in the Nurses' Health Study and the Health Professionals' Follow Up Study (MacIntosh et al. 1996). In subsequent analyses, the researchers analyzed arsenic and mercury concentrations in toenail samples collected from a subset of these participants (MacIntosh et al. 1997). Using the FCCR approach, estimated arsenic and mercury exposures were compared to measured levels in the toenails, and the authors found significant, though somewhat weak, correlations (Spearman correlation coefficients of 0.15 and 0.35). These coefficients are within the range found for similarly estimated dietary intake of chlorpyrifos and urinary 3,4,5-TCPY concentrations (MacIntosh et al. 2001) and chlordecone from diet and in blood (Guldner et al. 2010). They are also similar to coefficients observed between FFQ-based estimates of intake and biomarkers of dietary carotene (Russell-Briefel et al. 1985) and polyunsaturated fat (Hunter et al. 1992).

In this study, we chose not to focus on correlation coefficients between the FCCR-based OP exposure estimates and the urinary DAP metabolites. From the outset, we did not hypothesize a strong direct correlation between individual results from these assessment methods, due to the temporal mismatch between the biological markers and the dietary information captured in the FFQ. The short half-lives of OP metabolites, and their correspondingly high intra-individual variability, are well established (Griffith et al. 2011; Kissel et al. 2005) and the purpose of this work was to develop a metric for long-term dietary OP exposure – which the DAP biomarkers

simply cannot do. Therefore, we focused the DAP analysis on providing a check of the face validity of our exposure estimates, rather than evaluating the direct agreement between the two assessment techniques, and the results of this study suggest that this approach was successful and informative.

We estimated chronic dietary OP exposure for the MESA population in units of methamidophos equivalents per kg body weight per day. Unlike urinary biomarkers, these estimates can be used to inform risk. Within the MESA cohort, the 95th percentile of exposure was 11 ng/kg-day. For comparison, the EPA's 2006 OP Cumulative Risk Assessment estimated a 95th percentile of single day dietary exposure for adults over the age of 50 years of 92 ng/kg-day (USEPA 2006). This difference may reflect both the earlier data used in that risk assessment (prior to the more recent reductions in OP use) and the fact that the EPA was predicting a single-day maximum, whereas we are predicting typical exposure over the course of a year. Given that the EPA determined that those estimates were protective of health within an 870-fold margin of error (USEPA 2006), our results do not suggest unacceptable risk using the risk benchmarks employed in that assessment, which are based on thresholds of cholinesterase inhibition.

While these levels are below current risk thresholds, those thresholds may not adequately account for the potential synergistic effects of exposure to a mixture of pesticides – effects which have been observed in several recent animal studies (Laetz et al. 2009; Laetz et al. 2013; Laetz et al. 2014). Further, these thresholds may not reflect important mechanisms of low-level toxicity, which are only beginning to be understood. The importance of understanding chronic, low-level exposures to OPs is underscored by the results of several studies of the effects of low levels of OP exposure to infants and children. Mother-child cohort studies have found prenatal maternal

urinary DAP levels to be significantly associated with attention problems and attention deficit/hyperactivity disorder in children at 5 years of age (Marks et al. 2010), poorer intellectual development at 7 years of age (Bouchard et al. 2011), and decreased cognitive development in children at one year of age and at 6-9 years (Engel et al. 2011). Another mother-child cohort study found that prenatal chlorpyrifos exposure, assessed using umbilical cord blood plasma, was associated with deficits in both memory and IQ at 7 years of age (Rauh et al. 2011). Mothers in these studies were thought to have either agricultural or residential OP exposure, in addition to dietary exposure.

Interestingly, a more recent study in a fourth cohort of mother/infant pairs found that higher prenatal maternal urinary DAP levels were associated with improved neurobehavioral outcomes among infants at five weeks of age (Yolton et al. 2013). However, the mothers with higher OP exposure also reported more frequent consumption of fruits and vegetables than those with lower OP exposure, suggesting that total produce intake (and other correlated residual confounders associated with socioeconomic status) may be critical to consider, particularly when diet is likely to be the only significant source of OP exposure.

The current study adds to the existing literature regarding the relationship between organic food consumption and DAP levels (Curl et al. 2003; Lu et al. 2006). Here, we found that—*when matched on produce intake*— individuals who reported eating organic produce at least occasionally had significantly lower urinary DAP levels than those who ate primarily conventional produce. This finding only held when produce intake was considered, as the median DAP level among individuals who consumed organic produce was higher than the median level in the conventional consumer group as a whole. On its face, this finding is

counterintuitive and perhaps even concerning, as it might suggest that organic produce is not actually free of OP pesticides. However, we hypothesize that this reflects the difference in total produce consumption among these groups. This study did not include a group of individuals who exclusively ate organic produce and it is difficult to know exactly how much of a participant's diet is organic when they report that organic produce is "often" eaten. We suspect that the conventional fraction of the total produce intake among participants who "often" consume organic produce is responsible for the DAPs present in their urine. Alternate explanations for this finding would be higher exposure to OPs (or DAPs) from environmental sources or in organic food itself, but our study includes no data to support these explanations.

While this work provides a method for assessing long-term dietary OP exposure, and supports that method with the results of urinary biomonitoring, there are several limitations. Notably, we were only able to compare these estimates to the very biomarkers we find lacking. Unfortunately, this is the state of the science, as no gold standard is available. Another limitation is that these FCCR-based estimates are based on data acquired from FFQs, which can be limited by recall bias. However, among all dietary assessment tools, FFQs are best suited to provide information for studies where typical, long-term diet is the conceptually important exposure, rather than intake on a few specific days (Willett 1998). This is a strength of using FFQ data in the current study, as individuals are known to do a better job of recalling their usual diets rather than describing what foods were eaten in any specific meal in the past (Willett 1998).

While the PDP database provides the most comprehensive OP residue data available, it is a national database, and does not reflect the specific residues to which individuals are exposed. We also did not include every food item to which OPs are applied. Some items were excluded

because no OPs were detected on any samples of that type, which is unlikely to affect the results of this study. Other items were excluded because they were not included on the MESA FFQ. We also did not include fruit and vegetable components of mixed dishes as we were concerned that this would increase uncertainty in our analyses. Future studies might benefit from including an FFQ specifically designed for pesticide exposure assessment purposes. Further, while this study benefited from the large sample size available in the Multi-Ethnic Study of Atherosclerosis, that study is limited to older adults. Future research on the ability of this method to estimate exposures among children and younger adults is warranted.

A final limitation of this study is that we did not allow produce consumption frequency and organic produce choice to vary freely within the analytic datasets. Because of the costs associated with determining the DAP concentration, we made some statistical selections to maximize efficiency while avoiding confounding. Among conventional consumers, we evaluated the relationship between FCCR-based exposure predictions and urinary DAP concentrations using two different sampling strategies and found consistent results. The first comparison included individuals who were randomly selected from across the spectrum of estimated exposure, removing the possibility of selection bias. However, the FCCR-based estimates are, by definition, highly correlated with produce consumption, which is in turn related to demographic and socioeconomic factors. Distributions of demographics and socioeconomics were notably different across the randomly selected groups. By evaluating DAP concentrations among groups matched on these characteristics, as in our second comparison, we can be fairly confident that the observed differences in DAPs were not related to these factors.

While there were limitations to this study, it also had several notable strengths. We employed a large and well-characterized cohort, for which data was collected using standardized data collection methods. The exposure estimation approach we propose allows identification of parent compounds, which in turn allows evaluation of risk. Compared to urinary metabolite analysis, the FCCR-based assessment approach is non-invasive, inexpensive, and can be easily implemented in cohorts in which FFQ data are already available (though information on organic food consumption habits may still need to be acquired). Further, this methodology provides estimates of long-term exposure, which cannot be obtained from existing biomarker methods.

The food composition – chemical residue method described in the present study may prove useful in future epidemiological studies of long-term dietary OP exposure, particularly if paired with information on organic food consumption, which may modify the observed exposure-response relationship. As concern grows regarding potential effects of low-level OP exposures, the need increases for more sophisticated exposure assessment methods. These methods must consider the relevant time frame of exposure and be able to define the parent compounds to which individuals are exposed in order to truly assess risk.

References

- Attfield KR, Hughes MD, Spengler JD, Lu C. 2014. Within- and Between-Child Variation in Repeated Urinary Pesticide Metabolite Measurements over a 1-Year Period. *Environ Health Perspect* 122(2):201-206.
- Barr DB, Wong LY, Bravo R, Weerasekera G, Odetokun M, Restrepo P, et al. 2011. Urinary concentrations of dialkylphosphate metabolites of organophosphorus pesticides: National Health and Nutrition Examination Survey 1999-2004. *Int J Environ Res Public Health* 8:3063-3098.
- Bellows AC, Alcaraz VG, Hallman WK. 2010 Gender and food, a study of attitudes in the USA towards organic, local, U.S. grown, and GM-free foods. *Appetite* 55:540-50.
- Bild DE, Bluemke DA, Burke GL, Detrano R, Diez Roux AV, Folsom AR, et al. 2002. Multi-ethnic study of atherosclerosis: objectives and design. *Am J Epidemiol* 156(9):871-881.
- Bouchard MF, Chevrier J, Harley KG, Kogut K, Vedar M, Calderon N, et al. 2011. Prenatal Exposure to Organophosphate Pesticides and IQ in 7-Year-Old Children. *Environ Health Perspect* 119(8):1189-1195.
- Bravo R, Caltabiano LM, Weerasekera G, Whitehead RD, Fernandez C, Needham LL, et al. 2004. Measurement of dialkyl phosphate metabolites of organophosphorus pesticides in human urine using lyophilization with gas chromatography-tandem mass spectrometry and isotope dilution quantification. *J Expo Anal Environ Epidemiol* 14(3):249-59.
- CDC (Centers for Disease Control and Prevention). 2009. National Report on Human Exposure to Environmental Chemicals. Available: <http://www.cdc.gov/exposurereport> [accessed 17 July 2014].
- Clune AL, Ryan PB, Barr DB. 2012. Have regulatory efforts to reduce organophosphorus insecticide exposures been effective? *Environ Health Perspect* 120(4):521-5.
- Curl CL, Fenske RA, Elgethun K. 2003. Organophosphorus pesticide exposure of urban and suburban preschool children with organic and conventional diets. *Environ Health Perspect* 111(3):377-382.
- De Alwis GK, Needham LL, Barr DB. 2009. Automated solid phase extraction, on-support derivatization and isotope dilution-GC/MS method for the detection of urinary dialkyl phosphates in humans. *Talanta* 77(3):1063-1067.

- Engel SM, Wetmur J, Chen J, Zhu C, Barr DB, Canfield RL, et al. 2011. Prenatal exposure to organophosphates, paraoxonase 1, and cognitive development in childhood. *Environ Health Perspect* 119(8):1182-1188.
- Forsberg ND, Rodriguez-Proteau R, Ma L, Morre J, Christensen JM, Maier CS, et al. 2011. Organophosphorus pesticide degradation product *in vitro* metabolic stability and time-course uptake and elimination in rats following oral and intravenous dosing. *Xenobiotica* 41:422-429.
- Food Quality Protection Act of 1996. 1996. Public Law 104–170.
- Garfitt SJ, Jones K, Mason HJ, Cocker J. 2002. Oral and dermal exposure to propetamphos: a human volunteer study. *Toxicol Lett* 134(1-3):115-118.
- Griffin P, Mason H, Heywood K, Cocker J. 1999. Oral and dermal absorption of chlorpyrifos: a human volunteer study. *Occup Environ Med* 56(1):10-13.
- Griffith W, Curl CL, Fenske RA, Lu CA, Vigoren EM, Faustman EM. 2011. Organophosphate pesticide metabolite levels in pre-school children in an agricultural community: Within- and between-child variability in a longitudinal study. *Environ Res* 111(6):751-756.
- Grube DD, Kiely T, Wu L. 2011. Pesticide Industry Sales and Usage: 2006 and 2007 Market Estimates. Washington, DC:Office of Chemical Safety and Pollution Prevention; US Environmental Protection Agency. Available: http://www.epa.gov/opp00001/pestsales/07pestsales/market_estimates2007.pdf [accessed 5 January 2015].
- Guldner L, Multigner L, Heraud F, Monfort C, Thome JP, Giusti A, et al. 2010. Pesticide exposure of pregnant women in Guadeloupe: ability of a food frequency questionnaire to estimate blood concentration of chlordecone. *Environ Res* 110(2):146-151.
- Hornung RW and Reed LD. 1990. Estimation of average concentration in the presence of non-detectable values. *App Occup Environ Hyg* 5:46-51.
- Hunter DJ, Rimm EB, Sacks FM, Stampfer MJ, Colditz GA, Litin LB, et al. 1992. Comparison of measures of fatty acid intake by subcutaneous fat aspirate, food frequency questionnaire, and diet records in a free-living population of US men. *Am J Epidemiol* 135(4):418-427.

- Jensen AF, Petersen A, Granby K. 2003. Cumulative risk assessment of the intake of organophosphorus and carbamate pesticides in the Danish diet. *Food Addit Contam* 20(8):776-85.
- Kissel JC, Curl CL, Kedan G, Lu C, Griffith W, Barr DB, et al. 2005. Comparison of organophosphorus pesticide metabolite levels in single and multiple daily urine samples collected from preschool children in Washington State. *J Expo Anal Environ Epidemiol* 15(2):164-171.
- Koch D, Lu C, Fisker-Andersen J, Jolley L, Fenske RA. 2002. Temporal association of children's pesticide exposure and agricultural spraying: report of a longitudinal biological monitoring study. *Environ Health Perspect* 110(8):829-833.
- Kwong TC. 2002. Organophosphate pesticides: biochemistry and clinical toxicology. *Ther Drug Monit* 24(1):144-149.
- Laetz CA, Baldwin DH, Collier TK, Herbert V, Stark JD, Scholz NL. 2009. The synergistic toxicity of pesticide mixtures: Implications for risk assessment and the conservation of endangered Pacific salmon. *Environ Health Perspect* 117:348-53.
- Laetz CA, Baldwin DH, Herbert V, Stark JD, Scholz NL. 2013. Interactive neurobehavioral toxicity of diazinon, malathion, and ethoprop to juvenile coho salmon. *Environ Sci Technol* 47:2925-2931.
- Laetz CA, Baldwin DH, Herbert V, Stark JD, Scholz NL. 2014. Elevated temperatures increase the toxicity of pesticide mixtures to juvenile coho salmon. *Aquat Toxicol* 146:38-44.
- Loewenherz C, Fenske RA, Simcox NJ, Bellamy G, Kalman D. 1997. Biological monitoring of organophosphorus pesticide exposure among children of agricultural workers in central Washington State. *Environ Health Perspect* 105(12):1344-1353.
- Lu C, Barr DB, Pearson MA, Waller LA. 2008. Dietary intake and its contribution to longitudinal organophosphorus pesticide exposure in urban/suburban children. *Environ Health Perspect* 116(4):537-42.
- Lu C, Bravo R, Caltabiano LM, Irish RM, Weerasekera G, Barr DB. 2005. The presence of dialkylphosphates in fresh fruit juices: implication for organophosphorus pesticide exposure and risk assessments. *J Toxicol Environ Health A* 68(3):209-227.

- Lu C, Fenske RA, Simcox NJ, Kalman D. 2000. Pesticide exposure of children in an agricultural community: evidence of household proximity to farmland and take home exposure pathways. *Environ Res* 84(3):290-302.
- Lu C, Toepel K, Irish R, Fenske RA, Barr DB, Bravo R. 2006. Organic diets significantly lower children's dietary exposure to organophosphorus pesticides. *Environ Health Perspect* 114(2):260-263.
- MacIntosh DL, Kabiru C, Echols SL, Ryan PB. 2001. Dietary exposure to chlorpyrifos and levels of 3,5,6-trichloro-2-pyridinol in urine. *J Expo Anal Environ Epidemiol* 11(4):279-285.
- MacIntosh DL, Spengler JD, Ozkaynak H, Tsai L, Ryan PB. 1996. Dietary exposures to selected metals and pesticides. *Environ Health Perspect* 104(2):202-209.
- MacIntosh DL, Williams PL, Hunter DJ, Sampson LA, Morris SC, Willett WC, et al. 1997. Evaluation of a food frequency questionnaire-food composition approach for estimating dietary intake of inorganic arsenic and methylmercury. *Cancer Epidemiol Biomarkers Prev* 6(12):1043-1050.
- Marks AR, Harley K, Bradman A, Kogut K, Barr DB, Johnson C, et al. 2010. Organophosphate pesticide exposure and attention in young Mexican-American children: the CHAMACOS study. *Environ Health Perspect* 118(12):1768-1774.
- Nettleton JA, Rock CL, Wang Y, Jenny NS, Jacobs DR. 2009. Associations between dietary macronutrient intake and plasma lipids demonstrate criterion performance of the Multi-Ethnic Study of Atherosclerosis (MESA) food-frequency questionnaire. *Br J Nutr* 102(8):1220-1227.
- Onyango BM, Hallman WK, Bellows AC. 2007. Purchasing organic food in US food systems - A study of attitudes and practice. *Br Food J* 109:399-411.
- Quirós-Alcalá L, Bradman A, Smith K, Weerasekera G, Odetokun M, Barr DB, et al. 2012. Organophosphorous pesticide breakdown products in house dust and children's urine. *J Expo Sci Environ Epidemiol* 22(6):559-68.
- Rauh V, Arunajadai S, Horton M, Perera F, Hoepner L, Barr DB, et al. 2011. Seven-year neurodevelopmental scores and prenatal exposure to chlorpyrifos, a common agricultural pesticide. *Environ Health Perspect* 119(8):1196-1201.

- Russell-Briefel R, Caggiula AW, Kuller LH. 1985. A comparison of three dietary methods for estimating vitamin A intake. *Am J Epidemiol* 122(4):628-636.
- Smith T, Huang C, Lin B. 2009. Does price or income affect organic choice? Analysis of US fresh produce users. *J Agric App Econ* 41:731-44.
- Sudakin DL, Stone DL. 2011. Dialkyl phosphates as biomarkers of organophosphates: the current divide between epidemiology and clinical toxicology. *Clin Toxicol (Phila)* 49(9):771-781.
- Timchalk C, Busby A, Campbell JA, Needham LL, Barr D.B. 2007. Comparative pharmacokinetics of the organophosphorus insecticide chlorpyrifos and its major metabolites diethylphosphate, diethylthiophosphate and 3,5,6-trichloro-2-pyridinol in the rat. *Toxicology* 23:145-157.
- USDA (United States Department of Agriculture). 2006. What we eat in America, NHANES 2003-2004 Data. Available: <http://sepri.ars.usda.gov/Services/docs.htm?docid=13793> [accessed 20 July 2014].
- USDA (United States Department of Agriculture). 2012. Pesticide Data Program Annual Summary, Calendar Year 2010. Available: <http://www.ams.usda.gov/pdp> [accessed 5 January 2015].
- USDA (United States Department of Agriculture) Pesticide Data Program (PDP). 2014. Available: <http://www.ams.usda.gov/AMSV1.0/PDP> [accessed 17 July 2014].
- Willett W. 1998. *Nutritional Epidemiology*. 2nd ed. New York: Oxford University Press
- USEPA. 2006. Organophosphorous Cumulative Risk Assessment - 2006 Update. Washington, DC. Available: http://www.epa.gov/pesticides/cumulative/2006-op/op_cra_main.pdf [accessed 5 January 2015].
- Yolton K, Xu Y, Sucharew H, Succop P, Altaye M, Popelar A, et al. 2013. Impact of low-level gestational exposure to organophosphate pesticides on neurobehavior in early infancy: a prospective study. *Environ Health* 12(1):79.
- Zepeda L, Li J. 2007. Characteristics of organic food shoppers. *J Agric App Econ* 39:17-28.
- Zhang X, Driver JH, Li Y, Ross JH, Krieger RI. 2008. Dialkylphosphates (DAPs) in fruits and vegetables may confound biomonitoring in organophosphorus insecticide exposure and risk assessment. *J Agric Food Chem* 56(22):10638-10645.

Zhang F, Huang CL, Lin B-H. 2008. Modeling fresh organic produce consumption with scanner data: a generalized double hurdle model approach. *Agribusiness* 24:510-22.

Table 1. Demographic distributions of all MESA participants completing an FFQ at Exam 5 (n=4,466), and demographics of participants in subgroups based on self-report of organic produce consumption habits.

	n	Gender		Race/Ethnicity				Age			Annual Household Income ^a			Education ^b		
		Female	Male	White	Chinese	Black	Hispanic	<65 yrs	65-74 yrs	>75 yrs	<\$30K	\$30K-\$75K	>\$75K	High school or less	Some college	Bachelor's or higher
Full cohort	4466	53%	47%	41%	12%	26%	22%	35%	32%	33%	33%	39%	28%	31%	29%	39%
Self-reported frequency of organic produce consumption																
"Rarely or never"	2670	51%	49%	40%	12%	26%	23%	31%	31%	38%	37%	40%	23%	37%	28%	35%
"Sometimes"	1574	55%	45%	43%	11%	27%	19%	39%	33%	27%	27%	38%	35%	23%	32%	45%
"Often or always"	222	65%	35%	39%	10%	23%	27%	44%	34%	23%	34%	35%	32%	24%	29%	48%

^a147 participants were missing information on income. ^b7 participants were missing information on education.

Table 2. Percentiles of FCCR-based exposure estimates (ng methamidophos equivalents / kg body weight-day) for all participants completing the Exam 5 FFQ, and for subgroups based on self-report of organic produce consumption habits.

	<i>n</i>	Percentile of FCCR-Based Exposure Estimates ^a (ng methamidophos equivalents/kg body weight-day)				
		10%	25%	50%	75%	90%
Full cohort	4466	0.69	1.4	2.8	5.2	8.6
Self-reported frequency of organic produce consumption						
“Rarely or never”	2670	0.57	1.2	2.4	4.6	7.8
“Sometimes”	1574	0.93	1.8	3.4	5.7	9.4
“Often or always”	222	1.5	2.3	4.0	6.8	11.0

^aThese exposure estimates do not incorporate information on organic consumption habits; they are based exclusively on self-reported produce intake and residue levels in foods. The higher exposure estimates among individuals reporting that they “often or always” consume organic food is reflective of the fact that this group eats more produce than those who eat organic food less frequently or not at all.

Table 3. Demographic distributions of participants who were selected for urinary metabolite analysis.

	n	Gender		Race/Ethnicity				Age			Annual Household Income			Education		
		Female	Male	White	Chinese	Black	Hispanic	<65 yrs	65-74 yrs	>75 yrs	<\$30K	\$30K-\$75K	>\$75K	High school or less	Some college	Bachelor's or higher
Subgroups Selected for Urinary Metabolite Comparison – Conventional Consumers^a																
Random Sample^b																
Tertile 1	80	38%	63%	45%	9%	26%	20%	35%	38%	28%	28%	38%	35%	35%	28%	38%
Tertile 2	79 ^c	48%	52%	51%	18%	16%	15%	25%	37%	38%	41%	30%	29%	29%	32%	39%
Tertile 3	80	59%	41%	56%	8%	20%	16%	28%	30%	43%	26%	43%	31%	21%	28%	51%
Demographically-Matched Sample^d																
Tertile 1	80	55%	45%	56%	15%	14%	15%	31%	41%	28%	39%	34%	28%	30%	28%	43%
Tertile 2	80	56%	44%	55%	16%	14%	15%	31%	41%	28%	39%	34%	28%	30%	28%	43%
Tertile 3	80	56%	44%	56%	16%	14%	14%	31%	43%	27%	38%	35%	28%	30%	28%	43%
Subgroups Selected for Urinary Metabolite Comparison – by Organic Produce Consumption Habits																
“Rarely or never”	80	60%	40%	53%	9%	26%	13%	31%	36%	33%	21%	46%	33%	24%	25%	51%
“Sometimes”	80	66%	34%	43%	9%	21%	28%	40%	35%	25%	26%	48%	26%	25%	33%	43%
“Often or always”	80	66%	34%	45%	13%	20%	23%	43%	34%	24%	29%	38%	34%	20%	26%	54%

^aComparisons among conventional consumers are across tertiles of estimated dietary exposure to OPs. The lowest tertile (Tertile 1) includes individuals with estimated exposures of less than 1.8 nmol/day; the middle tertile (Tertile 2) includes individuals with estimated exposures ranging from 1.8-4.7 nmol/day; and the highest tertile (Tertile 3) includes individuals with estimated exposures greater than 4.7 nmol/day. ^b80 participants were randomly selected from each tertile of predicted exposure. ^cOne participant was excluded due to an implausibly high urinary DAP measurement (>30,000 nmol DAP/g creatinine). ^dParticipants in this analysis were selected to provide three groups of 80 participants with similar frequencies of each demographic characteristic shown.

Table 4. Percentiles of FCCR-based exposure estimates (nmol OPs/day) for of participants who were selected for urinary metabolite analysis.

	<i>n</i>	Percentile of FCCR-Based Exposure Estimates (nmol OPs/day)				
		10%	25%	50%	75%	90%
Subgroups Selected for Urinary Metabolite Comparison – Conventional Consumers^a						
Random Sample^b						
Tertile 1	80	0.3	0.5	1.0	1.5	1.7
Tertile 2	79 ^c	2.1	2.4	3.2	3.9	4.6
Tertile 3	80	5.2	6.0	7.5	10.7	13.0
Demographically-Matched Sample^d						
Tertile 1	80	0.5	0.9	1.1	1.6	1.7
Tertile 2	80	2.3	2.5	3.2	4.0	4.6
Tertile 3	80	5.5	5.9	7.2	9.4	12.3
Subgroups Selected for Urinary Metabolite Comparison – by Organic Produce Consumption Habits^e						
“Rarely or never”	80	5.9	6.9	9.1	11.3	13.8
“Sometimes”	80	6.0	7.0	9.0	11.4	13.8
“Often or always”	80	6.1	6.9	8.9	11.6	13.8

^aComparisons among conventional consumers are across tertiles of estimated dietary exposure to OPs. The lowest tertile (Tertile 1) includes individuals with estimated exposures of less than 1.8 nmol/day; the middle tertile (Tertile 2) includes individuals with estimated exposures ranging from 1.8-4.7 nmol/day; and the highest tertile (Tertile 3) includes individuals with estimated exposures greater than 4.7 nmol/day. ^b80 participants were randomly selected from each tertile of predicted exposure. ^cOne participant was excluded due to an implausibly high urinary DAP measurement (>30,000 nmol DAP/g creatinine). ^dParticipants in this analysis were selected to provide three groups of 80 participants with similar frequencies of relevant demographic characteristics. ^eParticipants in this analysis were selected to provide three groups who were intentionally matched on FCCR-based exposure estimate (a metric of produce intake weighted by frequency and magnitude of OP residues detected in each food item). This is reflected in the similar values across the percentiles of exposure.

Table 5. Percentiles of urinary DAP concentrations (nmol DAPs/g creatinine) by tertile of FCCR-based exposure estimates among conventional consumers and by self-report of organic produce consumption frequency.

	<i>n</i>	Percentile of urinary DAP concentration (nmol DAPs/g creatinine) ^a				
		10%	25%	50%	75%	90%
Subgroups Selected for Urinary Metabolite Comparison – Conventional Consumers						
<i>Random Sample</i>						
Tertile 1	80	24	33	56	115	228
Tertile 2	79 ^b	33	48	79	158	275
Tertile 3	80	36	67	104	241	489
<i>Demographically-Matched Sample</i>						
Tertile 1	80	29	42	63	115	197
Tertile 2	80	31	47	70	137	225
Tertile 3	80	40	63	110	217	414
Subgroups Selected for Urinary Metabolite Comparison – by Organic Produce Consumption Habits						
“Rarely or never”	80	48	80	163	365	638
“Sometimes”	80	39	58	121	237	474
“Often or always”	80	36	54	106	204	321

^aDAP detection frequencies were as follows: DMP = 80%; DEP = 73%; DMTP = 51%; and DETP = 16%. At least one DAP was detected in 93% of the samples analyzed. ^bOne participant was excluded due to an implausibly high urinary DAP measurement (>30,000 nmol DAP/g creatinine).

Figure Legends

Figure 1. Distribution of exposure estimates (nmols OPs / day) in the MESA population (n=4,466). This distribution ranges from 0-49.3 nmols/day (median = 3.8; interquartile range = 1.7-6.9). For analyses occurring within the subset of participants who rarely or never consume organic food (n=2,670), the tertiles of exposure were demarcated as follows: Tertile 1 < 1.8 nmol/day; Tertile 2 = 1.8-4.7 nmol/day; Tertile 3 > 4.7 nmol/day.

Figure 2. Urinary dialkylphosphate concentrations (nmol DAPs / g creatinine) by tertile of FCCR-based exposure estimates (nmol OPs / day). Boxes extend from the 25th to the 75th percentile, horizontal bars represent the median, whiskers extend 1.5 times the length of the interquartile range (IQR) above and below the 75th and 25th percentiles, respectively, and outliers are represented as stars. **a.** This “random sample” comparison includes 80 participants randomly selected from each tertile of FCCR-based exposure estimates. One participant in the second tertile was excluded due to an implausibly high urinary DAP concentration (>30,000 nmol DAP/g creatinine). Urinary DAPs were significantly different across the three groups (p<0.04). **b.** This “demographically-matched” comparison includes 80 participants from each tertile, which were selected to provide groups with similar age, gender, race/ethnicity, income and education distributions. Urinary DAPs were significantly different across the three groups (p<0.03). One outlier, which was in the “High estimated exposure” group, was not shown to preserve scale (value of 2017 nmol DAPs/g creatinine).

Figure 3. Urinary dialkylphosphate concentrations (nmol DAPs / g creatinine) by self-report frequency of organic produce consumption. Boxes extend from the 25th to the 75th percentile, horizontal bars represent the median, whiskers extend 1.5 times the length of the interquartile range (IQR) above and below the 75th and 25th percentiles, respectively, and outliers are represented as stars. Urinary DAP concentrations were significantly different across groups (p<0.02). Two outliers, both of which were in the “rarely or never” group, were not shown to preserve scale (values of 3187 and 3707 nmol DAPs/g creatinine).

Figure 1.





Figure 2.

a.



b.




Figure 3.

