

Prospective Study of Ambient Particulate Matter Exposure and Risk of Pulmonary Embolism in the Nurses' Health Study Cohort

Vivian C. Pun, Jaime E. Hart, Christopher Kabrhel, Carlos A. Camargo Jr., Andrea A. Baccarelli, and Francine Laden

<http://dx.doi.org/10.1289/ehp.1408927>

Received: 5 July 2014

Accepted: 6 May 2015

Advance Publication: 8 May 2015

This article will be available in its final, 508-conformant form 2–4 months after Advance Publication. If you need assistance accessing this article before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Prospective Study of Ambient Particulate Matter Exposure and Risk of Pulmonary Embolism in the Nurses' Health Study Cohort

Vivian C. Pun,^{1,2} Jaime E. Hart,^{2,3} Christopher Kabrhel,^{3,4} Carlos A. Camargo Jr.,^{3,4,5} Andrea A. Baccarelli,^{2,5} and Francine Laden^{2,3,5}

¹Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Shatin, Hong Kong; ²Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; ³Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA; ⁴Department of Emergency Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA; ⁵Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

Address correspondence to Francine Laden, Channing Division of Network Medicine, 401 Park Drive, Landmark Center, 3rd Floor West, Boston, MA 02215 USA. Telephone: 617.525.2711. E-mail: francine.laden@channing.harvard.edu

Running title: Particulate matter and pulmonary embolism risk

Acknowledgments: This work was funded by NIH grants P01 CA87969, R01 ES017017, R01 HL116854 and UM1 CA186107. Dr. Pun was supported by Global Scholarship Programme for Research Excellence-CNOOC grant from the Chinese University of Hong Kong.

Competing financial interests: The authors declare they have no actual or potential competing financial interests.

Abstract

Background: Pulmonary embolism (PE) is the most serious manifestation of venous thromboembolism and a leading cause of sudden death. Several studies have suggested associations of venous thromboembolism with short-term particulate matter (PM) exposure; evidence on long-term PM and traffic exposure is mixed.

Objectives: We examined the association of long-term exposure to $PM_{2.5}$, $PM_{2.5-10}$ and PM_{10} , and distance to roadways with overall incident PE and with PE subtypes in a cohort of US women.

Methods: The study included 115,745 women from the Nurses' Health Study, followed from 1992–2008. Incident PE cases were self-reported biennially. Nonidiopathic PE were cases for which the medical record revealed an underlying health condition related to PE (i.e., surgery, trauma or malignancy); idiopathic PE were cases with no such history. We used spatiotemporal models combining spatial smoothing and geographic covariates to quantify exposure at residential addresses, and Cox proportional hazards models to calculate hazard ratios (HR) and 95% confidence intervals (CI).

Results: $PM_{2.5}$ averaged over 1 month (HR=1.22, 95% CI: 1.04, 1.44) or 12 months (HR=1.17, 95% CI: 0.93, 1.48) was associated with incident PE, after adjusting for known risk factors and $PM_{2.5-10}$. Equivalent analyses restricted to PE subtypes showed a positive association for $PM_{2.5}$ with nonidiopathic PE, but not with idiopathic PE. We did not find evidence of an association between distance to roadways and PE risk.

Conclusions: We provide evidence that PM in the prior 1 and 12 months is associated with PE risk. Our results also suggest that women with underlying health conditions may be more susceptible to PE after PM exposure.

Introduction

Exposure to particulate matter (PM) air pollution has been associated with increased short- and long-term morbidity and mortality from heart disease and stroke (Brook et al. 2004; 2010; Dockery et al. 1993; Hoek et al. 2013; Miller and Siscovick 2007; Pope et al. 2002). One of the plausible mechanistic pathways mediating such effects is hypercoagulability and enhanced thrombosis (Baccarelli et al. 2007; Brook et al. 2004; Utell and Frampton 2002). Procoagulant abnormalities are a hallmark of venous thromboembolism (VTE), which is the third most common cardiovascular disease after acute myocardial infarction and stroke, with annual incidence of 1–2 per 1,000 persons in developed countries (Oger 2000; Previtali et al. 2011; Rosendaal 1999). VTE encompasses deep vein thrombosis (DVT) and pulmonary embolism (PE). DVT refers to the formation of one or more blood clots in one of the body's large veins, most commonly in the lower limbs; PE, the most serious manifestation of VTE, typically occurs when a portion of the blood clot breaks off and travels to the heart, ultimately lodging in the arteries of the lungs, potentially leading to death and disability (Office of the Surgeon General 2008; Piazza and Goldhaber 2006). PE is a leading cause of sudden death (Kürkciyan and Meron 2000). Studies have showed that the mortality rate for patients with PE is 2- to 4-fold higher than that for patients with DVT, and 10–20% of PE patients die within 3 months after the event (Goldhaber et al. 1999; Naess et al. 2007).

Until recently, little attention has been given to understanding the association between PM exposure and incident VTE or PE risk compared to the attention given to heart disease and stroke. A limited number of short- and long-term studies have provided some insight into a potential association, though the findings are conflicting (Baccarelli et al. 2008; 2009; Dales et

al. 2010; Kan et al. 2011; Martinelli et al. 2012; Milojevic et al. 2014; Shih et al. 2011).

Therefore, in this study, we used data from the Nurses' Health Study (NHS), a large and well-defined prospective cohort, to examine the association between long-term particulate exposures, distance to roadways and the risk of incident PE.

Methods

Study population

The NHS began in 1976 with 121,701 female registered nurses ages 30–55 years. These nurses lived in one of 11 states in the U.S. (California, Texas, Florida, Massachusetts, Pennsylvania, Ohio, New York, New Jersey, Michigan, Connecticut, and Maryland) at enrollment, but later relocated throughout the United States, with at least 10 nurses in each of the 48 contiguous states by 1992. The NHS is approved by the institutional review board of the Brigham and Women's Hospital, and participants provided implied informed consent through completion of study questionnaires. For the current study, we included nurses who had at least a single residential address in the contiguous United States between 1992–2008 if they had not previously reported a PE and had information available on the exposures of interest.

Outcome assessment

Biennial questionnaires on risk factors and health outcomes have been mailed to nurses since 1976 (Colditz et al. 1997). Details of the case identification and confirmation have been reported elsewhere (Kabrhel et al. 2011). Briefly, on each questionnaire, nurses were asked to report whether they had received a physician-diagnosis of PE during the previous 2 years. For nurses who reported PE diagnosis and had no history of malignancy, medical records were obtained and

reviewed by study physicians blind to exposure status. Imaging was considered diagnostic of PE if a ventilation/perfusion lung scan was read by a radiologist as high probability for PE, or if there was a filling defect on contrast enhanced computed tomography of the pulmonary vasculature or on catheter-based pulmonary angiography. Idiopathic PE was defined by the absence of recent surgery, major trauma or active malignancy listed in the medical record. A diagnosis of nonidiopathic PE was made based on the presence of either: (i) surgery or major trauma within one month of PE diagnosis as listed in the medical record or (ii) history of active malignancy and self-reported physician-diagnosed PE. Given the severity of diagnosis, followed by extensive treatment, therapeutic monitoring and clinical follow-up required of patients with VTE, PE cases are likely reported with high accuracy in our population of nurses with high health literacy. This is supported by our ongoing medical record review of PE cases, and by a validation study of 101 self-reported cases of DVT for which medical records were available. We found that 95 (94%) cases were confirmed, 2 (2%) cases were probable, and only 4 (4%) cases were not confirmed. Unlike PE, DVT diagnosis has not been routinely tracked in the NHS; therefore, the current study focused only on PE cases.

Exposure assessment

Monthly predictions of particulate matter less than 10 (PM_{10}), less than 2.5 ($PM_{2.5}$), or between 2.5 and 10 microns ($PM_{2.5-10}$) were available for all residential addresses from nationwide spatio-temporal models. Details of the model development process have been provided elsewhere (Paciorek et al. 2009; Yanosky et al. 2008; 2009; 2014). Briefly, the generalized additive mixed models (GAMM) incorporated monitoring data from the U.S. Environmental Protection Agency Air Quality System (AQS) (US EPA 2009), the Visibility Information Exchange Web System

(VIEWS 2010), the Interagency Monitoring of Protected Visual Environments (IMPROVE) network (CIRA 2013), Stacked Filter Unit (a predecessor to IMPROVE) (National Park Service), Clean Air Status and Trends (CASTNet) networks (US EPA), and Harvard research studies (Spengler and Koutrakis 1996; Suh et al. 1997) along with a number of Geographic Information System (GIS)-derived and meteorological covariates and spatial smoothing to estimate monthly prediction surfaces from 1989 forward. Due to the lack of PM_{2.5} monitoring data before 1999, separate GAMM models were constructed for 1988–1998 and 1999–2007 (Yanosky et al. 2014). The pre-1999 model added extinction coefficients and a simpler spatiotemporal structure to estimate the PM_{2.5} to PM₁₀ ratio seasonally, whereas the post-1999 PM_{2.5} model included point-source PM_{2.5} emissions. The models were evaluated using cross-validation; they exhibited little bias (slopes of the regression between predicted and measured of 0.87, 0.77, and 0.76 for PM_{2.5}, PM₁₀, and PM_{2.5-10}, respectively) and high precision (mean absolute errors of 1.81, 5.44, and 4.73 µg/m³ for PM_{2.5}, PM₁₀, and PM_{2.5-10}, respectively) (Yanosky et al. 2014).

We used distance from each residence to the nearest road as a proxy for traffic-related air pollution exposure. Specifically, the distance (in meters, m) from each residential address to the closest U.S. Census Feature Class Code A1 (roads with limited access, typically interstates), A2 (major, noninterstate roads), and A3 (secondary roads, typically with more than two lanes) road segments from the ESRI StreetMap Pro 2007 file were determined using a GIS (ArcGIS, version 9.2; ESRI, Redlands, CA). Distance to the closest road was categorized as 0–49 m, 50–199 m, 200–499 m or ≥ 500 m, based on the distribution of distance in the cohort and exposure studies showing exponential decay in a number of exposures with increasing distance from a road (Karner et al. 2010).

Covariates

To assess potential confounding, time-varying data from the biennial questionnaires were used for the following covariates chosen *a priori* based on previous associations with either PE or air pollution: age (in months), calendar year, season, region of residence (Northeast, Midwest, South, West), body mass index (BMI), smoking status (never, former, or current), pack-years of smoking (number of packs/day multiplied by number of years of cigarette smoking), having a high-risk diet according to the Alternative Healthy Eating Index score (Chiuve et al. 2012), total caloric intake, physical activity [< 3 , 3 to < 9 , 9 to < 18 , 18 to < 27 , or ≥ 27 metabolic equivalent (MET) hour/week], parity (nulliparous, 1, 2–3, or ≥ 4 children), menopausal status and hormone use, nonaspirin nonsteroidal anti-inflammatory drug use, warfarin use, multivitamin use, hypertension, coronary heart disease (any history of angina, coronary artery stenosis, or myocardial infarction), and rheumatologic disease (any history of systemic lupus erythematosus or rheumatoid arthritis). Race (white or nonwhite) and spouse's highest educational attainment (high school, college, or graduate school) obtained from the 1992 questionnaire were included. Medication and vitamin use, hypertension, coronary heart disease, and rheumatological disease were treated as dichotomous variables. Area-level socioeconomic status at the U.S. Census tract level was adjusted for using median household value and median family income (US Census Bureau 2000).

Statistical analysis

Time-varying Cox proportional hazards models were used to assess the association of PE for each $10 \mu\text{g}/\text{m}^3$ increase in predicted $\text{PM}_{2.5}$, PM_{10} , and $\text{PM}_{2.5-10}$ exposures (after assessing linearity with cubic regression splines) and categories of distance to road. All survival models

were based on a monthly time scale, and were used to estimate hazard ratios (HRs) and 95% confidence intervals (CIs). Person-months of follow-up time were calculated from baseline (30 June 1992) until PE diagnosis, the end of follow-up (30 June 2008), censoring (moving outside the geographic region of interest or loss to follow-up) or death. We estimated the effect of average particulate exposure in the prior 12 months, because previous research in this cohort has shown time period to be the most relevant exposure for cardiovascular outcomes (Puett et al. 2009). In separate models, we also considered average exposure in the prior 1 month, given that shorter-term exposures may be more relevant for PE (Dales et al. 2010; Martinelli et al. 2012; Office of the Surgeon General 2008). We examined $PM_{2.5}$, PM_{10} , and $PM_{2.5-10}$ individually in single pollutant models. In sensitivity analysis, we assessed the robustness of the associations by examining $PM_{2.5}$ and $PM_{2.5-10}$ simultaneously in a multipollutant model. Basic Cox models were stratified by age in months and adjusted for year (linear term), region of residence and season. Confounding from covariates was evaluated individually in separate models and also simultaneously in full multivariable models. In addition, we conducted models restricted to after 2000 (when direct monitoring information on $PM_{2.5}$ became available) and models restricted to women who had not moved for at least 5 years. All statistical analyses were conducted using SAS version 9.3 software (SAS Institute, Inc., Cary, North Carolina).

Results

A total of 115,745 nurses were available for analysis, and selected participant characteristics during follow-up are presented in Table 1. Throughout the follow-up period, most were never or former smokers, white, and had given birth to at least two children. Nearly half of the nurses reported < 9 MET hours per week of physical activity, and were postmenopausal with no or past

HRT use. Mean (\pm standard deviation) levels of $PM_{2.5}$, $PM_{2.5-10}$ and PM_{10} exposures in the previous 12 months were 12.6 ± 3.0 , 20.8 ± 5.8 , and $8.2 \pm 4.2 \mu\text{g}/\text{m}^3$, respectively. The majority had residential addresses located ≥ 200 m from the nearest road, and only 15% were <50 m from the nearest road.

Over 1,725,024 person-years of follow-up, we identified 1,091 total cases of incident PE; 323 (29.6%) were confirmed idiopathic, 526 (48.2%) were nonidiopathic (283 in subjects with malignancy, 243 associated with surgery or trauma), and the subtype could not be determined for 242 (22.2%) cases (Table 2). The associations for spatiotemporal model predicted ambient average $PM_{2.5}$, $PM_{2.5-10}$ and PM_{10} in the previous 1 or 12 months with incident PE and the idiopathic and nonidiopathic subtypes are presented in Table 2. In basic models adjusted for age, calendar time, season and region of residence, each $10 \mu\text{g}/\text{m}^3$ increase in prior 1-month or 12-month average $PM_{2.5}$, PM_{10} , or $PM_{2.5-10}$ was associated with an increased risk of incident PE, and most associations were statistically significant. However, these increased risks were mostly observed within the nonidiopathic subtype for both time periods.

After adding each potential time-varying confounder one at a time to the basic model, the relationships of PM exposures with PE did not change appreciably relative to the basic models (data not shown). Multivariable adjusted models included race, BMI, smoking status, pack-years, diet, total caloric intake, physical activity, husband's education attainment, parity, menopausal status, hypertension, coronary heart disease, rheumatologic disease, nonaspirin nonsteroidal anti-inflammatory drug use, warfarin use, multivitamin supplement use, median family income in census tract of residence and median house value in census tract of residence (Table 2). Results from multivariable models were similar to those from basic models (Table 2), as were results

from models restricted to after 2000 or to women who had not moved for at least 5 years (data not shown).

Table 3 shows the sensitivity analyses where we considered exposures to PM_{2.5} and PM_{2.5-10} simultaneously. Most HRs for incident PE were modestly attenuated relative to HRs in the single size fraction models, except for the estimate of 10 µg/m³ increment in PM_{2.5} level averaged over the prior 1 month [HR=1.22 (95% CI: 1.04, 1.44)]. Equivalent analyses restricted to specific PE subtypes showed that PM_{2.5} exposure averaged over the prior 1 and 12 months was associated with increased risk of nonidiopathic PE [1 month HR=1.30 (95% CI: 1.03, 1.65); 12 month HR=1.20 (95% CI: 0.86, 1.67)] upon adjusting for PM_{2.5-10}, but not with idiopathic PE. Although elevated incident PE risk was linked to PM_{2.5-10} exposure in single size fraction models, the inclusion of PM_{2.5} led to a decrease in HRs, and the associations were no longer statistically significant.

Lastly, we did not find evidence of an association between distance to roadways and PE risk (Table 4). For nurses living within 49 m of roads compared to those living further away (≥ 500 m), the multivariable adjusted HRs for incident PE, idiopathic PE and nonidiopathic PE were 1.02 (95% CI: 0.83, 1.25), 0.92 (95% CI: 0.63, 1.33), and 1.20 (95% CI: 0.90, 1.61), respectively.

Discussion

In the NHS cohort of US women, we observed an increased risk of incident PE for each 10 µg/m³ increase in prior 1 (HR=1.22; 95% CI: 1.04, 1.44) and 12 month (HR=1.17; 95% CI: 0.93, 1.48) average exposures to PM_{2.5} after controlling for known risk factors and PM_{2.5-10}.

Equivalent analyses limited to specific PE subtypes showed statistically significantly positive

association for PM_{2.5} with nonidiopathic PE, but not with idiopathic PE. We did not find evidence of an association between distance to roadways and incident PE risk.

To our knowledge, this is the first study to report positive association of particulate exposures with incident PE risk in a North American population. To date, only three epidemiologic studies have examined the association of long-term exposure to PM with VTE risk, and there has been little consistency in outcomes or exposures across the studies. In the first studies to examine the impact of pollution on DVT, Baccarelli et al. (2008) observed that each 10 µg/m³ increase in PM₁₀ during the prior year was associated with a 70% elevation in DVT risk in healthy subjects in the Lombardia Region of Italy. The risk was smaller in women (40% increase) and limited to those who were not using oral contraceptive or hormone therapy at the time of diagnosis. Two prospective American cohorts, the Women's Health Initiative (WHI) Hormone Therapy trials and the Atherosclerosis Risk in Communities (ARIC) Study, reported no association between measures of PM₁₀ and PM_{2.5} exposures and VTE risk (Kan et al. 2011; Shih et al. 2011). Our finding on distance to roadways and incident PE generally agrees with another American study reporting a null association of traffic density and distance to roads with VTE risk (Kan et al. 2011), but is in contrast to an Italian case-control study that found inverse association between distance to roads and risk of DVT (Baccarelli et al. 2009). A possible explanation for the discrepant findings is the different primary disease outcomes across studies. In the ARIC and WHI studies, the main outcome was idiopathic and nonidiopathic combined VTE (Kan et al. 2011; Shih et al. 2011), whereas the main outcome in the Italian studies was first incident of lower limb DVT with or without symptomatic PE (Baccarelli et al. 2008; 2009). We focused our analysis on incident PE, which is the most serious manifestation of VTE. We found evidence that

of an association between incident PE and PM_{2.5}, but not with distance to roadway that is proxy measure for traffic-related pollution exposure. This suggests that other, non-traffic PM_{2.5} pollution exposures (e.g., industrial emissions) may underlie the association of PE with PM_{2.5}.

A strength of our study was the ability to estimate the relative risks of idiopathic and nonidiopathic PE subtypes separately in association with PM. We found that the PM_{2.5} exposure was mainly associated with nonidiopathic PE, and the association between PM_{2.5-10} and idiopathic PE, though consistently positive, was non-significant. This is contrary to our hypothesis that air pollution would be more strongly associated with idiopathic PE cases given the general concept that air pollution induces a state of hypercoagulability and enhanced thrombosis. While precise biological pathways involved remain to be determined, our findings suggest a more complicated mechanism that long-term PM exposure may be additive or interact with preexisting health conditions (in this case, active malignancy, recent surgery or trauma) to promote PE, but may not participate in determining PE among individuals if they are not in the setting of some other risk factors of PE. Such findings may contribute to collective evidence indicating that certain characteristics, such as life stage, genetic polymorphisms, preexisting diseases and socioeconomic status, may increase the susceptibility of populations to PM-related health effects (Peled 2011; Sacks et al. 2011). Our ability to examine PE subtypes and the fact that we observed different associations by subtype may help explain the conflicting results in previous studies. If the ratio of idiopathic to nonidiopathic VTE varies between studies, heterogeneity in findings may arise. Other explanations for the differential associations observed by PE subtypes include differential random variability, biases between groups, and fewer

idiopathic cases than nonidiopathic cases. Further investigation of the effects of air pollution exposures on PE subtypes is warranted.

In addition to long-term effects of PM exposure, Dales et al. (2010) estimated increased hospital admissions for VTE and PE with daily levels of PM_{2.5} in a time-series analysis in Santiago, Chile, whereas Martinelli et al. (2012) estimated increases in emergency hospital admissions for VTE and PE associated with same day elevations of PM_{2.5-10}, but not with PM_{2.5}, in Verona, Italy. Although the underlying mechanisms of how particulate pollution may influence VTE pathogenesis (Virchow's Triad) remain to be determined, it has been suggested that short- and long-term PM exposure enhances thrombotic and coagulation potentials by increasing circulating histamine, inflammatory cytokines and platelets, plasma levels of coagulation proteins, such as procoagulant factor VIII, von Willebrand factor, and fibrinogen, and shortening prothrombin time (Baccarelli et al. 2007; Brook et al. 2010; Esmon 2003; Nemmar et al. 2006; R uckerl et al. 2006; Utell and Frampton 2002).

Several limitations of our study warrant consideration. First, we did not have information on the incidence of overall DVT or VTE, and we were not able to distinguish subtype for 22% of the PE cases. This limits our ability to examine the impact of PM on these other previously examined outcomes, as well as the power to detect statistically significant associations in our subtype analyses. Also, although self-report of PE has been well validated in this population, it is possible that some cases were not reported, leading to an underestimate of the effects of air pollution on PE. Second, our exposure estimates are predicted ambient, not personal, levels of PM. We do not have information on indoor sources, building characteristics that would influence the penetration of ambient exposures, or the amount of time each woman spent at her residence.

These factors would all lead to misclassification in exposure, and would likely bias the estimates of association towards the null. Third, since we derived PM_{2.5-10} estimates from those of PM₁₀ and PM_{2.5}, they are subjected to two sources of random error in measurements, which may lead to lower statistical power. However, Yanosky et al. (2009) demonstrated that long-term average PM_{2.5-10} was reasonably well estimated (cross-validation $R^2 = 0.63$ and 0.65 for long-term post-1999 and pre-1999 PM_{2.5-10}, respectively). Third, the GIS-based temporal spatial smoothing of the PM predictive models reduced variability relative to measured concentrations; however, this Berkson measurement error should not cause substantial bias toward the null (Gryparis et al. 2009). Findings from the current study may not be generalizable to nonwhite racial/ethnic groups, to men, to younger women and to socioeconomic deprived populations. Last but not least, it is possible that our results from this observational study are due to confounding. Adjustment for a variety of known confounding variables did not eliminate the observed associations; though there remains the possibility of residual confounding or confounding by unmeasured covariates (e.g. long-distance travel, recent prolonged inactivity), and/or by additional pollutants.

Nonetheless, our study had several strengths including the ascertainment of incident PE cases, and our ability to subtype cases based on detailed medical record review. PE is a reliable diagnosis based on evidence from objective methods such as ventilation/perfusion lung scan, thus the chance of false-positive PE cases is slim. The application of GIS-based spatiotemporal statistical models allowed us to assign exposures specific to biennially updated residential addresses for the entire follow-up period with little bias and high precision relative to other modeling approaches (Yanosky et al. 2008; 2009). Missing exposure data were minimized

because of the use of our highly specific GIS-based spatial smoothing model. The availability of updated information on individual-level covariate status (e.g., smoking) throughout the follow-up period, enabled us to take into account possible confounding effects.

Conclusion

We reported evidence of an association between exposure to PM_{2.5} and elevated risk of PE among women in the United States. By contrast, distance to road was not a risk factor for incident PE. Our findings suggest that women with underlying health conditions may be more susceptible to increased PE risk after PM exposure.

References

- Baccarelli A, Martinelli I, Pegoraro V, Melly S. 2009. Living near major traffic roads and risk of deep vein thrombosis. *Circulation* 119: 3118–3124.
- Baccarelli A, Martinelli I, Zanobetti A, Grillo P, Hou L-F, Bertazzi P a, et al. 2008. Exposure to particulate air pollution and risk of deep vein thrombosis. *Arch. Intern. Med.* 168: 920–7.
- Baccarelli A, Zanobetti A, Martinelli I, Hou L, Grillo P, Giacomini S, et al. 2007. Effects of exposure to air pollution on blood coagulation. *J. Thromb. Haemost.* 5: 252–60.
- Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, et al. 2004. Air pollution and cardiovascular disease: a statement for healthcare professionals from the Expert Panel on Population and Prevention Science of the American Heart Association. *Circulation* 109: 2655–71.
- Brook RD, Rajagopalan S, Pope CA, Brook JR, Bhatnagar A, Diez-Roux A V, et al. 2010. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation* 121: 2331–78.
- Chiuve S, Fung T, Rimm E. 2012. Alternative dietary indices both strongly predict risk of chronic disease. *J. Nutr.* 142: 1009–1018.
- CIRA (Cooperative Institute for Research in the Atmosphere). 2013. Interagency Monitoring of Protected Visual Environments (IMPROVE). Available: <http://vista.cira.colostate.edu/improve/> [accessed 8 January 2015].
- Colditz GA, Manson J, Hankinson S. 1997. The Nurses' Health Study: 20-year contribution to the understanding of health among women. *J Womens Heal.* 6:49–62.
- Dales RE, Cakmak S, Vidal CB. 2010. Air pollution and hospitalization for venous thromboembolic disease in Chile. *J. Thromb. Haemost.* 8: 669–74.
- Dockery DW, Pope CA, Xu X, Spengler JD, Ware JH, Fay ME, et al. 1993. An association between air pollution and mortality in six U.S. cities. *N. Engl. J. Med.* 329: 1753–9.
- Esmon CT. 2003. Inflammation and thrombosis. *J. Thromb. Haemost.* 1:1343–8.
- Goldhaber SZ, Visani L, Rosa M De. 1999. Acute pulmonary embolism: clinical outcomes in the International Cooperative Pulmonary Embolism Registry (ICOPER). *Lancet* 353:1386–9.
- Gryparis A, Paciorek CJ, Zeka A, Schwartz J, Coull BA. 2009. Measurement error caused by spatial misalignment in environmental epidemiology. *Biostatistics* 10: 258–74.

- Hoek G, Krishnan RM, Beelen R, Peters A, Ostro B, Brunekreef B, et al. 2013. Long-term air pollution exposure and cardio- respiratory mortality: a review. *Environ. Health* 12: 43.
- Kabrhel C, Varraso R, Goldhaber SZ, Rimm E, Camargo CA, Jr CAC. 2011. Physical inactivity and idiopathic pulmonary embolism in women : prospective study. *BMJ* 343: d3867.
- Kan H, Folsom A, Cushman M, Rose K, Rosamond W, Liao D, et al. 2011. Traffic exposure and incident venous thromboembolism in the atherosclerosis risk in communities (ARIC) study. *J. Thromb Haemost* 9: 672–678.
- Karner AA, Eisinger DS, Niemeier DA. 2010. Near-roadway air quality: synthesizing the findings from real-world data. *Environ. Sci. Technol.* 44: 5334–44.
- Kürkciyan I, Meron G. 2000. Pulmonary embolism as cause of cardiac arrest: presentation and outcome. *JAMA Intern. Med.* 160:1529–1535.
- Martinelli N, Girelli D, Cigolini D, Sandri M, Ricci G, Rocca G, et al. 2012. Access rate to the emergency department for venous thromboembolism in relationship with coarse and fine particulate matter air pollution. *PLoS One* 7: e34831.
- Miller K, Siscovick D. 2007. Long-term exposure to air pollution and incidence of cardiovascular events in women. *N. Engl. J. Med.* 356:447–458.
- Milojevic A, Wilkinson P, Armstrong B, Bhaskaran K, Smeeth L, Hajat S. 2014. Short-term effects of air pollution on a range of cardiovascular events in England and Wales: case-crossover analysis of the MINAP database, hospital admissions and mortality. *Heart* 1093–1098.
- Naess IA, Christiansen SC, Romundstad P, Cannegieter S, Rosendaal FR, Hammerstrøm J, et al. 2007. Incidence and mortality of venous thrombosis: a population-based study. *J. Thromb. Haemost.* 5: 692–9.
- National Park Service. Stack Filter Unit Network. Available:
http://vista.cira.colostate.edu/improve/data/Other/Data_nps_sfu.htm [accessed 8 January 2015].
- Nemmar A, Hoylaerts MF, Nemery B. 2006. Effects of particulate air pollution on hemostasis. *Clin. Occup. Environ. Med.* 5: 865–81.

- Office of the Surgeon General. 2008. Acting Surgeon General Issues “Call to Action to Prevent Deep Vein Thrombosis and Pulmonary Embolism.” U.S. Dep. Heal. Hum. Sci. Available: <http://www.surgeongeneral.gov/news/2008/09/pr20080915.html> [accessed 8 January 2015].
- Oger E. 2000. Incidence of Venous Thromboembolism: A Community-based Study in Western France. *Thromb. Haemost.* 83:657–660.
- Paciorek CJ, Yanosky JD, Puett RC, Laden F, Suh HH. 2009. Practical large-scale spatio-temporal modeling of particulate matter concentrations. *Ann. Appl. Stat.* 3: 370–397.
- Peled R. 2011. Air pollution exposure: Who is at high risk? *Atmos. Environ.* 45: 1781–1785.
- Piazza G, Goldhaber SZ. 2006. Acute pulmonary embolism: part I: epidemiology and diagnosis. *Circulation* 114: e28–32.
- Pope CAI, Burnett RTR, Thun MMJ, Calle EE, Krewski D, Ito K, et al. 2002. Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA* 287:1132–41.
- Previtali E, Bucciarelli P, Passamonti SM, Martinelli I. 2011. Risk factors for venous and arterial thrombosis. *Blood Transfus.* 9: 120–38.
- Puett RC, Hart JE, Yanosky JD, Paciorek C, Schwartz J, Suh H, et al. 2009. Chronic fine and coarse particulate exposure, mortality, and coronary heart disease in the Nurses’ Health Study. *Environ. Health Perspect.* 117: 1697–701.
- Rosendaal FR. 1999. Venous thrombosis: a multicausal disease. *Lancet* 353:1167–73.
- Rückerl R, Ibaldo-Mulli A, Koenig W, Schneider A, Woelke G, Cyrys J, et al. 2006. Air pollution and markers of inflammation and coagulation in patients with coronary heart disease. *Am. J. Respir. Crit. Care Med.* 173: 432–41.
- Sacks JD, Stanek LW, Luben TJ, Johns DO, Buckley BJ, Brown JS, et al. 2011. Particulate matter-induced health effects: who is susceptible? *Environ. Health Perspect.* 119: 446–54.
- Shih R a, Griffin BA, Salkowski N, Jewell A, Eibner C, Bird CE, et al. 2011. Ambient particulate matter air pollution and venous thromboembolism in the Women’s Health Initiative Hormone Therapy trials. *Environ. Health Perspect.* 119: 326–31.
- Spengler J, Koutrakis P. 1996. Health effects of acid aerosols on North American children: air pollution exposures. *Environ. Health Perspect.* 104:492–499.

- Suh HHH, Nishioka Y, Allen GAG, Washington S, Study DCF, Suh HHH, et al. 1997. The metropolitan acid aerosol characterization study: results from the summer 1994 Washington, DC field study. *Environ. Health Perspect.* 105:826–834.
- US Census Bureau. 2000. *Census 2000 CD*. U.S. Government Printing Office., Washington, DC.
- US EPA (Environmental Protection Agency). 2009. Air Quality System. Available: <http://www.epa.gov/ttn/airs/airsaqs/> [accessed 4 January 2015].
- US EPA (Environmental Protection Agency). Clean Air Status and Trends (CASTNet) networks. Available: <http://epa.gov/castnet/javaweb/index.html> [accessed 8 January 2015].
- Utell M, Frampton M. 2002. Cardiovascular effects associated with air pollution: potential mechanisms and methods of testing. *Inhal. Toxicol.* 14: 1231–1247.
- VIEWS (Visibility Information Exchange Web System). 2010. Visibility Information Exchange Homepage. Available: <http://views.cira.colostate.edu/web/> [accessed 6 January 2015].
- Yanosky J, Paciorek C, Schwartz J. 2008. Spatio-temporal modeling of chronic PM 10 exposure for the Nurses' Health Study. *Atmos. Environ. (Oxford, Engl. 1994)* 42: 4047–4062.
- Yanosky JD, Paciorek CJ, Laden F, Hart JE, Puett RC, Liao D, et al. 2014. Spatio-temporal modeling of particulate air pollution in the conterminous United States using geographic and meteorological predictors. *Environ. Health* 13: 63.
- Yanosky JD, Paciorek CJ, Suh HH. 2009. Predicting chronic fine and coarse particulate exposures using spatiotemporal models for the Northeastern and Midwestern United States. *Environ. Health Perspect.* 117: 522–9.

Table 1. Age-standardized characteristics of the Nurses' Health Study population throughout the follow-up period 1992–2008.

Characteristic	Study population
No. of women	115,745
Age, years (mean ± SD)^a	65.9 ± 8.3
White race/ethnicity (%)	96.8
BMI, kg/m² (mean ± SD)	25.6 ± 7.4
Smoking status (%)	
Never	43.8
Past	43.4
Current	12.6
Missing	0.2
Pack-years of smoking^b (mean ± SD)	24.4 ± 21.5
High-risk diet score (mean ± SD)	164.3 ± 100.4
Total caloric intake, kcal/day (mean ± SD)	1515.1 ± 724.7
Physical activity^c (%)	
<3 MET hr/week	20.6
3 to <9 MET hr/week	20.7
9 to <18 MET hr/week	18.5
18 to <27 MET hr/week	11.4
≥27 MET hr/week	19.3
Missing	9.4
Husband's education attainment (%)	
Less than high school	4.0
High school	25.8
Greater than high school	35.2
Missing	35.0
Parity (%)	
Nulliparous	5.8
1	7.3
2-3	55.0
≥4	29.9
Missing	2.1
Menopausal status^d (%)	
Premenopausal	5.1
Postmenopausal and no HRT use	22.2
Postmenopausal and past HRT use	28.8
Postmenopausal and estrogen replacement therapy use	14.0
Postmenopausal and estrogen-progesterone replacement therapy use	9.5
Missing	20.4
Comorbidities	
Hypertension (%)	46.7
Coronary heart disease (%)	3.0
Rheumatologic disease (%)	3.8
Nonaspirin nonsteroidal anti-inflammatory drug use (%)	22.1
Warfarin use (%)	2.2
Multivitamin supplement use (%)	45.2

Characteristic	Study population
Median family income in US^e \$, thousands (mean ± SD)	63.7 ± 24.7
Median house value in US^e \$, 10 thousands (mean ± SD)	17.1 ± 12.7
Predicted PM_{2.5}, µg/m³ (mean ± SD)^f	12.6 ± 3.0
Interquartile range	4.1
Predicted PM₁₀, µg/m³ (mean ± SD)^f	20.8 ± 5.8
Interquartile range	6.4
Predicted PM_{2.5-10}, µg/m³ (mean ± SD)^f	8.2 ± 4.2
Interquartile range	4.6
Distance to road (%)	
0–49 m	15.1
50–199 m	27.6
200–499 m	32.1
≥500 m	25.1

^a SD refers to standard deviation; ^b Among smokers; ^c MET refers to metabolic equivalent; ^d HRT refers to hormone-replacement therapy; ^e Estimated for census tract of residence using data from the US Census Bureau (2000); ^f for 12 months prior.

Table 2. Hazard ratios (95% confidence intervals) for pulmonary embolism (PE)^a and its subtypes with a 10 µg/m³ increase in averaged predicted PM exposure in the 1 or 12 months before diagnosis (1,725,024 total person-years), single pollutant models, 1992–2008 (n=115,745).

Pollutants	Incident PE (1,091 cases)		Idiopathic PE (323 cases)		Nonidiopathic PE (526 cases) ^b	
	Basic ^c	Multivariable adjusted ^d	Basic ^c	Multivariable adjusted ^d	Basic ^c	Multivariable adjusted ^d
1-month average						
PM _{2.5}	1.24 (1.06, 1.45)*	1.22 (1.05, 1.43)*	0.98 (0.73, 1.31)	0.98 (0.73, 1.30)	1.26 (1.01, 1.58)*	1.26 (1.00, 1.57)*
PM ₁₀	1.11 (1.01, 1.22)*	1.10 (1.00, 1.20)*	0.97 (0.82, 1.16)	0.97 (0.81, 1.15)	1.07 (0.94, 1.22)	1.06 (0.93, 1.21)
PM _{2.5-10}	1.08 (0.94, 1.23)	1.06 (0.92, 1.21)	0.96 (0.74, 1.24)	0.95 (0.73, 1.23)	0.98 (0.80, 1.20)	0.97 (0.79, 1.19)
12-month average						
PM _{2.5}	1.27 (1.02, 1.57)*	1.24 (1.00, 1.54)	1.04 (0.69, 1.54)	1.03 (0.69, 1.53)	1.22 (0.89, 1.65)	1.21 (0.88, 1.65)
PM ₁₀	1.18 (1.04, 1.32)*	1.15 (1.02, 1.30)*	1.08 (0.87, 1.35)	1.07 (0.86, 1.33)	1.10 (0.93, 1.30)	1.09 (0.92, 1.29)
PM _{2.5-10}	1.23 (1.03, 1.46)*	1.19 (1.00, 1.42)*	1.17 (0.85, 1.62)	1.16 (0.83, 1.60)	1.09 (0.84, 1.40)	1.07 (0.83, 1.38)

^a Regression results for 242 (22.2%) of the 1,091 cases had unknown PE subtype are not shown.

^b Nonidiopathic PE includes confirmed secondary PE, and PE and previous cancer.

^c Adjusted for age in months, region, year, season.

^d Adjusted for age in months, region, year, season, race/ethnicity, BMI (continuous), smoking status, pack-years, Alternate Healthy Eating Index score, total caloric intake, physical activity, husband's education attainment, parity, menopausal status, hypertension, coronary heart disease, rheumatologic disease, nonaspirin nonsteroidal anti-inflammatory drug use, warfarin use, multivitamin supplement use, median family income in census tract of residence and median house value in census tract of residence.

* P<0.05.

Table 3. Hazard ratios (95% confidence intervals) for pulmonary embolism (PE)^a and its subtypes with a 10 µg/m³ increase in averaged predicted PM exposure in the 1 or 12 months before diagnosis (1,725,024 total person-years), multipollutant models, 1992–2008 (n=115,745).

Pollutants	Incident PE (1,091 cases)		Idiopathic PE (323 cases)		Nonidiopathic PE (526 cases) ^b	
	Basic ^c	Multivariable adjusted ^d	Basic ^c	Multivariable adjusted ^d	Basic ^c	Multivariable adjusted ^d
1-month average						
PM _{2.5}	1.23 (1.05, 1.45)*	1.22 (1.04, 1.44)*	0.99 (0.74, 1.33)	0.99 (0.74, 1.33)	1.30 (1.03, 1.65)*	1.30 (1.03, 1.65)*
PM _{2.5-10}	1.02 (0.88, 1.18)	1.00 (0.87, 1.16)	0.96 (0.73, 1.26)	0.95 (0.73, 1.25)	0.91 (0.74, 1.13)	0.90 (0.73, 1.12)
12-month average						
PM _{2.5}	1.19 (0.94, 1.49)	1.17 (0.93, 1.48)	0.96 (0.63, 1.48)	0.96 (0.62, 1.47)	1.20 (0.86, 1.67)	1.20 (0.86, 1.67)
PM _{2.5-10}	1.17 (0.97, 1.41)	1.14 (0.94, 1.37)	1.18 (0.84, 1.67)	1.17 (0.83, 1.66)	1.03 (0.78, 1.36)	1.01 (0.77, 1.34)

^a Regression results for 242 (22.2%) of the 1,091 cases had unknown PE subtype are not shown.

^b Nonidiopathic PE includes confirmed secondary PE, and PE and previous cancer.

^c Adjusted for age in months, region, year, season.

^d Adjusted for age in months, region, year, season, race/ethnicity, BMI (continuous), smoking status, pack-years, Alternate Healthy Eating Index score, total caloric intake, physical activity, husband’s education attainment, parity, menopausal status, hypertension, coronary heart disease, rheumatologic disease, nonaspirin nonsteroidal anti-inflammatory drug use, warfarin use, multivitamin supplement use, median family income in census tract of residence and median house value in census tract of residence.

* P<0.05.

Table 4. Hazard ratios (95% confidence intervals) for pulmonary embolism (PE)^a and its subtypes with proximity to nearest road (1,483,727 total person-years), 1992–2008 (n=108,337).

Exposure	Person-years	Incident PE (954 cases)			Idiopathic PE (283 cases)			Nonidiopathic PE (454 cases) ^b		
		Cases	Basic ^c	Multivariable adjusted ^d	Cases	Basic ^c	Multivariable adjusted ^d	Cases	Basic ^c	Multivariable adjusted ^d
Distance to road (m)										
0–49	224,188	156	1.04 (0.85, 1.28)	1.02 (0.83, 1.25)	46	0.94 (0.65, 1.36)	0.92 (0.63, 1.33)	83	1.21 (0.91, 1.62)	1.20 (0.90, 1.61)
50–199	410,044	250	0.93 (0.78, 1.12)	0.92 (0.77, 1.10)	67	0.76 (0.55, 1.06)	0.75 (0.54, 1.04)	128	1.03 (0.80, 1.34)	1.03 (0.80, 1.34)
200–499	476,700	314	1.03 (0.86, 1.21)	1.01 (0.85, 1.20)	92	0.91 (0.67, 1.23)	0.89 (0.66, 1.21)	136	0.97 (0.75, 1.24)	0.97 (0.75, 1.25)
≥500	372,796	234	1.00 (referent)	1.00	78	1.00	1.00	107	1.00	1.00

^a Regression results for 217 (22.7%) of the 954 cases had unknown PE subtype are not shown.

^b Nonidiopathic PE includes confirmed secondary PE, and PE and previous cancer.

^c Adjusted for age in months, region, year, season.

^d Adjusted for age in months, region, year, season, race/ethnicity, BMI (continuous), smoking status, pack-years, Alternate Healthy Eating Index score, total caloric intake, physical activity, husband's education attainment, parity, menopausal status, hypertension, coronary heart disease, rheumatologic disease, nonaspirin nonsteroidal anti-inflammatory drug use, warfarin use, multivitamin supplement use, median family income in census tract of residence and median house value in census tract of residence.