

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Air Pollution Exposure during Pregnancy and Childhood Autistic Traits in Four European Population-Based Cohort Studies: The ESCAPE Project

Mònica Guxens, Akhgar Ghassabian, Tong Gong, Raquel Garcia-Esteban, Daniela Porta, Lise Giorgis-Allemand, Catarina Almqvist, Aritz Aranbarri, Rob Beelen, Chiara Badaloni, Giulia Cesaroni, Audrey de Nazelle, Marisa Estarlich, Francesco Forastiere, Joan Forns, Ulrike Gehring, Jesús Ibarluzea, Vincent W.V. Jaddoe, Michal Korek, Paul Lichtenstein, Mark J. Nieuwenhuijsen, Marisa Rebagliato, Rémy Slama, Henning Tiemeier, Frank C. Verhulst, Heather E. Volk, Göran Pershagen, Bert Brunekreef, and Jordi Sunyer

Table of Contents

Methods S1. Description of the air pollution assessment

Methods S2. Description of the autistic traits assessment

Autism-Tics, Attention deficit and hyperactivity disorders, and other Comorbidities (A-TAC) inventory (Swedish cohort)

Pervasive Developmental Problems (PDP) subscale of the Child Behavior Checklist for Toddlers (CBCL1½-5) (Dutch and Italian cohorts)

Adapted 18-item version of the Social Responsiveness Scale (SRS) (Dutch cohort)

Childhood Autism Spectrum Test (CAST) (Spanish cohort)

Table S1. Distribution of the autistic traits scales

Table S2. Power sample calculation

Table S3. Spearman correlations between air pollution levels during pregnancy and traffic indicator variables

Table S4. Minimally adjusted combined associations between air pollution exposure during pregnancy and autistic traits within the borderline/clinical range

Table S5. Fully adjusted associations between nitrogen dioxide exposure during pregnancy, potential confounding variables, and autistic traits within the borderline/clinical range across cohorts

Table S6. Fully adjusted associations between air pollution exposure during pregnancy and autistic traits as a quantitative trait across cohorts

Table S7. Fully adjusted combined associations between air pollution exposure during pregnancy and autistic traits within the borderline/clinical range, assessing the influence of a single cohort in the meta-analysis estimates

Table S8. Fully adjusted combined associations between nitrogen oxides exposure during pregnancy and autistic traits within the borderline/clinical range among cohorts with particulate matter variables available

Table S9. Fully adjusted combined associations between air pollution during pregnancy and autistic traits within the percentile 90th of each scale

Table S10. Fully adjusted combined associations between air pollution exposure during pregnancy and autistic traits within the borderline/clinical range stratified by type of evaluator of the test

Table S11. Fully adjusted combined associations between non-back-extrapolated air pollution exposure at child's birth address and autistic traits within the borderline/clinical range

Table S12. Sensitivity analyses of fully adjusted combined association between air pollution exposure during pregnancy and autistic traits within the borderline/clinical range

Table S13. Fully adjusted combined associations between air pollution exposure during pregnancy and autistic traits within the borderline/clinical range by child's sex

References