



**Cord Blood Methylmercury and Fetal Growth Outcomes  
in Baltimore Newborns: Potential Confounding and  
Effect Modification by Omega-3 Fatty Acids, Selenium,  
and Sex**

**Ellen M. Wells, Julie B. Herbstman, Yu Hong Lin, Jeffery Jarrett,  
Carl P. Verdon, Cynthia Ward, Kathleen L. Caldwell,  
Joseph R. Hibbeln, Frank R. Witter, Rolf U. Halden,  
and Lynn R. Goldman**

<http://dx.doi.org/10.1289/ehp.1408596>

**Received: 23 April 2014**

**Accepted: 24 June 2015**

**Advance Publication: 26 June 2015**

This article will be available in a 508-conformant form upon final publication. If you require a 508-conformant version before then, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Cord Blood Methylmercury and Fetal Growth Outcomes in Baltimore Newborns: Potential Confounding and Effect Modification by Omega-3 Fatty Acids, Selenium, and Sex**

Ellen M. Wells,<sup>1</sup> Julie B. Herbstman,<sup>2</sup> Yu Hong Lin,<sup>3</sup> Jeffery Jarrett,<sup>4</sup> Carl P. Verdon,<sup>4</sup> Cynthia Ward,<sup>4</sup> Kathleen L. Caldwell,<sup>4</sup> Joseph R. Hibbeln,<sup>3</sup> Frank R. Witter,<sup>5</sup> Rolf U. Halden,<sup>6</sup> and Lynn R. Goldman<sup>7</sup>

<sup>1</sup>School of Health Sciences, Purdue University, West Lafayette, Indiana, USA; <sup>2</sup>Columbia Center for Children's Environmental Health, Department of Environmental Health Sciences, Columbia University Mailman School of Public Health, New York, New York, USA; <sup>3</sup>Laboratory of Membrane Biochemistry and Biophysics, National Institute of Alcohol Abuse and Alcoholism, National Institutes of Health, Rockville, Maryland, USA; <sup>4</sup>Division of Laboratory Sciences, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA; <sup>5</sup>Department of Gynecology and Obstetrics, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA; <sup>6</sup>Center for Environmental Security, Biodesign Institute, Global Security Initiative, Arizona State University, Tempe, Arizona, USA; <sup>7</sup>Milken Institute School of Public Health, Department of Environmental and Occupational Health, George Washington University, Washington, DC, USA

**Address correspondence to** Ellen M. Wells, Purdue University, School of Health Sciences; Hampton Hall of Civil Engineering 1269; 550 Stadium Mall Drive; West Lafayette, Indiana 47907 USA. Telephone: 765-496-3535. Fax: 765-496-1377. E-mail: [wells54@purdue.edu](mailto:wells54@purdue.edu)

**Running title:** Methylmercury and fetal growth

**Acknowledgments:** We thank Benjamin Apelberg, Bruce Craig, Ruth Quinn, Norman Salem, Jr., and John Bernert for their assistance. This study received funding from the Maryland Cigarette Restitution Program Research Grant, NIEHS grant 1R01ES015445 (RUH), and the US EPA STAR Fellowship Program (EMW). The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the CDC, NIH Institutes of Health or US EPA; nor have these organizations officially endorsed this report.

**Competing financial interests:** The authors have no competing financial interests.

## Abstract

**Background:** Methylmercury (MeHg) may affect fetal growth; however, prior research often lacked assessment of mercury speciation, confounders, and interactions.

**Methods:** This cross-sectional study includes 271 singletons born in Baltimore, Maryland, 2004-5. Umbilical cord blood was analyzed for speciated mercury, serum omega-3 highly unsaturated fatty acids (n-3 HUFAs), and selenium. Multivariable linear regression models controlled for gestational age, birth weight, maternal age, parity, pre-pregnancy body mass index, smoking, hypertension, diabetes, selenium, n-3 HUFAs, and inorganic mercury (IHg).

**Results:** Geometric mean cord blood MeHg was 0.94  $\mu\text{g/L}$  (95% confidence interval (CI): 0.84, 1.07). In adjusted models for ponderal index,  $\beta\ln(\text{MeHg}) = -0.045 \text{ (g/cm}^3\text{)} \times 100$  (95% CI: -0.084, -0.005). There was no evidence of a MeHg\*sex interaction with ponderal index. Contrastingly, there was evidence of a MeHg\*n-3 HUFAs interaction with birth length (among low n-3 HUFAs,  $\beta\ln(\text{MeHg}) = 0.40 \text{ cm}$ , 95% CI: -0.02, 0.81; among high n-3 HUFAs,  $\beta\ln(\text{MeHg}) = -0.15$ , 95% CI: -0.54, 0.25;  $p$ -interaction = 0.048) and head circumference (among low n-3 HUFAs,  $\beta\ln(\text{MeHg}) = 0.01 \text{ cm}$ , 95% CI: -0.27, 0.29; among high n-3 HUFAs,  $\beta\ln(\text{MeHg}) = -0.37$ , 95% CI: -0.63, -0.10;  $p$ -interaction = 0.042). The association of MeHg with birthweight and ponderal index was affected by n-3 HUFAs, selenium, and IHg. For birthweight,  $\beta\ln(\text{MeHg})$  without these variables was -16.8 grams (-75.0, 41.3) versus -29.7 (-93.9, 34.6) with all covariates. Corresponding values for ponderal index were -0.030  $\text{(g/cm}^3\text{)} \times 100$  (-0.065, 0.005) and -0.045 (-0.084, -0.005).

**Conclusion:** We observed an association of increased MeHg with decreased ponderal index. There is evidence for interaction between MeHg and n-3 HUFAs; infants with higher MeHg and n-3 HUFAs had lower birth length and head circumference. These results should be verified with additional studies.

## Introduction

Impaired fetal growth has been associated with neonatal morbidity and mortality (Ashworth 1998; Cheung et al. 2002). Additionally, the impact of fetal growth restriction has been shown to extend to chronic diseases of adulthood (Barker et al. 2002). There are several contributors to preterm birth and restricted fetal growth (Lang et al. 1996), including environmental exposures (Triche and Hossain 2007).

Methylmercury (MeHg), best recognized as a developmental neurotoxicant, may also impair fetal growth. This was initially demonstrated in settings with high MeHg exposures (Amin-Zaki et al. 1974; Harada 1978). More recently, Ramirez et al. (2000) reported an inverse association between gestational age at birth and cord blood mercury, while Xue et al. (2007) reported that maternal hair mercury was positively associated with very preterm birth (<35 weeks), but not with all preterm births combined (<37 weeks) (Ramirez et al. 2000; Xue et al. 2007). Mercury concentrations in hair and cord blood have also been associated with reduced birthweight (Foldspang and Hansen 1990; Lee et al. 2010; Ramon et al. 2009; Sikorski et al. 1986). Consistent with this, two longitudinal studies have reported associations of cord blood mercury with reductions of attained weight in early childhood (Grandjean et al. 2003; Kim et al. 2011). A significant association of cord blood mercury with reduced birth length was reported among a Spanish cohort (Ramon et al. 2009) and an association of maternal hair mercury with decreased head circumference at 20-24 weeks' gestation among French infants with non-overweight mothers was reported ( $p < 0.06$ ) (Drouillet-Pinard et al. 2010).

In contrast, some researchers report no association of cord blood mercury and/or maternal blood mercury with gestational age (Foldspang and Hansen 1990; Grandjean et al. 2001; Lederman et al. 2008), birthweight (Ding et al. 2013; Grandjean et al. 2001; Lederman et al.

2008), length (Ding et al. 2013; Lederman et al. 2008), or head circumference (Ding et al. 2013; Gundacker et al. 2010; Lederman et al. 2008). Two studies using biomarkers for inorganic mercury (IHg) exposure did not observe any association with birth outcomes (Bashore et al. 2014; Daniels et al. 2007). Additionally, there was a significant, positive bivariate correlation of maternal hair mercury with birth length in an Austrian cohort (Gundacker et al. 2010) and a significant, positive adjusted association of cord blood mercury with head circumference in a Filipino cohort (Ramirez et al. 2000). A French cohort reported significant positive associations of maternal hair mercury with birthweight and length, as well as borderline significant positive associations with head circumference among infants with overweight mothers (Drouillet-Pinard et al. 2010).

Interpretation of this literature is challenging due to the potential influence of confounders and mercury speciation on mercury's effect on fetal growth. Seafood consumption is the main source of exposure to MeHg; however, this is also a major source for n-3 highly unsaturated fatty acids (HUFAs) and selenium, nutrients which enhance fetal growth (Choi et al. 2008). Use of total mercury (including IHg and MeHg) as a proxy for MeHg may result in overestimation of MeHg, particularly in populations with low fish consumption (Stern and Smith 2003). As the toxicokinetics and target organs of MeHg and IHg differ (Clarkson and Magos 2006), this overestimation may reduce the ability of studies to identify associations with MeHg. Meanwhile, IHg itself is toxic and it is possible that IHg exposure may result in toxicity in addition to MeHg (Chan 2011). Mercury toxicity may also be modified by sex (Gao et al. 2007; Goodrich et al. 2013). A recent review noted all of these concerns and recommended additional, careful investigations of MeHg toxicity (Karagas et al. 2012).

The current study was designed to address these concerns. Our goal is to determine the cross-sectional association between cord blood MeHg with measures of fetal growth, controlling for potential effects of co-exposure to n-3 HUFAs, selenium and IHg, while assessing potential effect modification by sex.

## **Methods**

This research includes 271 infants from the Baltimore THREE Study (Tracking Health Related to Environmental Exposures), a cross-sectional birth study in Baltimore, Maryland described in more detail in a previous publication (Apelberg et al. 2007). This research was in compliance with international human subject protection guidelines, and was approved by the Johns Hopkins School of Medicine Institutional Review Board (IRB). The IRB approved a waiver from the Health Insurance Portability and Accountability Act, and confirmed that it was not necessary to obtain informed consent because the study was determined to be of minimal risk, only required collection of specimens which would otherwise be discarded, and all study data were anonymized following data collection. Analytical services provided by the United States Centers for Disease Control and Prevention (CDC) were determined to not constitute active engagement in research.

*Study population.* This study potentially included all singleton births delivered at the Johns Hopkins Hospital in Baltimore, Maryland between November 2004 and March 2005. There were 597 singleton births (609 deliveries) during the study period. Umbilical cord blood was collected for 341 births; 300 had sufficient quantity for initial laboratory analyses that included mercury and this is the Baltimore THREE Study population. For this publication, 29 additional births were excluded due to missing data for: birth length (n=5); maternal smoking

(n=1); serum selenium (n=10); and serum fatty acids (n=12). Additionally, one infant was excluded due to an unusually small birth length. Thus, the sample size for this study is 271.

*Laboratory analyses.* Umbilical cord blood samples were collected, processed, and stored using standardized techniques (Witter et al. 2001). Samples were transferred to the CDC for metal and cotinine analyses.

Umbilical cord whole blood MeHg and IHg were determined using high performance liquid chromatography linked with inductively coupled plasma mass spectrometry (HPLC-ICP-MS) (Verdon et al. 2008). The limit of detection (LOD) was 0.48  $\mu\text{g/L}$  (MeHg) and 0.75  $\mu\text{g/L}$  (IHg). There were n=47, 17.3% (MeHg) and n=208, 76.8% (IHg) samples <LOD. A common method for addressing values <LOD is replacing them with a given value, such as LOD/2; however, this can introduce bias, particularly when the percentage of values <LOD is >10%. Therefore, we requested additional data that was <LOD but > the instrumentation limit of detection; i.e, these values were measured but too small to be declared significantly different than a blank. While these values are notably less precise than values >LOD, they are likely more accurate than other commonly used imputation methods (Succop et al. 2004). There were n=25, 9.2% (MeHg) and n=130, 48.0% (IHg) values observed, yet <LOD; these values were used as replacements. This left only n=22, 8.12% (MeHg) and n=78, 28.9% (IHg) for which there was no information. These were replaced by the lowest observed value divided by  $\sqrt{2}$ . Total mercury was assessed using ICP-MS with an LOD of 0.33  $\mu\text{g/L}$ ; seven values were < LOD.

Umbilical cord serum selenium was determined using inductively coupled plasma dynamic reaction cell mass spectrometry (ICP-DRC-MS) (Xiao 2006). The LOD for selenium was 5  $\mu\text{g/L}$ ; all values were >LOD. Umbilical cord serum cotinine was measured using liquid chromatography in conjunction with atmospheric pressure ionization mass spectrometry (LC-

API-MS) (Bernert et al. 1997). The LOD for cotinine was 0.015 ng/mL; 75 samples were <LOD. Cotinine was combined with reported smoking for analysis.

Cord serum fatty acids, including total n-3 HUFAs, were determined at the National Institute of Alcohol Abuse and Alcoholism using automated high throughput extraction based on transesterification followed by fast gas chromatography (GC) (Lin et al. 2012). Samples were analyzed for common individual n-3 HUFAs as well as total n-3 HUFAs. All samples analyzed were above the detection limit for all analytes. However, there was insufficient serum for analysis for 12 samples.

*Medical record abstraction.* Maternal smoking during pregnancy was classified as umbilical cord serum cotinine  $\geq 10$  ng/mL (Hukkanen et al. 2005) or smoking reported in the medical record. Pre-pregnancy body mass index (BMI) was calculated from height (m) and pre-pregnancy weight (kg) as  $\text{kg/m}^2$ . Missing data for pre-pregnancy weight (n=3) and height (n=7) were imputed using multivariable imputation (Wells et al. 2011). Hypertension includes chronic (pregestational) hypertension, gestational hypertension or preeclampsia, or hypertensive medication use. Diabetes includes chronic (pregestational) diabetes, gestational diabetes, or diabetic medication use. Small for gestational age was defined as <10<sup>th</sup> percentile of growth among all infants of that gestational age; large for gestational age was defined as >90<sup>th</sup> percentile, otherwise infants are considered appropriate for gestational age. Ponderal index was calculated as  $[(\text{birthweight, g})/(\text{birth length, cm})^3] \times 100$ .

*Data analysis.* Statistical analysis was performed with Stata 13.1. A  $p$ -value  $\leq 0.05$  was considered statistically significant, whereas  $p$ -values ranging from 0.05 to 0.10 were considered borderline statistically significant. The relationship of birth outcome with increasing MeHg was determined with multivariable linear regression models. Model covariates were determined by

descriptive statistics ( $p$ -value  $\leq 0.10$  in bivariate comparisons of covariates with MeHg or birth outcome) and prior literature (selenium, n-3 HUFAs) (Choi et al. 2008). Mode of delivery was considered as a covariate for head circumference; however, this was not a significant contributor to models. IHg was evaluated as a potential confounder. Various forms for covariates, including dichotomous, categorical, linear, quadratic, cubic splines were explored visually using lowess curves and statistically by comparing AIC/BIC values from models differing only by the covariate format. MeHg and IHg were lognormally distributed and therefore described using geometric means and natural-log transformed for regressions. Gestational age, birthweight, birth length, head circumference, and ponderal index were all evaluated as birth outcomes in regression models. Regression models are presented as adjusted only for gestational age; adjusted for demographics and medical history including gestational age (continuous), maternal age (quadratic), primiparity (dichotomous), prepregnancy BMI (continuous), maternal race (white/black/Asian), maternal smoking during pregnancy (dichotomous), maternal pregestational and gestational hypertension (dichotomous), maternal pregestational and gestational diabetes (dichotomous); or a fully adjusted model which also included umbilical cord serum selenium (continuous), umbilical cord serum n-3 HUFAs (continuous), and umbilical cord blood IHg (natural log transformed). We assessed model fit, heteroskedasticity, and influence. Variance inflation factors (VIF) were used to test for multicollinearity.

In a non-transformed regression equation the expected change in  $y$  given a one-unit change in  $x$  equals  $\beta(x+1)-\beta(x)$ . With a 1% change in a natural-log transformed  $x$  variable, this can be written as  $\beta(\ln(x+1\%))- \beta(\ln(x))$ ; given the properties of natural logs this can be rewritten as  $\beta[\ln((x+1\%)/x)]$ , which is equivalent to  $\beta\ln(101/100)$ . For a 100% increase (approximately a doubling), this would be  $\beta\ln(200/100)$ , which is roughly  $\beta(0.693)$ .

Potential effect modification by infant sex, selenium, or n-3 HUFAs was investigated using stratified models (i.e., subgroup of males and subgroup of females) as well as fully adjusted models computing an interaction term across the entire cohort. In these analyses these three variables were dichotomized: sex (male/female), selenium (below the median/above the median) and n-3 HUFAs (below the median/above the median). Post-hoc sensitivity analyses included determination of the individual impact of selenium, n-3 HUFAs and IHg; comparison of models using total mercury versus MeHg; and analyses with IHg and MeHg as categorical variables.

## Results

Population characteristics and cord blood analytes are presented in Table 1. Average maternal age was 25.7 years (95% confidence interval (CI): 25.0, 26.5). Mother's average pre-pregnancy BMI was 26.4 kg/m<sup>2</sup> (95% CI: 25.6, 27.3). Average n-3 HUFAs were 52.6 µg/mL (95% CI: 50.7, 54.4) and average selenium was 70.0 µg/L (95% CI: 68.5, 71.4). Geometric mean umbilical cord blood IHg was 0.13 µg/L (95% CI: 0.10, 0.17). Geometric mean umbilical cord blood MeHg was 0.94 µg/L (95% CI: 0.84, 1.07) and ranged from non-detectable levels to 15.4 µg/L. Cord blood MeHg was positively correlated with IHg, total Hg, and n-3 HUFAs, with Spearman's  $\rho$  of 0.51, 0.91, and 0.23, respectively (Supplemental Material, Table S1). MeHg was not correlated with serum selenium (Spearman's  $\rho$  = -0.02).

Cord blood MeHg increased with increasing maternal age (Table 1). This association was borderline significant (defined as  $p \leq 0.1$  but  $> 0.05$ ) based on an unadjusted linear regression model of maternal age (in years) and ln-transformed cord blood MeHg ( $\beta_{\ln(\text{MeHg})} = 0.019$ ; 95% CI: -0.0002, 0.037). Unadjusted regression model results indicated that infants born to black ( $\beta_{\ln(\text{MeHg})} = 0.50$ ; 95% CI: 0.21, 0.78) or Asian ( $\beta_{\ln(\text{MeHg})} = 1.08$ ; 95% CI: 0.59, 1.56)

mothers had significantly higher MeHg levels compared to children of white mothers). Higher MeHg was significantly related to higher n-3 HUFAs ( $\beta_{\ln(\text{MeHg})} = 0.02$ ; 95% CI: 0.01, 0.02) and IHg ( $\beta = 0.22$ ; 95% CI: 0.17, 0.28), but not selenium ( $\beta = -0.002$ ; 95% CI: -0.01, 0.01), in unadjusted regression models.

Average measures of fetal growth and development are summarized in Table 2. There were  $n=34$  (12.5%) preterm (<37 weeks gestation) and  $n=29$  (10.7%) with low birthweight (<2500 g) infants. For birth length,  $n = 4$  (1.5%) and  $n=52$  (19.2%) were classified as small and large for gestational age, respectively. For head circumference, these were  $n=14$  (5.2%) and  $n=30$  (11.1%). There were no significant differences between MeHg concentrations by preterm birth (vs. not), low birth weight (vs. not) or small for gestational age categories (vs. appropriate for gestational age) in bivariate analyses compared to  $\ln(\text{MeHg})$  using either t-tests, oneway anovas, or unadjusted regression models (data not shown). Exclusion of preterm or low birth weight infants from regression models did not substantially change our results (data not shown).

Multivariable regression model results are presented in Table 3, Figure 1 and Supplemental Material, Table S2. In fully adjusted models for the entire population (Model 3), higher MeHg was significantly associated with decreasing ponderal index, but not other outcomes. Adjusting for n-3 HUFAs, selenium, and IHg (i.e., based on Model 3 versus Model 2), increased estimated mean differences associated with a 1-unit increase in  $\ln(\text{MeHg})$  for birthweight (Model 2  $\beta = -16.8$ ; 95% CI: -75.0, 41.3; Model 3  $\beta = -29.7$ ; 95% CI: -93.9, 34.6) and ponderal index (Model 2  $\beta = -0.030$ ; 95% CI: -0.065, 0.005; Model 3  $\beta = -0.045$ ; 95% CI: -0.084, -0.005).

Supplemental Table S2 presents data on the effect of additionally including either IHg, n-3 HUFAs, or selenium as an additional covariate to Model 2. The results suggest that the

strength of the relationships between MeHg and birth outcomes were affected by inclusion of n-3 HUFAs or IHg, but not selenium. For example, in gestational age models without n-3 HUFAs or IHg  $\beta\ln(\text{MeHg})$  was 0.86 days (-0.81, 2.54); however, when also adjusting for n-3 HUFAs  $\beta\ln(\text{MeHg})$  was 0.38 (-1.33, 2.08), or when additionally adjusting for IHg  $\beta\ln(\text{MeHg})$  was 0.31 (-1.54, 2.16). When predicting birthweight using Model 2,  $\beta\ln(\text{MeHg})$  was -16.84 grams (-74.96, 41.27); however, when IHg was also included as a covariate  $\beta\ln(\text{MeHg})$  was -39.67 (-103.51, 24.17). Corresponding results in ponderal index models were: Model 2  $\beta\ln(\text{MeHg}) = -0.030$  ( $\text{g}/\text{cm}^3$ ) $\times 100$  (-0.065, 0.005); Model 2+IHg  $\beta\ln(\text{MeHg}) = -0.045$  (-0.084, -0.007).

In stratified analyses (Supplemental Material, Table S3) we observed a statistically significant association between MeHg with ponderal index among the subgroup of males only ( $\beta\ln(\text{MeHg}) = -0.063$  ( $\text{g}/\text{cm}^3$ ) $\times 100$ , 95% CI: -0.119, -0.007); between MeHg with head circumference among the subgroup of infants with cord serum n-3 HUFAs  $> 50.4$   $\mu\text{g}/\text{mL}$  ( $\beta\ln(\text{MeHg}) = -0.30$  cm, 95% CI: -0.60, -0.01); and between MeHg with length among the subgroup of infants with cord serum n-3 HUFAs  $\leq 50.4$   $\mu\text{g}/\text{mL}$  ( $\beta\ln(\text{MeHg}) = 0.46$  cm, 95% CI: 0.005, 0.91).

Results from interaction models are shown in Figure 1 and Supplemental Table S3. Our data suggest there was an interaction between MeHg and n-3 HUFAs with length ( $\beta\ln(\text{MeHg})$  among those with low n-3 HUFAs = 0.40 cm, 95% CI: -0.02, 0.81; with high n-3 HUFAs = -0.15, 95% CI: -0.54, 0.25;  $p$ -interaction = 0.048) and head circumference ( $\beta\ln(\text{MeHg})$  among those with low n-3 HUFAs = 0.01 cm, 95% CI: -0.27, 0.29; with high n-3 HUFAs = -0.37, 95% CI: -0.63, -0.10;  $p$ -interaction = 0.042). Additionally, there was a possible interaction between MeHg and sex with gestational age ( $\beta\ln(\text{MeHg})$  among males = 1.29 days, 95% CI: -1.00, 3.58; with females = -1.59, 95% CI: -3.99, 0.81;  $p$ -interaction = 0.062) as well as between MeHg and

selenium with birthweight ( $\beta\ln(\text{MeHg})$  among infants with low Se = -77.8 g, 95% CI: -162.9, 7.4; with high Se = 25.8, 95% CI: -58.2, 109.7;  $p$ -interaction = 0.067).

There were statistically significant associations of selenium with increasing gestational age and increasing birthweight. N-3 HUFAs were significantly associated with decreasing head circumference and infant length. Increasing IHg was related to statistically significant increases in birthweight and ponderal index (data not shown).

The MeHg-ponderal index association in models using categorical variables for either MeHg or IHg was attenuated, but similar, to models that used continuous variables for both MeHg and IHg (data not shown). Similarly, an association for total mercury with ponderal index was attenuated, but similar to the association of MeHg with ponderal index (data not shown).

## Discussion

We identified a statistically significant decrease in ponderal index associated with higher cord blood MeHg concentration among newborns in models controlling for selenium, n-3 HUFAs and IHg. Additionally, our results provide evidence of an interaction of MeHg and n-3 HUFAs with birth length and head circumference.

Our goal was to evaluate the relationship of mercury with prenatal growth accounting for mercury speciation, negative confounding, and potential effect modification by sex. Although we did not observe substantial differences between using total versus MeHg (for ponderal index:  $\beta\ln(\text{totalHg}) = -0.044 \text{ (g/cm}^3\text{)} \times 100$ , 95% CI: -0.099, 0.011; whereas  $\beta\ln(\text{MeHg}) = -0.045$  95% CI: -0.084, -0.005), or inclusion of confounders, even these small changes could plausibly explain at least part of the mixed results in existing literature.

An Icelandic cohort evaluated fish consumption and ponderal index; however, they did not see a significant association with either fish or fish oil consumption (Thorsdottir et al. 2004).

The estimated decrease in ponderal index associated with MeHg in our study population suggests that MeHg exposure during gestation may result in thinner newborns. Although the magnitude of the estimated difference in ponderal index was small, this finding still suggests a potential public health risk among more highly MeHg-exposed infants as well as infants exposed to other risk factors for reduced fetal growth. Barker and colleagues first demonstrated how alterations in early development could lead to chronic disease among adults (Barker et al. 2002), a concept now understood as the developmental origins of adult disease. Reduced ponderal index is one factor that has been associated with later life health implications, particularly cardiometabolic outcomes (Eriksson et al. 2001; Larnkjaer et al. 2011).

We did not observe an association of mercury and gestational age, similar to some (Foldspang and Hansen 1990; Grandjean et al. 2001; Lederman et al. 2008), but not all (Ramirez et al. 2000; Xue et al. 2007) reports. Part of this may be explained by population differences or variations in adjustment for confounders. Some studies report increased fish consumption is related to increased gestational age (Guldner et al. 2007; Heppe et al. 2011; Olsen and Secher 2002). Two studies measuring both cord blood mercury and n-3 HUFAs report a positive association of fatty acids, but no association of mercury with longer gestation (Grandjean et al. 2001; Lucas et al. 2004). Our results additionally suggest selenium has a small, but significant positive association with gestational age ( $\beta = 0.31$  days, 95% CI: 0.18, 0.45) and birthweight ( $\beta = 6.72$  grams, 95% CI: 1.71, 11.73).

Much existing literature reports an inverse association (Foldspang and Hansen 1990; Lee et al. 2010; Ramon et al. 2009; Sikorski et al. 1986) or no association (Ding et al. 2013; Grandjean et al. 2001; Lederman et al. 2008; Lucas et al. 2004) of cord blood, maternal blood, or maternal hair mercury with birthweight. Drouillet-Pinard and colleagues report increased

birthweight with higher maternal hair mercury among infants with overweight mothers; however, this did not persist when adjusting for fish consumption (Drouillet-Pinard et al. 2010). It is possible that this may be explained by collinearity of n-3 HUFAs and mercury in seafood; our data suggest n-3 HUFAs may confound mercury's association with birthweight (Table 3, Supplemental Material Table S2). This is also supported by studies demonstrating that consumption of different kinds of fish (which likely vary in mercury and n-3 HUFA concentrations) have different associations with birthweight (Guldner et al. 2007; Heppe et al. 2011; Thorsdottir et al. 2004).

Ponderal index and birthweight are correlated in our population (Spearman's  $\rho=0.46$ ), however, babies with higher ponderal index may have lower or average birthweight (if babies have shorter length) and babies with lower ponderal index may have average or higher birthweight (if babies are longer). At this point it would be premature to make clinical conclusions on the basis of these data. Our findings on ponderal index need to be replicated in other studies.

Similar to our study, three cohorts did not report any significant association of cord blood or maternal hair mercury with birth length (Ding et al. 2013; Lederman et al. 2008; Sikorski et al. 1986); there was an increased risk of small for gestational age for length within a Spanish birth cohort with increased cord blood mercury (Ramon et al. 2009); and others found a significant relationship of increasing maternal hair mercury with increased length (Drouillet-Pinard et al. 2010; Gundacker et al. 2010). However, the result from Gundacker and colleagues was unadjusted, and the positive association in Drouillet-Pinard's finding was limited to overweight mothers and did not persist after additionally controlling for seafood consumption (Drouillet-Pinard et al. 2010; Gundacker et al. 2010). Both negative (Halldorsson et al. 2007; Thorsdottir et

al. 2004) and positive (Thorsdottir et al. 2004) associations between fish consumption and length at birth have been reported in prior studies.

Several studies reported no association between cord blood or maternal hair mercury with head circumference (Ding et al. 2013; Gundacker et al. 2010; Lederman et al. 2008; Sikorski et al. 1986). Ramirez and colleagues report a statistically significant increase in head circumference with cord blood mercury exposure; however, it is unclear what confounders were included in their model (Ramirez et al. 2000). Drouillet-Pinard and colleagues report a borderline significant increase in head circumference with increasing maternal hair mercury among overweight mothers; however, this did not persist after adjustment for fish consumption (Drouillet-Pinard et al. 2010). Contrastingly, decreases in head circumference with increasing hair mercury were observed from the subset of non-overweight mothers among this French cohort and studies which evaluated fish consumption instead of mercury biomarkers (Drouillet-Pinard et al. 2010; Halldorsson et al. 2007; Heppe et al. 2011; Thorsdottir et al. 2004). Although not statistically significant, we also observed a decrease in head circumference with increasing MeHg.

We observed a negative association of cord serum n-3 HUFAs with head circumference, length and birthweight in both our main models as well as our interaction models; this was surprising given the extensive evidence of the beneficial effects of n-3 HUFAs on fetal development (Imhoff-Kunsch et al. 2012). The reason for this is unclear. It is possible this result may have been influenced by the relatively low concentrations of n-3 HUFAs found in this population (Lin et al. 2012) or the cross-sectional nature of this analysis (Drouillet et al. 2009). Although n-3 HUFAs and MeHg were not highly correlated (Spearman's  $\rho < 0.3$ ), it is still possible that there may have been some influence of collinearity in our analyses. However, our

finding of longer gestation is consistent with findings of reduced risk of early preterm delivery with fatty acid supplementation (Imhoff-Kunsch et al. 2012).

In stratified analyses we found MeHg-ponderal index and birthweight associations among males, but not females. It is possible that we had insufficient power to detect a relationship among females. However, our results are consistent with previous work. There is evidence that mercury toxicokinetics vary by sex (Ekstrand et al. 2010; Lie et al. 1982; Nielsen 1992). Additionally, there are reports of increased neurologic impacts of mercury exposure among males compared to females (Cauli et al. 2013; Gao et al. 2007; Philibert et al. 2008). We are not aware of previous work looking at sex differences of mercury's relationship with fetal growth.

Our ability to control for n-3 HUFAs, selenium, and IHg is a strength of this study. The importance of n-3 HUFAs and selenium as negative confounders is that without taking these opposite effects into consideration, one may underestimate mercury's toxicity (Choi et al. 2008; Karagas et al. 2012; Ou et al. 2014); indeed, our estimates for mercury's effect on birthweight and ponderal index appear to be strengthened after inclusion of these variables.

In comparison, while it is known that toxicokinetics for MeHg and IHg differ, little is understood regarding this potential impact on fetal growth. Limited prior research with IHg did not identify any association of IHg with birth outcomes (Bashore et al. 2014; Daniels et al. 2007). In contrast, we found that associations of IHg with birth outcomes were generally the inverse of MeHg. These differences may be due to the fact that we included both IHg and MeHg as covariates; this raises the possibility that IHg may act as a negative confounder with respect to MeHg exposure. Umbilical cord blood IHg could be the result of placental transport of IHg from the mother to the fetus, or endogenously formed from MeHg demethylation (Carneiro et al. 2014; Ou et al. 2014; Stern and Smith 2003). Therefore, one possible explanation for our results

would be that higher concentrations of IHg in umbilical cord serum may be indicative of rapid MeHg metabolism.

Although we had precise speciated mercury measurements, many values were <LOD; therefore, we included observed values <LOD in analyses. While more precise than a single replacement value, these are still less precise than higher concentrations; however, the fact that we observed similar results when modeling mercury as quantiles suggests this was not a major limitation of our study.

Average cord blood mercury concentrations among this study population (either MeHg or total mercury) are lower than concentrations reported in a recent review on low-level exposures (Karagas et al. 2012), with the exception of an Austrian birth cohort (Gundacker et al. 2010). It is possible that in our population these associations were only detectable among the most highly exposed. Another possibility is the association of MeHg with ponderal index or other birth outcomes is nonlinear. Future analyses should consider the nonlinear possibility in their analyses.

This study provides supporting evidence for a relationship between mercury and fetal growth restriction. Our data also suggest that n-3 HUFAs, selenium, and sex may modify these relationships; and that n-3 HUFAs and IHg may confound these relationships. Further consideration of confounding and effect modification should be applied to future work.

## References

- Amin-Zaki L, Elhassani S, Majeed MA, Clarkson TW, Doherty RA, Greenwood M. 1974. Intra-uterine methylmercury poisoning in Iraq. *Pediatrics* 54:587-595.
- Apelberg BJ, Witter FR, Herbstman JB, Calafat AM, Halden RU, Needham LL, et al. 2007. Cord serum concentrations of perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in relation to weight and size at birth. *Environ Health Perspect* 115:1670-1676.
- Ashworth A. 1998. Effects of intrauterine growth retardation on mortality and morbidity in infants and young children. *Eur J Clin Nutr* 52 Suppl 1:S34-41; discussion S41-32.
- Barker DJ, Eriksson JG, Forsen T, Osmond C. 2002. Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol* 31:1235-1239.
- Bashore CJ, Geer LA, He X, Puett R, Parsons PJ, Palmer CD, et al. 2014. Maternal mercury exposure, season of conception and adverse birth outcomes in an urban immigrant community in Brooklyn, New York, U.S.A. *Int J Environ Res Public Health* 11:8414-8442.
- Bernert JT, Jr., Turner WE, Pirkle JL, Sosnoff CS, Akins JR, Waldrep MK, et al. 1997. Development and validation of sensitive method for determination of serum cotinine in smokers and nonsmokers by liquid chromatography/atmospheric pressure ionization tandem mass spectrometry. *Clin Chem* 43:2281-2291.
- Carneiro MFH, Grotto D, Barbosa F. 2014. Inorganic and Methylmercury Levels in Plasma Are Differentially Associated with Age, Gender, and Oxidative Stress Markers in a Population Exposed to Mercury through Fish Consumption. *J Toxicol Env Heal A* 77:69-79.

- Cauli O, Piedrafita B, Llansola M, Felipo V. 2013. Gender differential effects of developmental exposure to methyl-mercury, polychlorinated biphenyls 126 or 153, or its combinations on motor activity and coordination. *Toxicology* 311:61-68.
- Chan TYK. 2011. Inorganic mercury poisoning associated with skin-lightening cosmetic products. *Clin Toxicol* 49:886-891.
- Cheung YB, Yip PS, Karlberg JP. 2002. Size at birth and neonatal and postneonatal mortality. *Acta Paediatr* 91:447-452.
- Choi AL, Cordier S, Weihe P, Grandjean P. 2008. Negative confounding in the evaluation of toxicity: the case of methylmercury in fish and seafood. *Crit Rev Toxicol* 38:877-893.
- Clarkson TW, Magos L. 2006. The toxicology of mercury and its chemical compounds. *Crit Rev Toxicol* 36:609-662.
- Daniels JL, Rowland AS, Longnecker MP, Crawford P, Golding J. 2007. Maternal dental history, child's birth outcome and early cognitive development. *Paediatr Perinat Epidemiol* 21:448-457.
- Ding G, Cui C, Chen L, Gao Y, Zhou Y, Shi R, et al. 2013. Prenatal low-level mercury exposure and neonatal anthropometry in rural northern China. *Chemosphere* 92:1085-1089.
- Drouillet-Pinard P, Huel G, Slama R, Forhan A, Sahuquillo J, Goua V, et al. 2010. Prenatal mercury contamination: relationship with maternal seafood consumption during pregnancy and fetal growth in the 'EDEN mother-child' cohort. *Br J Nutr* 104:1096-1100.
- Drouillet P, Forhan A, De Lauzon-Guillain B, Thiebaugeorges O, Goua V, Magnin G, et al. 2009. Maternal fatty acid intake and fetal growth: evidence for an association in overweight women. The 'EDEN mother-child' cohort (study of pre- and early postnatal determinants of the child's development and health). *Br J Nutr* 101:583-591.

- Ekstrand J, Nielsen JB, Havarinasab S, Zalups RK, Soderkvist P, Hultman P. 2010. Mercury toxicokinetics--dependency on strain and gender. *Toxicol Appl Pharmacol* 243:283-291.
- Eriksson JG, Forsen T, Tuomilehto J, Osmond C, Barker DJ. 2001. Early growth and coronary heart disease in later life: longitudinal study. *BMJ* 322:949-953.
- Foldspang A, Hansen JC. 1990. Dietary intake of methylmercury as a correlate of gestational length and birth weight among newborns in Greenland. *Am J Epidemiol* 132:310-317.
- Gao Y, Yan CH, Tian Y, Wang Y, Xie HF, Zhou X, et al. 2007. Prenatal exposure to mercury and neurobehavioral development of neonates in Zhoushan City, China. *Environ Res* 105:390-399.
- Goodrich JM, Basu N, Franzblau A, Dolinoy DC. 2013. Mercury biomarkers and DNA methylation among Michigan dental professionals. *Environ Mol Mutagen* 54:195-203.
- Grandjean P, Bjerve KS, Weihe P, Steuerwald U. 2001. Birthweight in a fishing community: significance of essential fatty acids and marine food contaminants. *Int J Epidemiol* 30:1272-1278.
- Grandjean P, Budtz-Jorgensen E, Steuerwald U, Heinzow B, Needham LL, Jorgensen PJ, et al. 2003. Attenuated growth of breast-fed children exposed to increased concentrations of methylmercury and polychlorinated biphenyls. *FASEB J* 17:699-701.
- Guldner L, Monfort C, Rouget F, Garlantezec R, Cordier S. 2007. Maternal fish and shellfish intake and pregnancy outcomes: a prospective cohort study in Brittany, France. *Environ Health* 6:33.
- Gundacker C, Frohlich S, Graf-Rohrmeister K, Eibenberger B, Jessenig V, Gicic D, et al. 2010. Perinatal lead and mercury exposure in Austria. *Sci Total Environ* 408:5744-5749.

- Halldorsson TI, Meltzer HM, Thorsdottir I, Knudsen V, Olsen SF. 2007. Is high consumption of fatty fish during pregnancy a risk factor for fetal growth retardation? A study of 44,824 Danish pregnant women. *Am J Epidemiol* 166:687-696.
- Harada M. 1978. Congenital Minamata disease: intrauterine methylmercury poisoning. *Teratology* 18:285-288.
- Heppe DH, Steegers EA, Timmermans S, Breeijen H, Tiemeier H, Hofman A, et al. 2011. Maternal fish consumption, fetal growth and the risks of neonatal complications: the Generation R Study. *Br J Nutr* 105:938-949.
- Hukkanen J, Jacob P, 3rd, Benowitz NL. 2005. Metabolism and disposition kinetics of nicotine. *Pharmacol Rev* 57:79-115.
- Imhoff-Kunsch B, Briggs V, Goldenberg T, Ramakrishnan U. 2012. Effect of n-3 long-chain polyunsaturated fatty acid intake during pregnancy on maternal, infant, and child health outcomes: a systematic review. *Paediatr Perinat Epidemiol* 26 Suppl 1:91-107.
- Karagas MR, Choi AL, Oken E, Horvat M, Schoeny R, Kamai E, et al. 2012. Evidence on the human health effects of low-level methylmercury exposure. *Environ Health Perspect* 120:799-806.
- Kim BM, Lee BE, Hong YC, Park H, Ha M, Kim YJ, et al. 2011. Mercury levels in maternal and cord blood and attained weight through the 24 months of life. *Sci Total Environ* 410-411:26-33.
- Lang JM, Lieberman E, Cohen A. 1996. A comparison of risk factors for preterm labor and term small-for-gestational-age birth. *Epidemiology* 7:369-376.
- Larnkjaer A, Ingstrup HK, Schack-Nielsen L, Molgaard C, Michaelsen KF. 2011. Thin newborns are more insulin resistant at 10 years of age. *Acta Paediatr* 100:511-514.

- Lederman SA, Jones RL, Caldwell KL, Rauh V, Sheets SE, Tang D, et al. 2008. Relation between cord blood mercury levels and early child development in a World Trade Center cohort. *Environ Health Perspect* 116:1085-1091.
- Lee BE, Hong YC, Park H, Ha M, Koo BS, Chang N, et al. 2010. Interaction between GSTM1/GSTT1 polymorphism and blood mercury on birth weight. *Environ Health Perspect* 118:437-443.
- Lie A, Gundersen N, Korsgaard KJ. 1982. Mercury in urine.--Sex, age and geographic differences in a reference population. *Scand J Work Environ Health* 8:129-133.
- Lin YH, Salem N, Jr., Wells EM, Zhou W, Loewke JD, Brown JA, et al. 2012. Automated high-throughput fatty acid analysis of umbilical cord serum and application to an epidemiological study. *Lipids* 47:527-539.
- Lucas M, Dewailly E, Muckle G, Ayotte P, Bruneau S, Gingras S, et al. 2004. Gestational age and birth weight in relation to n-3 fatty acids among Inuit (Canada). *Lipids* 39:617-626.
- Nielsen JB. 1992. Toxicokinetics of mercuric chloride and methylmercuric chloride in mice. *J Toxicol Environ Health* 37:85-122.
- Olsen SF, Secher NJ. 2002. Low consumption of seafood in early pregnancy as a risk factor for preterm delivery: prospective cohort study. *BMJ* 324:447.
- Ou L, Chen L, Chen C, Yang T, Wang H, Tong Y, et al. 2014. Associations of methylmercury and inorganic mercury between human cord blood and maternal blood: a meta-analysis and its application. *Environmental pollution* 191:25-30.
- Philibert A, Bouchard M, Mergler D. 2008. Neuropsychiatric symptoms, omega-3, and mercury exposure in freshwater fish-eaters. *Archives of environmental & occupational health* 63:143-153.

- Ramirez GB, Cruz MC, Pagulayan O, Ostrea E, Dalisay C. 2000. The Tagum study I: analysis and clinical correlates of mercury in maternal and cord blood, breast milk, meconium, and infants' hair. *Pediatrics* 106:774-781.
- Ramon R, Ballester F, Aguinagalde X, Amurrio A, Vioque J, Lacasana M, et al. 2009. Fish consumption during pregnancy, prenatal mercury exposure, and anthropometric measures at birth in a prospective mother-infant cohort study in Spain. *Am J Clin Nutr* 90:1047-1055.
- Sikorski R, Paszkowski T, Szprengier-Juszkiewicz T. 1986. Mercury in neonatal scalp hair. *Sci Total Environ* 57:105-110.
- Stern AH, Smith AE. 2003. An assessment of the cord blood:maternal blood methylmercury ratio: implications for risk assessment. *Environ Health Perspect* 111:1465-1470.
- Succop PA, Clark S, Chen M, Galke W. 2004. Imputation of data values that are less than a detection limit. *J Occup Environ Hyg* 1:436-441.
- Thorsdottir I, Birgisdottir BE, Halldorsdottir S, Geirsson RT. 2004. Association of fish and fish liver oil intake in pregnancy with infant size at birth among women of normal weight before pregnancy in a fishing community. *Am J Epidemiol* 160:460-465.
- Triche EW, Hossain N. 2007. Environmental factors implicated in the causation of adverse pregnancy outcome. *Semin Perinatol* 31:240-242.
- Verdon CP, Caldwell KL, Jones RL. 2008. Blood mercury speciation performed by HPLC-ICP-DRC-MS. Centers for Disease Control and Prevention: Division of Laboratory Sciences Method 3010.2:1-97.

- Wells EM, Jarrett JM, Lin YH, Caldwell KL, Hibbeln JR, Apelberg BJ, et al. 2011. Body burdens of mercury, lead, selenium and copper among Baltimore newborns. *Environ Res* 111:411-417.
- Witter FR, Ten Broeck J, Fox HE. 2001. A new device for safer collection of postpartum cord blood. *Int J Gynaecol Obstet* 72:259-260.
- Xiao G. 2006. Zinc, copper and selenium in serum by ICP-DRC-MS (ICPDRCMS-3006.1). Centers for Disease Control and Prevention.
- Xue F, Holzman C, Rahbar MH, Trosko K, Fischer L. 2007. Maternal fish consumption, mercury levels, and risk of preterm delivery. *Environ Health Perspect* 115:42-47.

**Table 1: Population characteristics**

Characteristic	Category	N (%) <sup>a</sup>	Cord blood MeHg, µg/L <sup>b</sup>
Entire population		271 (100.0)	0.94 (0.84, 1.07)
Infant sex	Female	121 (44.7)	0.90 (0.74, 1.08)
	Male	150 (55.4)	0.98 (0.84, 1.16)
Maternal age	< 20 years	55 (20.3)	0.70 (0.54, 0.92)*
	20-29 years	135 (49.8)	0.96 (0.83, 1.11)*
	> 29 years	81 (29.9)	1.12 (0.86, 1.47)*
Maternal race	White/Caucasian	59 (21.8)	0.61 (0.44, 0.85)*
	African American	190 (70.1)	1.00 (0.89, 1.13)*
	Asian	22 (8.1)	1.79 (1.03, 3.11)*
Maternal smoking during pregnancy	None/passive	225 (83.0)	0.93 (0.81, 1.06)
	Active smoker	46 (17.0)	1.02 (0.78, 1.34)
Maternal parity	Primiparous	116 (42.8)	0.93 (0.77, 1.13)
	Not primiparous	155 (57.2)	0.95 (0.81, 1.12)
Maternal prepregnancy BMI	< 25 kg/m <sup>2</sup>	142 (52.4)	0.98 (0.82, 1.17)
	25-29.9 kg/m <sup>2</sup>	64 (23.6)	0.85 (0.66, 1.10)
	≥ 30 kg/m <sup>2</sup>	65 (24.0)	0.96 (0.76, 1.21)
Hypertension <sup>c</sup>	No	244 (90.0)	0.97 (0.85, 1.10)
	Yes	27 (10.0)	0.76 (0.48, 1.20)
Diabetes <sup>c</sup>	No	252 (93.0)	0.95 (0.83, 1.07)
	Yes	19 (7.0)	0.91 (0.53, 1.58)*
Cord serum n-3 HUFAs (quartiles)	22.70 – 42.00 µg/mL	68 (25.1)	0.72 (0.56, 0.92)*
	42.01 – 50.40 µg/mL	68 (25.1)	0.89 (0.71, 1.10)*
	50.41 – 60.25 µg/mL	68 (25.1)	0.96 (0.75, 1.24)*
	60.26 – 126.95 µg/mL	67 (24.7)	1.30 (1.01, 1.67)*
Cord serum selenium (quartiles)	42 – 62 µg/L	73 (26.9)	0.90 (0.71, 1.15)
	63 – 69 µg/L	67 (24.7)	1.12 (0.89, 1.41)
	70 – 78 µg/L	67 (24.7)	0.85 (0.66, 1.09)
	79 – 114 µg/L	74 (23.6)	0.93 (0.72, 1.20)
Cord blood IHg (tertiles)	ND – 0.08 µg/L	91 (33.6)	0.54 (0.44, 0.65)*
	0.09 – 0.57 µg/L	92 (34.0)	0.91 (0.75, 1.10)*
	0.58 – 2.17 µg/L	88 (32.5)	1.76 (1.48, 2.10)*

MeHg = methylmercury; G. mean = geometric mean; CI = confidence interval; BMI = body mass index; HUFAs = highly unsaturated fatty acids; IHg=inorganic mercury; ND = not detected.

a. Percentages may not sum to 100 due to rounding.

b. Values are geometric mean (95% CI).

c. Includes pregestational and gestational.

\*  $p \leq 0.05$ , oneway ANOVA.

**Table 2: Mean (95% CI) for birth outcomes stratified by MeHg quartile**

Range of MeHg, µg/L	Quartiles of cord blood MeHg				Entire population
	ND – 0.61 (Q1)	0.62 – 1.04 (Q2)	1.05 – 1.84 (Q3)	1.85 – 15.4 (Q4)	ND-15.4 µg/L
Gestational age, days	270.0 (266.3, 273.7)	272.1 (268.9, 275.3)	273.2 (270.1, 276.4)	273.7 (270.7, 276.8)	272.3 (270.6, 273.9)
Birthweight, grams	3171 (3033, 3308)	3215 (3061, 3370)	3275 (3141, 3410)	3186 (3038, 3333)	3212 (3140, 3284)
Birth length, cm	49.7 (49.0, 50.3)	49.8 (49.1, 50.6)	50.3 (49.7, 50.8)	50.3 (49.7, 51.0)	50.0 (49.7, 50.3)
Head circumference, cm	33.6 (33.1, 34.1)	33.7 (33.2, 34.1)	33.3 (32.9, 33.7)	33.5 (33.0, 33.9)	33.5 (33.3, 33.7)
Ponderal index, ((g/cm <sup>3</sup> )×100)	2.57 (2.50, 2.64)	2.58 (2.50, 2.66)	2.57 (2.50, 2.63)	2.48 (2.41, 2.55)	2.55 (2.51, 2.58)

MeHg = methylmercury; CI = confidence interval; ND = not detected; Q=quartile. N=271.

**Table 3: Estimated mean difference in birth outcomes with an increase in cord blood MeHg**

Outcome	Model	$\beta$ for ln(MeHg)
Gestational age, days	Model 1 <sup>a</sup>	1.23 (-0.38, 2.84)
	Model 2 <sup>b</sup>	0.86 (-0.81, 2.54)
	Model 3 <sup>c</sup>	-0.07 (-1.87, 1.73)
Birthweight, grams	Model 1 <sup>a</sup>	-19.58 (-76.14, 36.97)
	Model 2 <sup>b</sup>	-16.84 (-74.96, 41.27)
	Model 3 <sup>c</sup>	-29.65 (-93.89, 34.60)
Birth length, cm	Model 1 <sup>a</sup>	0.09 (-0.18, 0.35)
	Model 2 <sup>b</sup>	0.09 (-0.18, 0.37)
	Model 3 <sup>c</sup>	0.14 (-0.17, 0.45)
Head circumference, cm	Model 1 <sup>a</sup>	-0.13 (-0.31, 0.06)
	Model 2 <sup>b</sup>	-0.14 (-0.33, 0.04)
	Model 3 <sup>c</sup>	-0.16 (-0.37, 0.04)
Ponderal index, (g/cm <sup>3</sup> ) $\times$ 100	Model 1 <sup>a</sup>	-0.032 (-0.065, 0.002) <sup>*</sup>
	Model 2 <sup>b</sup>	-0.029 (-0.065, 0.005) <sup>*</sup>
	Model 3 <sup>c</sup>	-0.045 (-0.084, -0.005) <sup>#</sup>

MeHg = methylmercury; IHg = inorganic mercury; HUFAs = highly unsaturated fatty acids; CI = confidence interval

- Model 1 (gestational age adjustments) is adjusted for gestational age (except for gestational age outcome).
  - Model 2 (demographic/medical history adjustments) additionally adjusts for infant sex (entire cohort only), gestational age (except in gestational age model), maternal age (quadratic), primiparity, prepregnancy body mass index, maternal race, maternal smoking, maternal pregestational and gestational hypertension, and maternal pregestational and gestational diabetes.
  - Model 3 (all adjustments) additionally adjusts for cord serum selenium, cord serum n3-HUFAs, and cord blood IHg.
- <sup>\*</sup>*p* <0.10; <sup>#</sup>*p*  $\leq$ 0.05; N=271.

## Figure Legend

**Figure 1.** Differences and 95% confidence intervals for birth outcomes related to a one-unit increase in  $\ln(\text{MeHg})$ , i.e.,  $\beta\ln(\text{MeHg})$ . The outcomes include gestational age in days (Panel A); birthweight in grams (Panel B); birth length in centimeters (Panel C); head circumference in centimeters (Panel D); and ponderal index in  $(\text{g}/\text{cm}^3)\times 100$  (Panel E). Within each panel, from left to right, the lines indicate results from the entire cohort (N=271) and specific subgroups including females (N=121); males (N=150); serum n-3 HUFAS  $\leq 50.4 \mu\text{g}/\text{mL}$ , or “low n-3s” (N = 136); n-3 HUFAS  $>50.4 \mu\text{g}/\text{mL}$ , or “high n-3s” (N=135); serum selenium  $< 70 \mu\text{g}/\text{L}$ , or “low Se” (N=140); and selenium  $\geq 70 \mu\text{g}/\text{L}$ , or “high Se” (N=131). *P*-values of interaction terms ( $\ln(\text{MeHg})\text{*sex}$ ,  $\ln(\text{MeHg})\text{*n-3 HUFAs}$ , and  $\ln(\text{MeHg})\text{*selenium}$ , respectively) are indicated with “p-int”.

Figure 1.

