



ENVIRONMENTAL HEALTH PERSPECTIVES

<http://www.ehponline.org>

Evidence from Toxicology: The Most Essential Science for Prevention

Daniele Mandrioli and Ellen Kovner Silbergeld

<http://dx.doi.org/10.1289/ehp.1509880>

Received: 25 February 2015

Accepted: 12 June 2015

Advance Publication: 19 June 2015

This article will be available in a 508-conformant form upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.



Evidence from Toxicology: The Most Essential Science for Prevention

Daniele Mandrioli^{1,2} and Ellen Kovner Silbergeld¹

¹Department of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA; ²Cesare Maltoni Cancer Research Center, Ramazzini Institute, Bologna, Italy

Address correspondence to Daniele Mandrioli, Department of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health, 615 North Wolfe Street E6644, Baltimore, MD 21205 USA. Telephone: 410 955 8678. Fax: 410 955 9334. E-mail: dmandri1@jhu.edu

Running title: Toxicology and prevention

Acknowledgments: The authors thank continuing conversations with principals in the Cochrane Collaboration, especially Kay Dickersin and Roberta Scherer of the US Cochrane Center, Lisa Bero and Elizabeth Waters, Merel Ritskes-Hoitinga and other members of the working group on animal testing and other attendees at a working group held during the 23rd Cochrane Colloquium; Lori Rosman and Ana Navas Acien of Johns Hopkins; and our colleagues in the work of developing methods for evidence based toxicology Tracey Woodruff of UCSF, Khristina Thayer of NIEHS and Vincent Coglianò of EPA.

Competing financial interests: There were no funds received to support the writing or production of this paper. The authors declare no competing financial interests

Abstract

Background: The most essential goal of medicine and public health is to prevent harm [*primum non nocere*]. This goal is only fully achieved with primary prevention, which requires us to identify and prevent harms prior to human exposure through research and testing that does not involve human subjects. For that reason, public health policies place considerable reliance on nonhuman toxicological studies. But toxicology as a field has often not produced efficient and timely evidence for decision making in public health. In response to this, the US National Research Council called for the adoption of evidence-based methods and systematic reviews in regulatory decision making. EPA, FDA and the European Food Safety Agency have recently endorsed these methods in their assessments of safety and risk.

Objectives: In this article we summarize challenges and problems in current practices in toxicology as applied to decision making. We compare these practices with the principles and methods utilized in evidence based medicine and healthcare, with emphasis on the record of the Cochrane Collaboration.

Discussion: We propose a stepwise strategy to support the development, validation, and application of evidence based toxicology (EBT). We discuss current progresses in this field produced by the Office of Health Assessment and Translation of the National Toxicology Program (OHAT) and Navigation Guide works. We propose that adherence to the Cochrane principles is a fundamental prerequisite for the development and implementation of EBT.

Conclusion: The adoption of evidence based principles and methods will enhance the validity, transparency, efficiency and acceptance of toxicological evidence, with benefits in terms of reducing delays and costs for all stakeholders (researchers, consumers, regulators, industry).

Introduction

The most essential goal of medicine and public health is to prevent harm [in the words of Hippocrates, *primum non nocere*]. This goal is only fully achieved with primary prevention, which requires us to identify harms prior to human exposure. Toxicology, almost always involving nonhuman subjects, is the main source of such information. “Ethical principles of human subjects research have developed in response to several examples of morally reprehensible research involving humans over the past 70 years (Josefson 2001; Katz et al. 2006), prohibit the deliberate testing of humans for the purpose of establishing toxicity without expected benefit to the subjects of such testing (Silbergeld et al. 2004).

For preventing harms, we need to have reliable and sufficient evidence of safety for chemicals, drugs and food, prior to permitting human exposure, particularly in our chemical world with tens of thousands of chemicals in commerce and the environment. This ethic underlies the establishment of many regulations and guidance by governments and international institutions requiring pre-approval testing of substances developed for their biological activity, such as pharmaceuticals, in order to assess likely benefits and harms prior to testing in humans. The same principle is applied for testing other chemicals developed for their toxic properties, such as pesticides. For other chemicals produced by industry, the situation is less consistent (Silbergeld et al. 2015). For the many chemicals that are already on the market, nonhuman toxicological evidence can support prudent actions to reduce exposures without the delays and human costs of awaiting evidence from observational studies.

Despite its crucial position in science based public health policy, toxicology as a field has often failed to efficiently produce timely information for decision making and prevention of harms

(Gee et al. 2013). As a consequence, policy-making in environmental and occupational health, drug and product safety, lags far behind the need for prevention of harms. There are many reasons for this, including the failure of current methods in applying toxicological information to resolve controversies among stakeholders (Silbergeld et al. 2015). Part of this is certainly related to the economic and political importance of the issues for which toxicological information is generated, such as drug and chemical approvals and legally binding standards for air and water. But toxicology as a field contributes to its own failures to generate information expeditiously and to respond to controversies through its lack of systematic methods and evidence based principles similar to those that have been successfully applied to resolve controversies and reach decisions in other fields related to public health.

The wake-up call for the field of toxicology came with the recent US National Research Council recommendation to EPA for the adoption of evidence-based methods, similar to those widely used in medicine and healthcare, in its assessments of chemical hazards and risks. This NRC report included a strong critique in of the current reliance on nontransparent process such as “weight of evidence” (NRC 2014b). The US EPA (Cogliano 2014; NRC 2014b), FDA (FDA 2009) and the European Food Safety Agency (EFSA 2010) have made public commitments to the development and application of systematic methods for evaluating evidence from the toxicological sciences. The International Agency for Research on Cancer (IARC) has begun to utilize these methods in its monographs on carcinogens (Hamra et al. 2014). With these developments, there is now wider acceptance that evidence based methods, including systematic reviews, is “the road worth taking” for toxicology (Silbergeld and Scherer 2013). Less well understood is what this acceptance entails. In this paper, we define and discuss both the core principles and methods of evidence based practice that are applicable to toxicology, with specific

reference to the ones developed and used by the Cochrane Collaboration, an international not-for-profit organization preparing, maintaining and promoting the accessibility of systematic reviews of the effects of health care (Cochrane 2015c). Using a comparison between evidence based practice and current practices in toxicology, we examine the differences, limits and advantages of both principles and methods for toxicological research and application to public health policy.

Discussion

Toxicology: a matter (not just) for experts

The importance of toxicology is widely recognized and accepted in public health policy.

However, the reliability and validity of many toxicological methods – from study design to statistical analyses - have been challenged. These limitations have significant impacts for both improving and protecting health. Recent reviews have demonstrated the low predictive value of preclinical testing in identifying novel pharmaceuticals likely to have therapeutic benefits, as well as in detecting potential adverse effects early in drug development (Krauth et al. 2014). These failures may result in costs of millions of dollars in development as well as harms to patients (Kola and Landis 2004). For non-pharmaceutical chemicals, including food additives, current toxicological methods as well as practices do not resolve controversies because of their non transparent procedures and potential for conflict of interest. Too often decisions are based on information provided by and evaluated by parties with financial ties to the products without public disclosure (Abdel-Sattar et al. 2014; Neltner et al. 2013). As a consequence, debates over the hazards of many of these agents, already in production and use, go on for decades with controversies among regulatory agencies within and among countries, states, and stakeholders. In

a recent review, we also observed that the assessment of new chemicals prior to production relies heavily on non validated methods and nontransparent data submissions (Silbergeld et al. 2015).

Despite the increasing resources devoted to toxicity testing of drugs and chemicals in terms of animals, time, and expertise, the pace of regulatory decision-making by agencies such as the Environmental Protection Agency is best described as glacial. Recently, the US National Academy of Science was called on by Congress to review NTP Annual Report on Carcinogens listings of styrene and formaldehyde as carcinogens (NRC 2014a, c). These two major industrial chemicals are produced and used in many countries at a level of millions of tons per year, and panels with different experts have expressed divergent opinions on the hazards of these two chemicals (NRC 2014a, c). Toxicological information from US National Toxicology Program and the Ramazzini Institute on the hazards and risks of these two chemicals has been publicly available for decades (Conti et al. 1988; NTP 2011; Soffritti et al. 2002), yet definitive regulatory action has been delayed. Regulatory delays concerning styrene and formaldehyde, as well as delays reaching decisions with other chemicals, have prevented actions to reduce harms resulting from continued exposures, an example of what the European Environment Agency described as “late lessons from early warnings” (Gee et al. 2013). In many cases there are no early warnings, because the majority of chemicals are not tested before marketing or are marketed with insufficient evidence of safety. This still happens (for example in US and China) in full compliance with current chemical regulatory policies such as TSCA (Silbergeld et al. 2015). A tragic example of this practice is 4-methylcyclohexanemethanol. The accidental release of this chemical in West Virginia led to the shut down of drinking water for over 700,000 people because health hazards associated with its use were largely unknown (Manuel 2014).

The limits of the discipline of toxicology and the delayed promulgation and application of effective regulatory policies based on the use of toxicological principles contributed to the impetus for the precautionary principle largely in order to empower timely preventive actions (Gee et al. 2013; Ramazzini 2004). The increasing public pressure for more rapid action to protect public health and the environment has supported policies that reduce the requirements for full information. In fact the precautionary principle definition promulgated in 1992 by the UN Conference on Environment and Development, states “In order to protect the environment, the precautionary approach shall be widely applied by States according to their capabilities. Where there are threats of serious or irreversible damage, lack of full scientific certainty shall not be used as a reason for postponing cost-effective measures to prevent environmental degradation.” (UN 1993). But the precautionary principle does not remove the need for toxicological evidence for “threats of harm” and does not help decisions that require quantitation of harm such as most air and water quality standards. Others are placing hope in alternative methods, such as “Tox21” where high throughput molecular based systems are proposed to shift the assessment of chemical hazards away from traditional experimental animal toxicology studies to methods that reduce time and the burdens on animal use in experimentation by substituting mechanism-based in vitro assays and in silico assessments (Tice et al. 2013). The jury is still out on the utility of these methods to provide sufficient evidence of safety for either pharmaceuticals or chemical regulation (Schmidt 2009) and the Tox21 program “will likely take decades to fully achieve its goals” (Tice et al. 2013). In the meantime, other policies, such as the EU REACH chemical regulation (ECHA 2015), have attempted to reduce the “burden of proof” on governments to meet the demand for information, by placing responsibility on industry to generate toxicology data under the principle of “no data, no market” (Silbergeld et al. 2015). But the quality of these

toxicological data and the methods used for their evaluation are other concerns, as discussed below.

Why is toxicology failing? The methodological failures in current nonhuman testing described by Hooijmans and Ioannidis are endemic to the field of toxicology (Hooijmans and Ritskes-Hoitinga 2013; Ioannidis et al. 2014), including inappropriate study designs and inadequate statistical analyses. New tests have been adopted, such as structure-activity analysis and many in vitro methods, without appropriate validation (Knudsen et al. 2011) and the process of updating methods is extremely slow. In many respects, toxicology is its own worst enemy. The causes of its malaise are many but not hard to identify. The most critical afflictions of toxicology at present relate to its lack of principles commonly accepted as essential to evidence based practice, an aversion to transparency and persistent adherence to nonsystematic methods. As a consequence, toxicology in practice demonstrates little consistency in terms of even assembling the relevant literature, and no clear methods for screening this literature, extracting and evaluating information, in order to objectively test its reliability as evidence. As discussed below, all of these steps precede the integration of evidence for decision making.

Of greatest concern, toxicology has failed to adopt clear principles that could enhance its acceptability. Chief among these is the continuation by toxicology to extensively rely upon “expert judgment”. This concept is embedded in nontransparent and vague principles and practices such as “weight of evidence,” which was recently strongly criticized by the NAS (NRC 2014b). Douglas Weed succinctly characterized this term in his 2005 review, in which he concluded that it is not well defined nor does it refer to a consistent or transparent methodology (Weed 2005). Some of the “principles” often cited in toxicology as indicative of reliability and

quality are of unproven relevance in ensuring the reliability and quality of evidence derived from toxicological studies. For example, the Good Laboratory Practices code (OECD 1998) is a recipe for keeping adequate records, not for ensuring appropriately designed or valid studies. The Klimisch Score (Klimisch et al. 1997), currently widely used for assessing the reliability of toxicological studies, over-values compliance with GLP and guidelines and fails to address some of the most important criteria for assessing quality of studies, such as the validity and relevance of the study design, statistical rigor, and attention to sources of bias (Agerstrand et al. 2011; Myers et al. 2009).

The largest elephant in the room is the failure of toxicology as a field to examine its own biases in terms of conflicts of interest (LaDou et al. 2010). Bero and others have demonstrated that the source of the piper's pay in research, from clinical trials to tobacco studies, introduces a predictable risk of bias in results and conclusions (Lundh et al. 2012) (Barnes and Bero 1998; Bero et al. 2007). For this reason, conflict of interest (COI) was recently proposed as an independent item in the assessment of risk of bias in the Cochrane review process (Bero 2013). Several analyses suggest that the same topic is also important in toxicology and needs more examination as well (Barnes and Bero 1998; Neltner et al. 2013). One group working on evidence based toxicology in The Navigation Guide already embeds COI as an item in its risk of bias assessment (Woodruff and Sutton 2014).

Toxicology also has a history of service to private interests which indicates a particular need to evaluate sources of funding as related not only to study bias, but also claims of evidence based practices from interested stakeholders and their consultants (Ashford et al. 2002; EBTC 2015; Denison 2014; Guzelian et al 2005; Pearce et al 2015; Toxstrategies 2015). The case of the

Klimisch Score is paradigmatic: it was proposed by industry scientists of BASF and has been widely adopted by regulators, despite its lack of validation or relevance to any systematic assessment of the quality of the studies (Klimisch et al. 1997). There are other examples of the same pressures from industry and acquiescence by regulators in terms of the test methods of the OECD chemicals program that now form the basis for the EU REACH program (Ponti et al. 2014).

A Call To (Systematic) Action

Calls for the adoption of systematic methods to support the generation of evidence in toxicology are not new and there are several organizations claiming to use “evidence based toxicology”, although there is no common accepted definition of this term (Silbergeld and Scherer 2013). At this point in time, a wide community of participation is highly recommended, within some common understanding of what this term implies. In this commentary, we recommend that those interested in evidence based toxicology, especially regulators, can usefully learn from experience in the first “evidence based” fields, medicine and healthcare, which is embodied most fully in the international Cochrane Collaboration (Cochrane 2015c). Cochrane principles and methods were considered radical and highly disputed when presented several years ago (Dickersin and Manheimer 1998) and thus we can expect a similar context for the development of systematic methods in toxicology (Silbergeld and Scherer 2013). However, we may be able to shorten this initial “postnatal” period by learning from the past. The Cochrane Collaboration has worked for over 20 years to develop both principles and methods. Their systematic methods and reviews are internationally considered as the gold-standard in medicine and health care because of their demonstrated value and reliability through decades of development, validation, application and continuous improvement (Jorgensen et al. 2006; Tovey 2014). We present the

case that the new field of “evidence based toxicology,” which at present has multiple meanings and groups working on methodologies, can learn from both the principles and practices of systematic reviews within the Cochrane Collaboration to develop consensus approaches that can also be internationally accepted. We also consider the additional benefit that the introduction of evidence-based methods in toxicology will provide by enhancing the scientific development and the quality of studies in the field, in a manner similar to the experience in clinical trials in medicine.

Learning From Cochrane: Principles First

Seventy years ago, similar problems to the ones that toxicology is now facing characterized the challenge of obtaining reliable evidence for medical practice. The use of evidence based approaches first started with the need for the postwar UK National Health System to be able to reliably evaluate evidence of demonstrably efficacious interventions and treatments in order to approve payment. This was the birth of evidence based medicine (Dickersin and Manheimer 1998). From this very practical beginning, the Cochrane Collaboration grew into an essential global partner in ensuring evidence based practices and decision-making in health. Its methods now cover diagnostic and test methods as well as interventions and methods of outcome assessment (Cochrane 2015b).

Sir Archie Cochrane’s medicine can assist toxicology as well, by bringing this essential science into harmony with the principles and practices of evidence based medicine. As a first step in developing evidence based toxicology, the principles of evidence based medicine can be adopted straight from the Cochrane prescription. These principles have been proven solid and reliable, even when addressing controversial themes (Gotzsche and Jorgensen 2013; Jefferson et al.

2010). As shown in Appendix 1, these principles state the prerequisites for ensuring that work in Cochrane will produce reliable evidence for decision making (Cochrane 2015a). These principles include: the identification and reduction of bias (that is, factors that introduce systematic error and otherwise reduce confidence in results) and methods of work that enhance the achievement of this goal through transparency at all stages, open collaboration and access, validation and improvement of methods and continuous updating of reviews. These principles consider the legitimate interest of all the stakeholders (researchers, consumers, regulators and industry), where collaboration and public health interest prevail over single interests.

Many toxicologists at this time do not abide by these principles, as is clear from a recent position statement by a group of industry, government, and academic representatives “An Appeal for the integrity of Science and Public Policy” (February 11, 2015; personal communication by email signed by Gio Gori, Wolfgang Dekant, John Doull, and Alan Boobis; text available online at <http://www.eurotox.com/an-appeal-for-the-integrity-of-science-and-public-policy/>) in which they argue that the “rules of evidence of the scientific method” are to be preferred in establishing decisions regarding assurance of safety and prevention or risk. The appeal defines the scientific method without including the principles of transparency, participation, or adherence to the identification of sources of bias, including conflict of interest. This has been one source of toxicology’s present difficulties and a major contributor to the difficulty of resolving controversies.

Learning from Cochrane: Method, Follow

“In terms of methods, many of those already developed and validated by the Cochrane Collaboration can be adopted, some will require modification, and some adjustments specific to

toxicology may require the development and validation of new formulations to achieve an evidence-based approach.

The Cochrane Collaboration has developed protocols to guide steps in the process of systematic reviews that have been demonstrated to produce useful and reliable information. These are readily adaptable to toxicology; they include, clear formulation of the problem to be reviewed, comprehensive and explicit strategies for identifying sources of information, attention to all sources of bias, including inadequate study designs, unvalidated or inappropriate methods of generating and analyzing information, and public disclosure of financial conflicts of interest.

Differences between toxicology and evidence-based practice are illustrated in Table 1.

Well validated methods and practices of Systematic Reviews, as developed by the Cochrane Collaboration, can be largely translated to toxicology (Rooney et al. 2014):

- Clarity in formulation of the problem:, defining populations, exposures, comparators, outcomes, timings, and settings of interest (PECOTS):
- Transparent and replicable processes for research strategy
- Transparent methods of data extraction and presentation
- Validation of all methods and criteria in terms of relevance to reducing bias
- Comprehensive assessment of risk of bias (study design, appropriate statistical analyses, conflict of interest)
- Transparent criteria for determining if data integration is appropriate and conducting data integration, such as meta-analysis

But challenges in developing evidence based methods specific for toxicology will require also new adequate methods, that can't be directly inherited from Cochrane. For example, while sharing common problems (and perhaps some common solutions), nonhuman preclinical studies and toxicology tests require some different methods and policies because of their differing purposes: pre-clinical studies investigate efficacy (benefits), while toxicology investigates safety (harms) (Krauth et al. 2013). There are particular aspects of nonhuman studies that will require investments and efforts to develop methods, including:

- Attention to external validity of nonhuman toxicity tests for inferring risks to humans
- Challenges to integrating information:
 - Dealing with the diversity of nonhuman species currently used in toxicity tests as well as the use of in vitro systems, organotypic cultures, transformed cell lines, and ex vivo preparations
 - Assessing the validity of “toxicity pathway” studies
- Determining the contribution and value of mechanistic studies to overall evaluation of evidence
- Moving beyond harms: generating evidence to support decisions for setting regulatory standards (that is, dose:response)

The Office of Health Assessment and Translation of the National Toxicology Program (OHAT) “Handbook for Conducting a Literature Based Health Assessment Using OHAT Approach for Systematic Review and Evidence Integration”(NTP-OHAT 2015) and the “Navigation Guide Systematic Review Methodology” (Woodruff and Sutton 2014) are two important efforts to translate and embed many of the above-mentioned Cochrane ingredients in toxicology. There is

also ongoing work for implementing specific methods for integrating and grading the quality of evidence in toxicology (Rooney et al. 2014). Particularly relevant is the implementation of GRADE (Grades of Recommendation, Assessment, Development and Evaluation), a system for grading the quality of evidence used by several organizations worldwide (including Cochrane Collaboration and WHO), with specific scales that should be tailored for rewarding sensitivity of the studies to harm detection and prevention (the main outcomes of interest for toxicology), rather than efficacy (the main outcome of interest of clinical medicine and pre-clinical studies) (Guyatt et al. 2008). Harmonization and upgrades will be necessary following the first attempts of systematic reviews in toxicology, and adherence to common principles and methods will be the first necessary step toward the application of evidence-based approaches in toxicology.

Conclusions for a new beginning of Toxicology

Improving the methods of generating systematic evidence from toxicology will not only clarify and expedite the processes of decision-making, this will also enhance the international acceptability of a common evidence base that can be fitted into national policies (NRC 2014b). This is an important and significant challenge to our field; however, we come to this challenge on the shoulders of considerable achievements in developing and applying systematic methods in other relevant fields, such as the ones obtained by the Cochrane Collaboration in its work related to evidence based medicine and health care. As with experience in Cochrane, our dedication to generate systematic evidence by ensuring comprehensive and objective analyses will improve the process of decision making, thereby preventing harms, increasing public confidence and reducing costs. Moreover, success in this effort will improve and strengthen the science of toxicology, just as adoption of the systematic approach to evaluating information from clinical

trials has resulted in the adoption of more reliable methods, with lower risk of bias and more predictive value.

References

- Abdel-Sattar M, Krauth D, Anglemeyer A, Bero L. 2014. The relationship between risk of bias criteria, research outcomes, and study sponsorship in a cohort of preclinical thiazolidinedione animal studies: A meta-analysis. *Evidence-based Preclinical Medicine* 1:11–20.
- Agerstrand M, Breitholtz M, Ruden C. 2011. Comparison of four different methods for reliability evaluation of ecotoxicity data: A case study of non-standard test data used in environmental risk assessments of pharmaceutical substances. *Environmental Sciences Europe* 23:17.
- Ashford NA, Castleman B, Frank AL, Giannasi F, Goldman LR, Greenberg M, et al. 2002. The international commission on occupational health (icoh) and its influence on international organizations. *International journal of occupational and environmental health* 8:156–162.
- Barnes DE, Bero LA. 1998. Why review articles on the health effects of passive smoking reach different conclusions. *JAMA* 279:1566–1570.
- Bero L, Oostvogel F, Bacchetti P, Lee K. 2007. Factors associated with findings of published trials of drug–drug comparisons: Why some statins appear more efficacious than others. *PLoS Med* 4:e184.
- Bero LA. 2013. Why the cochrane risk of bias tool should include funding source as a standard item. *Cochrane Database Syst Rev* 12:[10.1002/14651858.ED000075](https://doi.org/10.1002/14651858.ED000075).
- Cochrane. 2015a. Cochrane organisational policy manual. Available: [Http://community.Cochrane.Org/organisational-policy-manual](http://community.Cochrane.Org/organisational-policy-manual) [accessed 10 April 2015]
- Cochrane. 2015b. Cochrane Library. Available: [Http://www.Cochranelibrary.Com/](http://www.Cochranelibrary.Com/) [accessed 10 April 2015].
- Cochrane. 2015c. Cochrane Collaboration. Available: [Http://www.Cochrane.Org/](http://www.Cochrane.Org/) [accessed 3 June 2015].
- Cogliano V. 2014. EPA IRIS workshop on the NRC recommendations. Available: [Http://www.Epa.Gov/iris/irisworkshops/nrc_workshop/presentations/welcome_nrcwork_cogliano.Pdf](http://www.Epa.Gov/iris/irisworkshops/nrc_workshop/presentations/welcome_nrcwork_cogliano.Pdf) [accessed 3 June 2015].

- Conti B, Maltoni C, Perino G, Ciliberti A. 1988. Long-term carcinogenicity bioassays on styrene administered by inhalation, ingestion and injection and styrene oxide administered by ingestion in sprague-dawley rats, and para-methylstyrene administered by ingestion in sprague-dawley rats and swiss mice. *Ann N Y Acad Sci* 534:203–234.
- Denison R. 2014. EDF's recommendations for IRIS conflicts-of-interest disclosures, and the strong precedents for them. Environmental Defense Fund. Available: <http://blogs.edf.org/health/2014/11/11/edfs-recommendations-for-iris-conflicts-of-interest-disclosures-and-the-strong-precedents-for-them/> [accessed 7 June 2015]
- Dickersin K, Manheimer E. 1998. The Cochrane Collaboration: evaluation of health care and services using systematic reviews of the results of randomized controlled trials. *Clin Obstet Gynecol* 41:315–331.
- EBTC. 2015. Evidence-based Toxicology Collaborations Sponsors. Available: <http://www.ebtox.com/sponsors/> [accessed 7 June 2015]
- ECHA. 2015. REACH Regulation. Available: <http://echa.europa.eu/regulations/reach> [accessed 3 June 2015].
- EFSA. 2010. Application of systematic review methodology to food and feed safety assessments to support decision making. *EFSA Journal* 8; doi: 10.2903/j.efsa.2010.1637. Available: <http://www.efsa.europa.eu/en/efsajournal/pub/1637.htm#> [accessed 3 June 2015].
- FDA. 2009. Guidance for industry: Evidence-based review system for the scientific evaluation of health claims-final. Retrieved from: Date Published: January. Available: <http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/LabelingNutrition/ucm073332.htm> [accessed 3 June 2015]
- Gee D, Grandjean P, Hansen SF, van denHove S, MacGarvin M, Martin J, et al. 2013. Late lessons from early warnings: Science, precaution, innovation. European Environment Agency. Available: <http://www.eea.europa.eu/publications/late-lessons-2> [accessed 3 June 2015]
- Gotzsche PC, Jorgensen KJ. 2013. Screening for breast cancer with mammography. *The Cochrane Database Syst Rev* 6:CD001877; doi: 10.1002/14651858.CD001877.pub5.
- Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. 2008. GRADE: An emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336:924–926.

- Guzelian PS, Victoroff MS, Halmes NC, James RC, Guzelian CP. 2005. Evidence-based toxicology: a comprehensive framework for causation. *Hum Exp Toxicol* 4:161–201.
- Hamra GB, Guha N, Cohen A, Laden F, Raaschou-Nielsen O, Samet JM, et al. 2014. Outdoor particulate matter exposure and lung cancer: A systematic review and meta-analysis. *Environ Health Perspect* 122:906–911.
- Hooijmans CR, Ritskes-Hoitinga M. 2013. Progress in using systematic reviews of animal studies to improve translational research. *PLoS Med* 10(7):e1001482; doi: 10.1371/journal.pmed.1001482.
- Ioannidis JPA, Greenland S, Hlatky MA, Khoury MJ, Macleod MR, Moher D, et al. 2014. Increasing value and reducing waste in research design, conduct, and analysis. *Lancet* 383:166–175.
- Jefferson T, Di Pietrantonj C, Rivetti A, Bawazeer GA, Al-Ansary LA, Ferroni E. 2010. Vaccines for preventing influenza in healthy adults. *Cochrane Database Syst Rev* (7):CD001269. doi: 10.1002/14651858.CD001269.pub4.
- Jorgensen AW, Hilden J, Gotzsche PC. 2006. Cochrane Reviews compared with industry supported meta-analyses and other meta-analyses of the same drugs: Systematic review. *BMJ* 333:782.
- Josefson D. 2001. Johns Hopkins faces further criticism over experiments. *BMJ* 323:531–531.
- Katz RV, Kegeles SS, Kressin NR, Green BL, Wang MQ, James SA, et al. 2006. The Tuskegee Legacy Project: Willingness of minorities to participate in biomedical research. *J Health Care Poor Underserved* 17:698–715.
- Klimisch HJ, Andreae M, Tillmann U. 1997. A systematic approach for evaluating the quality of experimental toxicological and ecotoxicological data. *Regul Toxicol Pharmacol* 25:1–5.
- Knudsen TB, Kavlock RJ, Daston GP, Stedman D, Hixon M, Kim JH. 2011. Developmental toxicity testing for safety assessment: New approaches and technologies. *Birth Defects Res B Dev Reprod Toxicol* 92:413–420.
- Kola I, Landis J. 2004. Can the pharmaceutical industry reduce attrition rates? *Nat Rev Drug Discov* 3:711–715.
- Krauth D, Woodruff TJ, Bero L. 2013. Instruments for assessing risk of bias and other methodological criteria of published animal studies: A systematic review. *Environ Health Perspect* 121:985–992.

- Krauth D, Anglemyer A, Philipps R, Bero L. 2014. Nonindustry-sponsored preclinical studies on statins yield greater efficacy estimates than industry-sponsored studies: A meta-analysis. *PLoS Biol* 12:e1001770.
- LaDou J, Castleman B, Frank A, Gochfeld M, Greenberg M, Huff J, et al. 2010. The case for a global ban on asbestos. *Environ Health Perspect* 118:897–901.
- Lundh A, Sismondo S, Lexchin J, Busuioac OA, Bero L. 2012. Industry sponsorship and research outcome. *Cochrane Database Syst Rev* 12:MR000033. doi: 10.1002/14651858.MR000033.pub2.
- Manuel J. 2014. Crisis and emergency risk communication lessons from the Elk River spill. *Environ Health Perspect* 122:A214–219.
- Myers JP, vom Saal FS, Akingbemi BT, Arizono K, Belcher S, Colborn T, et al. 2009. Why public health agencies cannot depend on good laboratory practices as a criterion for selecting data: The case of Bisphenol A. *Environ Health Perspect* 117:309–315.
- Neltner TG, Alger HM, O'Reilly JT, Krimsky S, Bero LA, Maffini MV. 2013. Conflicts of interest in approvals of additives to food determined to be generally recognized as safe: Out of balance. *JAMA Intern Med* 173:2032–2036.
- NRC. 2014a. Review of the styrene assessment in the National Toxicology Program 12th report on carcinogens. Washington, DC:The National Academies Press.
- NRC. 2014b. Review of EPA's Integrated Risk Information System (IRIS) process. Washington, DC:The National Academies Press.
- NRC. 2014c. Review of the formaldehyde assessment in the National Toxicology Program 12th report on carcinogens. Washington, DC:The National Academies Press.
- NTP. 2011. Report on carcinogens (12th ed.):DIANE Publishing Company. Available: ntp.niehs.nih.gov/go/roc12 [accessed 3 June 2015]
- NTP-OHAT. 2015. Handbook for conducting a literature based health assessment using ohat approach for systematic review and evidence integration. Available: https://ntp.Niehs.Nih.Gov/ntp/ohat/pubs/handbookjan2015_508.Pdf [accessed 10 April 2015].
- OECD. 1998. OECD principles on good laboratory practice (as revised in 1997). Available: http://www.oecd.org/chemicalsafety/testing/oecdseriesonprinciplesofgoodlaboratorypractice_glpandcompliancemonitoring.htm [accessed 3 June 2015]

- Pearce N, Blair A, Vineis P, Ahrens W, Andersen A, Anto JM et al. 2015. IARC Monographs: 40 Years of Evaluating Carcinogenic Hazards to Humans. *Environ Health Perspect* 6:507–14.
- Ponti B, Bettinetti R, Dossi C, Vignati DA. 2014. How reliable are data for the ecotoxicity of trivalent chromium to daphnia magna? *Environ Toxicol Chem* 33:2280–2287.
- Ramazzini C. 2004. The precautionary principle: Implications for research and policy-making. Statement of the Collegium Ramazzini. *Am J Ind Med* 45:380–381.
- Rooney AA, Boyles AL, Wolfe MS, Bucher JR, Thayer KA. 2014. Systematic review and evidence integration for literature-based environmental health science assessments. *Environ Health Perspect* 122:711–718.
- Schmidt CW. 2009. Tox 21: New dimensions of toxicity testing. *Environ Health Perspect* 117:A348–A353.
- Silbergeld E, Lerman S, Hushka L. 2004. Ethics. Human health research ethics. *Science* 305:949.
- Silbergeld E, Scherer RW. 2013. Evidence-based toxicology: Strait is the gate, but the road is worth taking. *ALTEX* 30:67–73.
- Silbergeld EK, Mandrioli D, Cranor CF. 2015. Regulating chemicals: Law, science, and the unbearable burdens of regulation. *Annu Rev Public Health* 36:175–91; doi: 10.1146/annurev-publhealth-031914-122654.
- Soffritti M, Belpoggi F, Lambertin L, Lauriola M, Padovani M, Maltoni C. 2002. Results of long-term experimental studies on the carcinogenicity of formaldehyde and acetaldehyde in rats. *Ann N Y Acad Sci* 982:87–105.
- Tice RR, Austin CP, Kavlock RJ, Bucher JR. 2013. Improving the human hazard characterization of chemicals: A tox21 update. *Environ Health Perspect* 121:756–765.
- Toxstrategies. 2015. Systematic Reviews and Evidence-Based Toxicology. ToxStrategies, Inc. Available: <https://www.toxstrategies.com/services/systematic-reviews-and-evidence-based-toxicology/> [accessed 7 June 2015].
- Tovey D. 2014. The role of the Cochrane Collaboration in support of the WHO nutrition guidelines. *Adv Nutr* 5:35–39.
- UN. 1993. Report of the United Nations conference on environment and development, Rio de Janeiro, 3-14 june 1992. New York: United Nations. Available: <http://www.un.org/documents/ga/conf151/aconf15126-1annex1.htm> [accessed 3 June 2015]

Weed DL. 2005. Weight of evidence: A review of concept and methods. *Risk Analysis* 25:1545–1557.

Woodruff TJ, Sutton P. 2014. The Navigation Guide systematic review methodology: A rigorous and transparent method for translating environmental health science into better health outcomes. *Environ Health Perspect* 122:1007–1014.

Appendix 1: Cochrane's Principles (Cochrane 2015a)

1. Collaboration: by fostering global co-operation, teamwork, and open and transparent communication and decision-making.
2. Building on the enthusiasm of individuals: by involving, supporting and training people of different skills and backgrounds.
3. Avoiding duplication of effort: by good management, co-ordination and effective internal communications to maximise economy of effort.
4. Minimising bias: through a variety of approaches such as scientific rigour, ensuring broad participation, and avoiding conflicts of interest.
5. Keeping up-to-date: by a commitment to ensure that Cochrane Systematic Reviews are maintained through identification and incorporation of new evidence.
6. Striving for relevance: by promoting the assessment of health questions using outcomes that matter to people making choices in health and health care.
7. Promoting access: by wide dissemination of our outputs, taking advantage of strategic alliances, and by promoting appropriate access models and delivery solutions to meet the needs of users worldwide.
8. Ensuring quality: by applying advances in methodology, developing systems for quality improvement, and being open and responsive to criticism.
9. Continuity: by ensuring that responsibility for reviews, editorial processes and key functions is maintained and renewed.
10. Enabling wide participation: in our work by reducing barriers to contributing and by encouraging diversity.

Table 1. Methods: Toxicology vs Evidence Based Toxicology

Toxicology	Evidence Based Toxicology
Unclear answers to unclear questions.	Clear formulation of problem (PECOTS)
Non-comprehensive research strategy	Comprehensive research strategy
Non Transparent methods	Transparent Methods
Unvalidated Methods	Requirement to validate methods prior to use
Inadequate study design (effect size; expected variability; etc)	Adequate study design
No or inconsistent assessment of risk of bias	Assessment of risk of bias
Inadequate or no statistical modeling	Appropriate statistical modeling based on appropriate study design
Conflict of Interest usually not disclosed	Conflict of Interest Disclosed
unvalidated or irrelevant guidelines for practice (Klimisch Scores and Good Laboratory Practices)	Specific evaluation of Risk of Bias and compliance with evidence based practice