



Lactational Exposure to Polybrominated Diphenyl  
Ethers and Its Relation to Early Childhood  
Anthropometric Measurements

Kate Hoffman, Michelle Mendez, Anna Maria Siega-Riz,  
Amy H. Herring, Andreas Sjödin, and Julie L. Daniels

<http://dx.doi.org/10.1289/EHP201>

Received: 29 May 2015  
Revised: 28 October 2015  
Accepted: 20 April 2016  
Published: 6 May 2016

**Note to readers with disabilities:** *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

# Lactational Exposure to Polybrominated Diphenyl Ethers and Its Relation to Early Childhood Anthropometric Measurements

Kate Hoffman,<sup>1,2</sup> Michelle Mendez,<sup>1</sup> Anna Maria Siega-Riz,<sup>1</sup> Amy H. Herring,<sup>1</sup> Andreas Sjödin,<sup>3</sup> and Julie L. Daniels<sup>1</sup>

<sup>1</sup>University of North Carolina Gillings School of Global Public Health, Chapel Hill, North Carolina, USA; <sup>2</sup>Nicholas School of the Environment, Duke University, Durham, North Carolina, USA; <sup>3</sup>Centers for Disease Control and Prevention, National Center for Environmental Health, Division for Laboratory Sciences, Atlanta, Georgia, USA

**Address correspondence to** Julie Daniels, Department of Epidemiology, University of North Carolina Gillings School of Global Public Health, McGavran-Greenberg Hall CB#7435, Chapel Hill, North Carolina 27599 USA. Telephone: 919-966-7096. Fax: 919-966-2089. E-mail: [juliedaniels@unc.edu](mailto:juliedaniels@unc.edu)

**Running title:** Lactational PBDE exposure and early-life anthropometric measurements

**Acknowledgments:** This research was supported by grants from the U.S. Environmental Protection Agency (RD832736) and the National Institute of Environmental Health Sciences (P30ES10126). The work of KH was supported by the NIEHS Environmental Biostatistics Training Program (T32ES007018).

**Disclaimer:** The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official position of the National Cancer Institute and the CDC.

**Declaration of Competing Interests:** All authors declare no actual or potential competing financial interests.

## **Abstract**

**Background:** Polybrominated diphenyl ethers (PBDEs) are ubiquitous environmental contaminants that may impact growth and development. We investigated the association between exposure to PBDEs via breast milk and anthropometric measurements in early childhood.

**Methods:** The Pregnancy Infection and Nutrition (PIN) Babies studies followed a cohort of North Carolina pregnant women and their children through 36 months of age. Breast milk samples obtained at 3 months postpartum were analyzed for PBDEs. We collected height and weight records from well-baby doctor visits and also measured children during study visits (n=246 children with >1400 anthropometric measurements). We assessed the relationship between breast milk concentrations of five PBDE congeners—BDE-28, 47, 99, 100, and 153—and child's weight-for-age, height-for-age, and weight-for-height z-scores (WAZ, HAZ, and WHZ, respectively), adjusting for age; maternal age, race, pre-pregnancy BMI; parity; smoking during pregnancy; and breastfeeding, and stratifying by sex.

**Results:** Overall PBDE exposures via breast milk were not associated with early life anthropometric measures in the PIN Babies cohort. When stratified by sex, PBDEs in milk were inversely associated with WHZ for boys; however, associations did not follow a consistent pattern across the concentration gradient and were imprecisely estimated. Among girls, PBDEs tended to be associated with increased WHZ, with the exception of BDE-153 which was inversely associated with WHZ, though all estimates were imprecisely estimated.

**Conclusions:** We observed little evidence of associations between early-life PBDE exposures via breast milk and anthropometric measurements overall; however, our results prompt the need for sex-specific investigations in larger cohorts.

## **Background**

Polybrominated diphenyl ethers (PBDEs) have been added to a variety of household products, including electronics, carpets and the polyurethane foam in furniture, to delay combustion and meet fire safety requirements (Darnerud et al. 2001). PBDEs are not chemically bound to the products to which they are added and leach into the environment over time (Sjodin et al. 2003). Resistant to degradation, PBDEs are persistent in the environment and have become ubiquitous contaminants (Darnerud et al. 2001; Hites 2004). Dust and diet are thought to be the primary sources of exposure to PBDEs in the general population (Fraser et al. 2009; Johnson et al. 2010; Stapleton et al. 2008; Stapleton et al. 2012; Wu et al. 2007); however, infants are also exposed to PBDEs prenatally and through breastfeeding (Johnson-Restrepo and Kannan 2009; Toms et al. 2009).

Toxicological data indicate that PBDEs disrupt endocrine function (Butt and Stapleton 2013; Ellis-Hutchings et al. 2006; Richardson et al. 2008; Stoker et al. 2004; Szabo et al. 2009; Zhou et al. 2001; Zhou et al. 2002). Prenatal and early life exposures to PBDEs have been shown to impact body weight in early development in animals (Ferne et al. 2006; Gee and Moser 2008; Kim et al. 2009; Kodavanti et al. 2010; Suvorov et al. 2009). The direction of association, however, has not been consistent across studies; some report increases in body weight following exposure to specific congeners (Suvorov et al. 2009), others report no association, or decreases in body weight (Kim et al. 2009; Kodavanti et al. 2010). Observed differences by sex, which have been inconsistent, further complicate our understanding of the relationship between PBDEs and early life growth and adiposity (Ferne et al. 2006; Gee and Moser 2008; Kim et al. 2009; Kodavanti et al. 2010; Suvorov et al. 2009). Few studies have investigated early life exposure in human populations; however, prenatal PBDE exposures have been inversely associated with

birth weight and body mass index (BMI) in newborn infants in some (Chao et al. 2007; Harley et al. 2011; Zhang et al. 2011), but not all studies (Mazdai et al. 2003; Tan et al. 2009). The implications of exposures to PBDEs in infancy on children's growth remain unknown. Thus, we prospectively investigate the association between early-life exposure to PBDEs through breast milk and early childhood weight-for age, height-for-age, and weight-for-height in a cohort of central North Carolina children.

## **Methods**

### ***Study Sample***

The Pregnancy, Infection and Nutrition (PIN) Babies Study began in 2004 to follow the infants born to women who participated in the PIN Pregnancy and Postpartum studies through 3 years of age (Daniels et al. 2010). Women in this study were enrolled from the University of North Carolina prenatal care clinics early in pregnancy. Self-administered questionnaires, telephone interviews and a brief questionnaire at the hospital after delivery were used to collect health and lifestyle information during the pregnancy. Information on postpartum health and lifestyle, as well as children's development, was collected during three in-home interviews at 3, 12, and 36 months postpartum. Children's physicians also provided information on growth and development at well-baby doctor visits. All children in the PIN Babies Study were singleton births and free from major birth defects (n=585). For these analyses we have further restricted to term infants (>37 weeks gestation). All study protocols were approved by the Institutional Review Board at the University of North Carolina-Chapel Hill and the Centers for Disease Control (CDC) and Prevention, and all mothers provided informed consent.

### ***Sample Collection***

Of the 585 women in the PIN Babies Study, 304 mothers were still lactating and provided breast milk samples at 3 months postpartum. Women providing breast milk samples were similar to those not providing samples with respect to age, but had higher educational attainment (96.4% vs. 88.2% had at least college education). Participants providing samples were also more likely to be white (11.4 % of participants with samples were non-white while 27.6% of participants without milk samples were non-white). Details of the breast milk sample collection are available in Daniels et al. 2010 (Daniels et al. 2010). Briefly, lactating women followed written instructions to pump both breasts, gently mix milk, and use a plastic pipette to transfer milk into three 1.5 mL tubes. Women stored the milk in their freezer until a research team picked up the sample during a home data collection visit later the same day. Samples were then transported on ice to -80 °C storage freezers until they were analyzed.

### ***Sample Analysis***

Breast milk samples were analyzed for 9 PBDE congeners by the Organic Analytical Toxicology Branch of the National Center for Environmental Health at the CDC and Prevention using previously described methods (Sjodin et al. 2004). Briefly, milk samples were added to diatomaceous earth packed in a solid phase extraction cartridge (3mL) and <sup>13</sup>C-labeled internal standards were added. Target analytes and lipids were extracted using an automated modular solid-phase extraction system which dried the sample onto the diatomaceous earth with pressurized nitrogen and eluted analytes and lipids with dichloromethane. Lipid content was determined gravimetrically, and the final analytical determination of PBDEs was performed by gas chromatography/isotope-dilution high-resolution mass spectrometry. Both wet weight and

lipid normalized concentrations (reported as nanograms per gram of lipid) were produced. Two quality controls and two blank samples were added to each batch of 16 study samples. Quality assurance practices in the laboratory were regularly monitored (Sjodin et al. 2004).

### ***Anthropometric Measurement***

Infant length (the usual method used before 24 months of age) or standing height and weight were recorded by health care professionals during well-baby visits, which are recommended by the American Academy of Pediatrics at 1, 2, 4, 6, 9, 12, 15, 18, 24, 30, and 36 months of age (American Academy of Pediatrics (AAP) 2008). Because there was some variability in the frequency and timing of visits, we linked measurements to the child's age at the time of the visit (reported by the health care provider) or, when age was not recorded, to age calculated using the visit date and the child's date of birth. If the day of the month was not reported for a visit, we used the 15<sup>th</sup> of the month in calculation of age. We excluded information for visits conducted before breast milk samples were collected ( $\leq 3$  months post-partum) to ensure that exposure temporally preceded outcomes. At the 36-month home visit, PIN Babies study staff also conducted in-home assessments of weight and standing height, according to the National Health and Nutrition Examination Surveys protocols (Centers for Disease Control and Prevention (CDC) 2007).

Weight and length/height of each child were compared to the CDC and Prevention/National Center for Health Statistics (NCHS) growth charts to calculate sex-standardize weight-for age, height-for-age, and weight-for-height z-scores (WAZ, HAZ, and WHZ, respectively)(Kuczmarski et al. 2002). Measurements that corresponded to z-scores  $>5$  or  $<5$  were considered implausible and were excluded from analyses (n=33 observations).

### ***Statistical Analysis***

A total of 1457 WAZ (mean=5.9; 1-10 per child), 1414 HAZ (mean=5.7; 1-13 per child), and 1432 WHZ (mean= 5.8; 1-10 per child) measurements were available for the 246 children the PIN cohort from 3 to 36 months of age. Measurements were concentrated in the first year (ages 3-12 months, n=843 measurements of 246 children); however, 350 measurements (n=229 children) were made between 12 and 24 months, and 277 (n=192 children) measurements were made after children were 24 months of age. To determine how PBDEs related to WAZ, HAZ, and WHZ in early childhood, we used linear mixed effects regression models. Mixed effects models easily accommodate data that are correlated (multiple measurements in each child) with varying numbers of measurements with variable spacing between measurements (Fitzmaurice et al. 2004). Because PBDE concentrations in breast milk are highly correlated used separate models to examine the association between quartiles of breast milk levels of each of the 5 most detected PBDEs (BDE 28, 47, 99, 100, and 153) and WAZ, HAZ, and WHZ. While a summary measure of PBDEs is sometimes calculated; because a number of previous studies have shown differences in the biological activity of PBDE congeners, we present results for each congener separately. We included random effects in all models to allow for child-specific variation in both size and growth rates. Because we anticipated that the relationship between age and anthropometric measurements may not be linear, we investigated a linear trend in age at the time of the measurement as well as a quadratic trend for the fixed and random effects. Using a restricted maximum likelihood ratio test, we determined including the quadratic term for random effects improved model fit.

As reported previously for this population, five PBDE congeners, BDE-28, 47, 99, 100, and 153 were detected in  $\geq 91\%$  of samples (Daniels et al. 2010). Milk samples with PBDE values below

the limit of detection (LOD) were assigned a concentration of the LOD divided by the square root of 2 for statistical analyses. The four PBDEs congeners infrequently detected (in less than 70% of the PIN Babies sample; BDE 66, 85, 154, 183 (Daniels et al. 2010)) were not evaluated in relation to childhood anthropometric measurements. Because PBDEs are lipophilic, levels in breast milk are often adjusted for lipid content in epidemiologic studies. Although investigators generally agree that adjustment is beneficial, the most appropriate method of adjustment remains unclear (O'Brien et al. 2015; Schisterman et al. 2005). Accordingly, we used multiple methods to account for confounding by the percent lipid in milk samples. First we conducted analyses using lipid normalized PBDE concentrations. We also included wet weight concentrations in regression models and included lipid as a covariate. Finally, we conducted analyses of lipid normalized concentrations with lipid included as a covariate (O'Brien et al. 2015). In general the results were similar using all methods. Here, we present results using lipid normalized concentrations, which are more commonly presented, and discuss differences observed using other methods of adjustment.

Based on our a priori expectation of their association with both PBDE milk concentrations and childhood anthropometric measurements, we adjusted analyses for maternal age ( $\leq 25$  years, 26-30 years, 31-35 years, and  $\geq 36$  years), race (white vs. non-white), smoking status during pregnancy (yes vs. no), pre-pregnancy body mass index (BMI;  $< 25$ , 25-30, and  $> 30$ ), parity (0 vs. 1+), and the child's age at the time of the measurement (continuous age in months and age<sup>2</sup>). We examined the appropriate form (continuous or nominal) for each covariate in relation to the outcomes (form of variables used is reflected in Table 1). We included a time varying covariate for breastfeeding status at the time of each anthropometric measurement, indicating that the child was currently breastfeeding, recently breastfed (within 2 months), or was no longer

breastfeeding. We did not adjust for maternal education or gestational weight gain because these factors were not strongly related to PBDE levels in breast milk and had little impact on the relationships between milk exposure and anthropometric measurements. Because we anticipated that the impact of exposure to PBDEs via lactation may vary with age, we also considered an interaction term between quartiles of PBDEs and age. Interaction terms between PBDE concentrations and age were not statistically significant and did not provide additional insights into the relationship between PBDEs and anthropometric measurements. Thus we present results for the average effect of PBDE exposure via breast milk from 3 months of age through 36 months of age.

We hypothesize the potential pathway by which PBDE exposures may be associated with growth is through endocrine disruption, which suggests the possibility of differential associations by sex. We conducted sex-stratified analyses to investigate differences in associations among boys and girls. All analyses were conducted in SAS (Version 9.2; SAS Institute Inc, Cary, NC).

## **Results**

Of the children whose mothers provided milk samples (n=304), 266 (87.5%) had height or weight measurements (children with measurements did not differ from other children with respect to included covariates). We excluded children who were born before 37 weeks gestation (n=20). Our final sample consisted of 246 children with at least one measurement of height or weight and a maximum of 13 measurements (>1400 measurements for each outcome; median measurements per child=6.0; interquartile range: 4-8 measurements). As in the larger PIN Babies cohort, mothers tended to be over age 31 at the time of their child's birth (52.6%) and were predominantly white (88.6%; Table 1). Children in our sample were slightly more likely to

be male (54.3%) and were generally breastfed longer than 10 months (64.2%). WAZ and HAZ for children in our sample were slightly higher at a given age than the 2000 CDC normative population (mean WAZ = 0.1 and mean HAZ = 0.5). However, children's WHZ tended to be lower, indicating that PIN Babies participants were slightly leaner than the normative population.

As reported previously in the PIN babies cohort, BDE-28, 47, 99, 100, and 153 were detected frequently in breast milk samples ( $\geq 90.7\%$  of samples; Table 2) and were generally correlated with each other (Daniels et al. 2010). Levels of BDE-47 tended to be higher than those of the other congeners (median breast milk level = 26.8 ng/g lipid) while the levels of BDE-28 were generally the lowest (median breast milk level = 2.0 ng/g lipid).

Associations between lipid normalized PBDE levels in breast milk and WAZ, HAZ, and WHZ are shown in Figures 1, 2, and 3, respectively (effect estimates and 95% confidence intervals (CIs) presented in supplemental tables S1-S3). PBDE exposures via breast milk were generally not associated with early-life anthropometric measures (WAZ, HAZ, and WHZ) in the PIN Babies cohort as a whole after adjusting for age, maternal age, race, pre-pregnancy BMI, parity, smoking during pregnancy, and breastfeeding status. Effect estimates generally hovered around the null, did not follow a consistent pattern across the concentration gradient, and were imprecisely estimated. Different methods of adjusting for the lipid content of milk (e.g. including lipid as a covariate) produced slightly different results; however, results with these methods remained largely null and did not provide additional insights regarding the relationship between locational exposure to PBDEs and anthropometric measures in early childhood.

When we stratified by sex, higher levels of lipid normalized PBDEs in milk tended to be associated with decreased WHZ for boys, particularly for BDE-28, 47 and 100; however, as in

the full cohort (boys and girls combined), the patterns were inconsistent across the concentration gradient and imprecisely estimated and generally not statistically significant (Figure 3B).

Results for WAZ were similar, with high PBDE levels in breast milk general associated with lower WAZ in boys comparing all quartiles of exposure to the lowest quartile, but not completely consistent and imprecise (Figure 1B). Associations between lipid normalized PBDE levels and anthropometric measurements were largely null for girls with the exception of BDE-153, which our data suggest may be inversely associated with WAZ and WHZ [Figure 1C and 2C; mean difference (95% CI) = -0.47 (-0.96, 0.03) and -0.24 (-0.69, 0.22) comparing the 4<sup>th</sup> quartile to the 1<sup>st</sup> respectively]. When we adjusted for lipids by adding a term for lipid to the regression model, effect estimates further attenuated with two notable exceptions: First, the inverse association between the highest quartile of BDE 153 and WHZ among girls, was unchanged (mean differences comparing the 4<sup>th</sup> quartile to the 1<sup>st</sup>=0.46, 95% CI: -0.98, 0.07). Second, the association between BDE-28 and WHZ among girls was more prominently positive (mean difference 2<sup>nd</sup> quartile=0.69, 95% CI: 0.19,1.22; mean difference 3rd quartile=0.21, 95% CI -0.32, 0.75; mean difference 4th quartile=0.68, 95% CI 0.06, 1.31).

## **Discussion**

Despite widespread exposure to PBDEs, relatively little is known about their potential to impact health, particularly regarding children's growth. In our cohort of central North Carolina children, higher PBDE levels in breast milk were generally not associated with anthropometric measurements from 3 to 36 months of age. There was some suggestion of decreased WHZ in boys more highly exposed via breast milk and increased WHZ in girls; however our small sample hindered our ability to more precisely assess differences in the associations by sex.

Although not a direct measurement of adiposity, WHZ is more reflective of fat mass than WAZ or HAZ.

Several previous toxicological studies have documented the endocrine disrupting properties of PBDEs and reported sex-differences in offspring weight gain in the early postnatal period, although the direction of reported associations has been inconsistent. For example, prenatal exposure to the commercial penta-PBDE mixture, DE71, was associated with decreased weight gain in female, but not male rat pups (Kodavanti et al. 2010). Conversely, prenatal exposure to BDE-209 was related to lower weight gain in early development in male rat pups but not female rat pups (Kim et al. 2009), a finding consistent with our results in boys which suggested lower WHZ with increasing PBDE levels in breast milk.

As we are aware, the only other study to assess postnatal exposure to PBDE found an inverse association between serum BDE-153 levels and BMI percentile at 7 years of age (Erkin-Cakmak et al. 2015 ); a finding which is consistent with our results suggesting lactational exposure to BDE-153 may be associated with lower WHZ. Additionally, previous studies have reported prenatal exposures to be associated with lower birth weight and BMI in newborn infants (Chao et al. 2007; Harley et al. 2011; Zhang et al. 2011). Our data for BDE-153 and WHZ are consistent with these results; however, we did not consistently observe similar associations for other congeners. There are several possible explanations for differences in the results of our work and those from previous research. A primary difference is that our results are based on post-natal PBDE exposure through breast milk, which is correlated with, but different from the prenatal exposure metrics used by most previous studies. We did not have exposure data to explore associations reflecting the prenatal period, which may be a more sensitive developmental stage for PBDEs exposure. It is possible that the associations between PBDEs and children's height

and weight differs over time. We did not see differences by age when including an interaction term in models, however the children in our study were relatively young, and the majority of the anthropometric measurements were collected in the first year of life, potentially limiting our ability to detect differences through three years of age (results not shown).

PBDE milk concentrations in our cohort were higher than concentrations reported by most similarly timed studies from the United States and considerable higher than those reported abroad (Daniels et al. 2010; Erdogrul et al. 2004; Frederiksen et al. 2009; Johnson-Restrepo et al. 2007; Schechter et al. 2003; She et al. 2007; Toms et al. 2007). Although a clear trend has yet to be established, exposure to PBDEs is likely to decrease in the coming decades because their use has been phased out in many countries.

Our work should be interpreted in the context of several limitations. PBDEs are ubiquitous contaminants in the home environment (e.g. (Stapleton et al. 2009)), raising the possibility that breast milk samples, which were collected by mothers in their own homes, may be contaminated. Although mothers were given instruction on sample collection to reduce this possibility, exposure misclassification is possible and may have biased results toward the null. Similarly, we have not accounted for competing environmental exposures or dietary patterns in these analyses. However, by design, all children included in analyses were breastfed through 3 months of age, and the majority was breastfed for longer than 10 months. Although we had over 1,400 anthropometric measurements, our samples size of 246 children was relatively small, limiting our ability to investigate difference in effect by sex or over time. Additionally, generalizability of our results is limited by the homogeneity of the PIN Babies cohort. PIN Babies included mothers that attended UNC prenatal care clinics, delivered generally healthy singleton infants, completed several years of follow-up, and were not very demographically

diverse. Mothers in the PIN cohort were highly educated. These mothers may have been more likely to intervene to mitigate sub-optimal growth, potentially masking any impact of early-life-PBDE exposures. Finally, the majority of height and weight measurements for the PIN Babies cohort were collected at doctor's offices. These measurements may be subject to random error because they were collected at multiple medical office sites by staff members that had not been trained using a standard method. Such error would likely diminish our ability to detect associations. However, we found excellent correlation between the health providers' measurements and our standardized study measurements; among 50 participants with both measures collected within a 2 week period around 36 months of age, the correlation between the two measurements was 0.85 ( $p < 0.001$  for correlation). This comparison offers assurance that health provider measurements provide reliable measures of growth between study visits and contribute favorably to our analysis. Additionally, we benefitted from ability to evaluate this association in a well-characterized, somewhat homogenous, group of mothers and children with reasonable exposure variability.

## **Conclusions**

We observed little evidence of associations between early-life PBDE exposures via breast milk and anthropometric measurements; however, our results suggesting possible differences in effects between boys and girls prompt the need for sex-specific investigations in larger cohorts. Further study is needed to determine whether PBDEs are adversely associated with children's growth and whether that persists throughout as children age, particularly as altered growth in early-life has long-term health implications.

## References

American Academy of Pediatrics (AAP). 2008. Recommendations for preventive pediatric health care.

Butt CM, Stapleton HM. 2013. Inhibition of thyroid hormone sulfotransferase activity by brominated flame retardants and halogenated phenolics. *Chemical research in toxicology* 26:1692-1702.

Centers for Disease Control and Prevention (CDC). 2007. National health and nutrition examination survey (nhanes): Anthropometry procedures manual. Available: [http://www.cdc.gov/nchs/data/nhanes/nhanes\\_07\\_08/manual\\_an.pdf](http://www.cdc.gov/nchs/data/nhanes/nhanes_07_08/manual_an.pdf) [accessed November 12 2012].

Chao HR, Wang SL, Lee WJ, Wang YF, Papke O. 2007. Levels of polybrominated diphenyl ethers (pbdes) in breast milk from central taiwan and their relation to infant birth outcome and maternal menstruation effects. *Environ Int* 33:239-245.

Daniels JL, Pan IJ, Jones R, Anderson S, Patterson DG, Jr., Needham LL, et al. 2010. Individual characteristics associated with pbde levels in u.S. Human milk samples. *Environ Health Perspect* 118:155-160.

Darnerud PO, Eriksen GS, Johannesson T, Larsen PB, Viluksela M. 2001. Polybrominated diphenyl ethers: Occurrence, dietary exposure, and toxicology. *Environ Health Perspect* 109 Suppl 1:49-68.

Ellis-Hutchings RG, Cherr GN, Hanna LA, Keen CL. 2006. Polybrominated diphenyl ether (pbde)-induced alterations in vitamin a and thyroid hormone concentrations in the rat during lactation and early postnatal development. *Toxicol Appl Pharmacol* 215:135-145.

Erdogru O, Covaci A, Kurtul N, Schepens P. 2004. Levels of organohalogenated persistent pollutants in human milk from kahramanmaras region, turkey. *Environ Int* 30:659-666.

Erkin-Cakmak A, Harley K, Chevrier J, Bradman A, Kogut K, Huen K, et al. 2015 In utero and childhood polybrominated diphenyl ether exposures and body mass at age 7 years: The chamacos study. *Environ Health Perspect* 123.

Fernie KJ, Laird Shutt J, Ritchie IJ, Letcher RJ, Drouillard K, Bird DM. 2006. Changes in the growth, but not the survival, of american kestrels (*falco sparverius*) exposed to environmentally relevant polybrominated diphenyl ethers. *J Toxicol Environ Health A* 69:1541-1554.

Fitzmaurice GM, Laird NM, Ware JH. 2004. Applied longitudinal analysis. New Jersey:John Wiley & Sons, Inc. .

Fraser AJ, Webster TF, McClean MD. 2009. Diet contributes significantly to the body burden of pbdes in the general u.S. Population. *Environ Health Perspect* 117:1520-1525.

Frederiksen M, Vorkamp K, Thomsen M, Knudsen LE. 2009. Human internal and external exposure to pbdes--a review of levels and sources. *International journal of hygiene and environmental health* 212:109-134.

Gee JR, Moser VC. 2008. Acute postnatal exposure to brominated diphenylether 47 delays neuromotor ontogeny and alters motor activity in mice. *Neurotoxicol Teratol* 30:79-87.

Harley KG, Chevrier J, Schall RA, Sjodin A, Bradman A, Eskenazi B. 2011. Association of prenatal exposure to polybrominated diphenyl ethers and infant birth weight. *Am J Epidemiol* 174:885-892.

Hites RA. 2004. Polybrominated diphenyl ethers in the environment and in people: A meta-analysis of concentrations. *Environ Sci Technol* 38:945-956.

Johnson-Restrepo B, Addink R, Wong C, Arcaro K, Kannan K. 2007. Polybrominated diphenyl ethers and organochlorine pesticides in human breast milk from massachusetts, USA. *Journal of environmental monitoring : JEM* 9:1205-1212.

Johnson-Restrepo B, Kannan K. 2009. An assessment of sources and pathways of human exposure to polybrominated diphenyl ethers in the united states. *Chemosphere* 76:542-548.

Johnson PI, Stapleton HM, Sjodin A, Meeker JD. 2010. Relationships between polybrominated diphenyl ether concentrations in house dust and serum. *Environ Sci Technol* 44:5627-5632.

Kim TH, Lee YJ, Lee E, Kim MS, Kwack SJ, Kim KB, et al. 2009. Effects of gestational exposure to decabromodiphenyl ether on reproductive parameters, thyroid hormone levels, and neuronal development in sprague-dawley rats offspring. *J Toxicol Environ Health A* 72:1296-1303.

Kodavanti PR, Coburn CG, Moser VC, MacPhail RC, Fenton SE, Stoker TE, et al. 2010. Developmental exposure to a commercial pbde mixture, de-71: Neurobehavioral, hormonal, and reproductive effects. *Toxicol Sci* 116:297-312.

Kuczumarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, et al. 2002. 2000 cdc growth charts for the united states: Methods and development. *Vital Health Stat* 11:1-190.

Mazdai A, Dodder NG, Abernathy MP, Hites RA, Bigsby RM. 2003. Polybrominated diphenyl ethers in maternal and fetal blood samples. *Environ Health Perspect* 111:1249-1252.

O'Brien KM, Upson K, Cook NR, Weinberg CR. 2015. Environmental chemical in urine and blood: Improving methods for creatinine and lipid adjustment. *Environ Health Perspect* (In press).

Richardson VM, Staskal DF, Ross DG, Diliberto JJ, DeVito MJ, Birnbaum LS. 2008. Possible mechanisms of thyroid hormone disruption in mice by bde 47, a major polybrominated diphenyl ether congener. *Toxicol Appl Pharmacol* 226:244-250.

Schechter A, Pavuk M, Papke O, Ryan JJ, Birnbaum L, Rosen R. 2003. Polybrominated diphenyl ethers (pbdes) in u.S. Mothers' milk. *Environ Health Perspect* 111:1723-1729.

Schisterman EF, Whitcomb BW, Louis GM, Louis TA. 2005. Lipid adjustment in the analysis of environmental contaminants and human health risks. *Environ Health Perspect* 113:853-857.

She J, Holden A, Sharp M, Tanner M, Williams-Derry C, Hooper K. 2007. Polybrominated diphenyl ethers (pbdes) and polychlorinated biphenyls (pcbs) in breast milk from the pacific northwest. *Chemosphere* 67:S307-317.

Sjodin A, Patterson DG, Jr., Bergman A. 2003. A review on human exposure to brominated flame retardants--particularly polybrominated diphenyl ethers. *Environ Int* 29:829-839.

Sjodin A, McGahee EE, 3rd, Focant JF, Jones RS, Lapeza CR, Zhang Y, et al. 2004. Semiautomated high-throughput extraction and cleanup method for the measurement of polybrominated diphenyl ethers and polybrominated and polychlorinated biphenyls in breast milk. *Anal Chem* 76:4508-4514.

Stapleton HM, Kelly SM, Allen JG, McClean MD, Webster TF. 2008. Measurement of polybrominated diphenyl ethers on hand wipes: Estimating exposure from hand-to-mouth contact. *Environ Sci Technol* 42:3329-3334.

Stapleton HM, Klosterhaus S, Eagle S, Fuh J, Meeker JD, Blum A, et al. 2009. Detection of organophosphate flame retardants in furniture foam and u.S. House dust. *Environ Sci Technol* 43:7490-7495.

Stapleton HM, Eagle S, Sjodin A, Webster TF. 2012. Serum pbdes in a north carolina toddler cohort: Associations with handwipes, house dust, and socioeconomic variables. *Environ Health Perspect* 120:1049-1054.

Stoker TE, Laws SC, Crofton KM, Hedge JM, Ferrell JM, Cooper RL. 2004. Assessment of de-71, a commercial polybrominated diphenyl ether (pbde) mixture, in the edsp male and female pubertal protocols. *Toxicol Sci* 78:144-155.

Suvorov A, Battista MC, Takser L. 2009. Perinatal exposure to low-dose 2,2',4,4'-tetrabromodiphenyl ether affects growth in rat offspring: What is the role of igf-1? *Toxicology* 260:126-131.

Szabo DT, Richardson VM, Ross DG, Diliberto JJ, Kodavanti PR, Birnbaum LS. 2009. Effects of perinatal pbde exposure on hepatic phase i, phase ii, phase iii, and deiodinase 1 gene expression involved in thyroid hormone metabolism in male rat pups. *Toxicol Sci* 107:27-39.

Tan J, Loganath A, Chong YS, Obbard JP. 2009. Exposure to persistent organic pollutants in utero and related maternal characteristics on birth outcomes: A multivariate data analysis approach. *Chemosphere* 74:428-433.

Toms LM, Harden FA, Symons RK, Burniston D, Furst P, Muller JF. 2007. Polybrominated diphenyl ethers (pbdes) in human milk from australia. *Chemosphere* 68:797-803.

Toms LM, Sjodin A, Harden F, Hobson P, Jones R, Edenfield E, et al. 2009. Serum polybrominated diphenyl ether (pbde) levels are higher in children (2-5 years of age) than in infants and adults. *Environ Health Perspect* 117:1461-1465.

Wu N, Herrmann T, Paepke O, Tickner J, Hale R, Harvey LE, et al. 2007. Human exposure to pbdes: Associations of pbde body burdens with food consumption and house dust concentrations. *Environ Sci Technol* 41:1584-1589.

Zhang L, Tian Y, Yang XF, Cui C, Gao Y, Wang XJ, et al. 2011. [concentration of polybrominated diphenyl ethers in umbilical cord serum and the influence on newborns birth outcomes in shanghai]. *Zhonghua Yu Fang Yi Xue Za Zhi* 45:490-493.

Zhou T, Ross DG, DeVito MJ, Crofton KM. 2001. Effects of short-term in vivo exposure to polybrominated diphenyl ethers on thyroid hormones and hepatic enzyme activities in weanling rats. *Toxicol Sci* 61:76-82.

Zhou T, Taylor MM, DeVito MJ, Crofton KM. 2002. Developmental exposure to brominated diphenyl ethers results in thyroid hormone disruption. *Toxicol Sci* 66:105-116.

Table 1: Selected characteristics of included PIN Babies participants.

Characteristic	n	%
Total population	246	100.0
Gender		
Male	133	54.3
Female	112	45.7
Maternal race		
White	218	88.6
Non-white	28	11.4
Maternal age		
<=25 Years	33	13.4
26-30 Years	82	33.3
31-35 Years	100	40.7
>=36 Years	31	12.6
Maternal Education		
High school graduate or less	9	3.7
College graduate	118	48.0
Graduate training or degree	119	48.4
Smoking during pregnancy		
Yes	9	3.7
No	232	94.3
Missing	5	2.0
Parity		
0	130	52.9
1+	116	47.2
Breast feeding duration		
<10 months	88	35.8
10-14 months	113	45.9
>=15 months	45	18.3
	<u>Mean</u>	<u>SD<sup>a</sup></u>
Maternal pre-pregnancy BMI (kg/m <sup>2</sup> ) <sup>a</sup>	23.4	± 4.7
Child WAZ <sup>b</sup>	0.1	± 1.1
Child HAZ <sup>c</sup>	0.5	± 1.0
Child WHZ <sup>d</sup>	-0.1	± 1.2

<sup>a</sup> SD: standard deviation; BMI: body mass index

<sup>b</sup> 1457 observations for 239 participants (measured between 1 and 13 times)

<sup>c</sup> 1414 observations for 240 participants (measured between 1 and 10 times)

<sup>d</sup> 1432 observations for 239 participants (measured between 1 and 10 times)

Table 2: Lipid-adjusted concentrations (ng/g lipid) of commonly detected PBDEs in 246 breast milk samples collected at 3 months postpartum in 2004-2006.

PBDE Congener	Percent Detect (LOD)	Mean	Quartile Ranges			
			Q1	Q2	Q3	Q4
BDE 28	97.1 (0.3)	3.2	ND to 1.3	1.4 to 2.0	2.1 to 3.7	3.8 to 49.6
BDE 47	99.6 (1.3)	50.2	ND to 15.7	15.8 to 27.7	27.8 to 54.2	54.3 to 1430
BDE 99	90.7 (1.1)	10.4	ND to 2.65	2.8 to 5.2	5.3 to 11.2	11.3 to 299
BDE 100	98.4 (0.3)	10.4	ND to 2.4	2.5 to 5.15	5.25 to 10.4	10.5 to 188
BDE 153	99.2 (0.3)	14.8	ND to 2.65	2.8 to 5.6	5.7 to 13.5	13.55 to 229

<sup>a</sup> ND (non-detectable) values are treated as the LOD divided by the square root of 2 in the calculation of the mean.

## Figure Legends

Figure 1: Mean difference estimates and 95% confidence intervals by quartile of PBDE congener (1<sup>st</sup> quartile, referent=circle; 2<sup>nd</sup> quartile=square; 3<sup>rd</sup> quartile=triangle; 4<sup>th</sup> quartile=diamond) for WAZ, adjusted for adjusting for age (age and age<sup>2</sup>), maternal age (<=25 years, 26-30 years, 31-35 years, and ≥36 years), race (white vs. non-white), pre-pregnancy BMI (<25, 25-30, and >30), parity (0 vs. 1+), smoking during pregnancy (yes vs. no), and breastfeeding status (current, within the last 2 months, and no longer breastfeeding): A) full cohort; B) boys only; C) girls only.

Figure 2: Mean difference estimates and 95% confidence intervals by quartile of PBDE congener (1<sup>st</sup> quartile, referent=circle; 2<sup>nd</sup> quartile=square; 3<sup>rd</sup> quartile=triangle; 4<sup>th</sup> quartile=diamond) for HAZ, adjusted for adjusting for age (age and age<sup>2</sup>), maternal age (<=25 years, 26-30 years, 31-35 years, and ≥36 years), race (white vs. non-white), pre-pregnancy BMI (<25, 25-30, and >30), parity (0 vs. 1+), smoking during pregnancy (yes vs. no), and breastfeeding status (current, within the last 2 months, and no longer breastfeeding): A) full cohort; B) boys only; C) girls only.

Figure 3: Mean difference estimates and 95% confidence intervals by quartile of PBDE congener (1<sup>st</sup> quartile, referent=circle; 2<sup>nd</sup> quartile=square; 3<sup>rd</sup> quartile=triangle; 4<sup>th</sup> quartile=diamond) for WHZ, adjusted for adjusting for age (age and age<sup>2</sup>), maternal age (<=25 years, 26-30 years, 31-35 years, and ≥36 years), race (white vs. non-white), pre-pregnancy BMI (<25, 25-30, and ≥30), parity (0 vs. 1+), smoking during pregnancy (yes vs. no), and breastfeeding status (current, within the last 2 months, and no longer breastfeeding): A) full cohort; B) boys only; C) girls only.

Figure 1.




Figure 2.




Figure 3.

