

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Male Breast Cancer Incidence and Mortality Risk in the Japanese Atomic Bomb Survivors – Differences in Excess Relative and Absolute Risk from Female Breast Cancer**

M.P. Little and D.M. McElvenny

#### **Table of Contents**

##### **Supplemental Material A. Supplemental Analysis Tables**

**Table A1.** Percent probability of causation of incident case of various types of cancer at age 65, associated with 50 mSv received at ages 35 and 55, using relative risk models developed by UNSCEAR (United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) 2008), BEIR VII (Committee to Assess Health Risks from Exposure to Low Levels of Ionizing Radiation 2006), and those of the present paper.

**Table A2.** Breast cancer cases and person years (PY) of follow-up by dose group, and whether not-in-city (NIC) survivors are included or not.

**Table A3.** Breast cancer deaths and person years (PY) of follow-up by dose group

**Table A4.** Male breast cancer cases and deaths and person years (PY) of follow-up by interval of follow-up.

##### **Supplemental Material B. Variables used for background models of breast cancer incidence and mortality in fits of generalized excess absolute risk model**

**Table B1.** Optimal variables used in baseline models, selected to minimize Akaike Information Criterion (AIC), and augmented to make polynomially complete.