

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Long-Term Exposure to Traffic-Related Air Pollution and Risk of Incident Atrial Fibrillation: A Cohort Study

Maria Monrad, Ahmad Sajadieh, Jeppe Schultz Christensen, Matthias Ketzel, Ole Raaschou-Nielsen, Anne Tjønneland, Kim Overvad, Steffen Loft, and Mette Sørensen

Table of Contents

Table S1: Association between 1- and 5-years residential exposure to NO₂ and NO_x and risk of atrial fibrillation

Table S2: Associations between established risk factor and risk of atrial fibrillation in the Diet, Cancer and Health cohort.

Figure S1: Exposure-response relationship between 1-year time-weighted mean NO_x and risk of atrial fibrillation. The vertical whiskers show incidence rate ratios with 95 % confidence interval at the median of exposure categories (deciles).

Figure S2: Exposure-response relationship between 1-year time-weighted mean NO₂ and risk of atrial fibrillation. The vertical whiskers show incidence rate ratios with 95 % confidence interval at the median of exposure categories (deciles).

Figure S3: Exposure-response relationship between 10-years time-weighted mean NO_x and risk of atrial fibrillation. The vertical whiskers show incidence rate ratios with 95 % confidence interval at the median of exposure categories (deciles).

Figure S4: Exposure-response relationship between 10-years time-weighted mean NO₂ and risk of atrial fibrillation. The vertical whiskers show incidence rate ratios with 95 % confidence interval at the median of exposure categories (deciles).