



ENVIRONMENTAL HEALTH PERSPECTIVES

<http://www.ehponline.org>

Perinatal Exposure to Traffic-Related Air Pollution and Autism Spectrum Disorders

Tong Gong, Christina Dalman, Susanne Wicks, Henrik Dal,
Cecilia Magnusson, Cecilia Lundholm, Catarina Almqvist,
and Göran Pershagen

<http://dx.doi.org/10.1289/EHP118>

Received: 16 March 2016

Revised: 23 June 2016

Accepted: 12 July 2016

Published: 5 August 2016

Note to readers with disabilities: *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.



Perinatal Exposure to Traffic-Related Air Pollution and Autism Spectrum Disorders

Tong Gong¹, Christina Dalman^{2,3}, Susanne Wicks^{2,3}, Henrik Dal^{2,3}, Cecilia Magnusson², Cecilia Lundholm¹, Catarina Almqvist^{1,4}, and Göran Pershagen^{5,6}

1 Department of Medical Epidemiology and Biostatistics, Karolinska Institutet

2 Department of Public Health Sciences, Karolinska Institutet

3 Center for Epidemiology and Community Medicine, Stockholm County Council

4 Astrid Lindgren Children's Hospital, Karolinska University Hospital

5 Institute of Environmental Medicine, Karolinska Institutet

6 Center for Occupational and Environmental Medicine, Stockholm County Council
Stockholm, Sweden

Corresponding author:

Tong Gong
Department of Medical Epidemiology and Biostatistics
Karolinska Institutet
SE-171 77 Stockholm, Sweden
T. +46 8 524 84450
Email. tong.gong@ki.se

Running title: Exposure to air pollution and autism spectrum disorders

Acknowledgements:

Financial support was provided through the Swedish Research Council for Health, Working Life and Welfare (FORTE 2012-0573 and 2015-00289), the Swedish Research Council (VR) 2011-3060, VR in partnership with FORTE, FORMAS and VINNOVA [Cross-disciplinary research program concerning children's and young people's mental health], VR through the Swedish Initiative for Research on Microdata in the Social And Medical Sciences (SIMSAM) framework grant 340-2013-5867, HKH Kronprinsessan Lovisas förening för barnsjukvård, and the Strategic Research Program in Epidemiology at Karolinska Institutet.

Conflict of interest: None.

Abstract

Background: Studies from the United States indicate that exposure to air pollution in early life is associated with autism spectrum disorders (ASD) in children, but the evidence is not consistent with European data.

Method: Children born and residing in Stockholm during 1993-2007 with an ASD diagnosis were identified through multiple health registers and classified as cases (n=5,136). A randomly selected sample of 18,237 children from the same study base constituted controls. Levels of nitrogen oxides (NO_x) and particulate matter with diameters <10µm (PM₁₀) from road traffic were estimated at residential addresses during mother's pregnancy and the child's first year of life by dispersion models. Odds ratios (OR) and 95% confidence intervals (CI) for ASD with or without intellectual disability (ID) were estimated using logistic regression models after conditioning on municipality and calendar year of birth as well as adjustment for potential confounders.

Result: Air pollution exposure during the prenatal period was not associated with ASD overall (OR 1.00, 95% CI 0.86-1.15 per 10 µg/m³ increase in PM₁₀ and OR 1.02, 95% CI 0.94-1.10 per 20 µg/m³ increase in NO_x during mother's pregnancy). Similar results were seen for exposure during the first year of life, and for ASD in combination with ID. An inverse association between air pollution exposure and ASD risk was observed among children of mothers who moved to a new residence during pregnancy.

Conclusion: Early life exposure to low levels of NO_x and PM₁₀ from road traffic does not appear to increase the risk of ASD.

Introduction

Autism spectrum disorders (ASD) are a range of childhood neurodevelopmental disorders characterized by deficits in communication and social interaction as well as restricted repetitive behaviors (American Psychiatric Association. 2013). The global prevalence appears to have increased over the past decades (Baird et al. 2006; Centers for Disease Control and Prevention 2014; Lauritsen et al. 2004; Nassar et al. 2009), partially due to the increased awareness and changes of clinical practices. In Sweden, 1.5% of children are currently diagnosed with ASD compared to 0.8% a decade ago (Idring et al. 2015; Lundstrom et al. 2015). Among ASD-affected children, intellectual disability (ID) is relatively common and those children often display significant clinical impairment (Buescher et al. 2014).

Early twin and family studies suggested ASD to be highly heritable (Bailey et al. 1995; Lichtenstein et al. 2010; Ritvo et al. 1989), however, recent studies have shown a moderate heritability for ASD (Ronald et al. 2011; Sandin et al. 2014), suggesting a substantial non-genetic component contributing to the etiology of the disorder. Moreover, twin studies have recently reported a modest genetic component but a significant environmental contribution to the correlation between ASD and ID (Hoekstra et al. 2009, 2010). Some maternal factors such as infections during pregnancy and a history of depression or antidepressant use have been associated with ASD (Lee et al. 2015; Rai et al. 2013). There is also growing evidence that perinatal and neonatal risk factors e.g. small for gestational age, preterm birth, low birth weight and cesarean delivery may affect the development of ASD (Buchmayer et al. 2009; Hultman et al. 2002). However, a recent meta-analysis did not indicate any specific pregnancy or delivery condition explaining the etiology of autism (Gardener et al. 2011).

Exposure to various air pollutants including ozone (Becerra et al. 2013; Jung et al. 2013; Volk et al. 2013), carbon monoxide (Becerra et al. 2013; Jung et al. 2013), nitrogen oxides (NO_x) (Becerra et al. 2013; Jung et al. 2013; Volk et al. 2013), sulfur dioxide (Jung et al. 2013),

particulate matter (PM) (Becerra et al. 2013; Jung et al. 2013; Kalkbrenner et al. 2010; Kalkbrenner et al. 2015; Raz et al. 2014; Windham et al. 2006; Volk et al. 2013), metals (Kalkbrenner et al. 2010; Palmer et al. 2009; Windham et al. 2006), and other hazardous air pollutants (Kalkbrenner et al. 2010; Windham et al. 2006) have been linked to a modestly increased risk of ASD, which may be explained by systemic inflammation affecting the central nervous system development in utero (Allen et al. 2014). However, a recent European meta-analysis could not confirm this from those exposed by NO_x and PM during perinatal life (Guxens et al. 2016). One explanation of the discrepant findings could be residual confounding from socioeconomic factors (Braveman et al. 2005; Hajat et al. 2013; Magnusson et al. 2012; Rai et al. 2012; Thomas et al. 2012). For example, a previous register-based study in Sweden found that lower familial socioeconomic status (SES) was associated with an increased risk of ASD through multiple dimensions of SES measures (Rai et al. 2012), which is contrary to findings from the United States (Thomas et al. 2012). Pollutant-specific effects could also contribute to the discrepant findings. In one study, the positive association was found for prenatal exposure to particulate matter with an aerodynamic diameter ≤ 2.5 micrometer (PM_{2.5}) but not to particulate matter with a diameter of ≤ 10 micrometer (PM₁₀) (Raz et al. 2014). Additionally, associations may differ for ASD with or without accompanying intellectual impairment, although this was not seen in a recent US study (Kalkbrenner et al. 2015).

Our aim was to investigate the association between exposure to air pollution from road traffic and the risk of ASD in children with or without presence of ID, with careful adjustment for individual- and area-level SES as well as other potential confounders. We selected a very large study base, allowing for more informative subgroup analyses than in earlier studies.

Methods

Population and study design

We conducted a case-control study based on the Stockholm Youth Cohort (SYC), a prospective cohort study including all children who resided in Stockholm County for at least four years during 2001-2007 (Idring et al. 2012). To investigate the effect of air pollution during prenatal and postnatal periods, we selected a sub-population from SYC including children born and living in Stockholm County all the time between 1993 and 2007 and with biological mothers living in Stockholm County one year before and one year after the child's birth (n=277, 478). Each child born in Sweden or each immigrant staying in Sweden for at least one year is assigned a unique personal identity number, which enables accurate linkage from different health registers to various socio-demographic background information while maintaining individual anonymity (Ludvigsson et al. 2009).

The cases constituted children with ASD (n=5,529) identified from the National Patient Register (NPR), the Clinical Database for Child and Adolescent Psychiatry in Stockholm (PASTILL), the Habilitation Register (HAB), and the Stockholm Regional Health Care Data warehouse (VAL) until December 31, 2011. These registers cover all public-financed health seeking pathways for ASD, with or without presence of ID, in Stockholm County during the whole study period, described in detail in a validation study (Idring et al. 2012). Diagnosis and care for ASD and ID was based on the International Classification of Diseases 9th or 10th revisions (ICD-9/10 codes: 299/F81 for ASD and 317-319/F70-79 for ID), the Diagnostic and Statistical Manual of mental disorders 4th version (DSM-IV code: 299 for ASD and 317-319 for ID), or any utilization of habilitation services followed by an ASD diagnosis. Cases were further divided into ASD with and without ID, regardless of whichever diagnosis came first. We excluded adopted children (n=2, 0.04%), multiple births (n=162, 3.1%), and births which were not recorded in the Medical Birth Register (n=229, 4.1%).

We selected a random sample of 20,000 children from the sub-population of SYC as controls and further excluded 420 (2.1%) who developed ASD during follow-up. Among the remaining 19,580 controls, we excluded adopted children (n=15, 0.1%), multiple births (n=585, 3.1%), and births which were not recorded in the Medical Birth Register (n=743, 3.8%).

Exposure assessment

Detailed descriptions of the air pollution exposure assessment methodology are available in previous publications (Bellander et al. 2001; Gruzieva et al. 2012). Briefly, a Gaussian air quality dispersion model was used to estimate the temporal and spatial distribution of NO_x and PM₁₀ in Stockholm County during the study period. This was based on emission databases for NO_x in 1990, 1995, 2000, 2002-2004, and interpolations estimated levels during the remaining years of the exposure period. PM₁₀ did not show any time trends during this period and model calculations were based on the year 2004. A street canyon contribution was added for the most polluted street segments in the inner city of Stockholm with multistory houses on both sides. Comparisons between model estimated and monitoring station measured NO_x levels during the exposure period provided an R² of 0.74-0.80, indicating a moderate-to-high model performance (Johansson et al. 2008). For PM₁₀ the R² was 0.61 (Eneroth et al. 2006). Relevant residential addresses of the study subjects were geocoded and pollutant levels emanating from road traffic were estimated at these coordinates from the dispersion models and used to calculate annual average concentrations for NO_x and PM₁₀. NO_x is used as a marker for tail pipe emissions, such as fine particles, while PM₁₀ reflects coarse particulates, mainly originating from road dust. To account for changes in exposure levels among those moving to another residence, time-weighted NO_x and PM₁₀ concentrations related to road traffic emissions were calculated based on all registered addresses during the pregnancy and the child's first year of life.

Covariates

Information on child characteristics including birth year, gender, sib order, gestational age, birth weight, congenital malformation and maternal characteristics including season of conception, smoking during pregnancy, and marital status at child birth were obtained from the Medical Birth Register. Data regarding family characteristics at child birth including disposable income within household, maternal and paternal education as well as employment of the mother and father were retrieved from the longitudinal integration database for health insurance and labor market studies, originally from Statistics Sweden. Information on maternal and paternal age at child birth, municipality of the mother at child birth, and parental birth countries was retrieved from the Total Population Register. Furthermore, area-based socioeconomic characteristics at birth year were measured by a neighborhood deprivation index (Sariaslan et al. 2013). Neighborhood was defined by the small-area market statistics (SAMS), which is based on small socioeconomically homogeneous areas with on average 1000 residents. Information on welfare beneficiaries, unemployment, immigrants, divorce rate, income, education, residential mobility, and criminal conviction rate from Statistics Sweden were linked with each SAMS area to calculate a neighborhood deprivation index using principal component analysis (Sariaslan et al. 2013). The neighborhood deprivation index was further categorized into tertiles. Information on family history of psychiatric disorders including schizophrenia, bipolar disorders, ID, non-affective psychosis, and other diagnoses in the mental and behavioral disorders chapter from ICD-10 (F-diagnoses) was obtained through NPR, VAL and PASTILL (see Table S1 for diagnostic codes).

Statistical analyses

To estimate the independence of pollutants over time, we calculated correlation coefficients for NO_x and PM_{10} over the pregnancy period of the mother and child's first year of life. Socio-demographic characteristics were compared among cases and controls using the t-test or chi-

square test. To assess the association between exposure to pollutants during each specific period and ASD in children and account for differences in diagnostic practices over time and across municipalities, we used conditional logistic regression models and conditioned on calendar year and municipality of birth (see Table S2 for detailed information on classification of municipalities). First, the two air pollution components were treated as continuous variables and fixed exposure increments per $20\mu\text{g}/\text{m}^3$ for NO_x and per $10\mu\text{g}/\text{m}^3$ for PM_{10} were used in all models to estimate the risk of ASD overall, with and without ID. Second, pollutants were categorized into quartiles based on their distribution in each time period and the lowest quartiles served as reference group when modeling the association with ASD overall. To assess effect modification, we examined the association between continuous exposure to either pollutant and ASD overall, with and without ID by gender, sib order, mother's marital status, neighborhood deprivation, highest education between parents at child birth, mother's smoking status during pregnancy as well as residential mobility during mother's pregnancy via inclusion of the interaction terms in the regression models. Wald tests were used to examine the statistical significance of interaction terms, using $p < 0.05$ as significance level. In sensitivity analyses, we excluded children with ASD diagnoses before 2 years of age, those born before year 2003 (when we have complete coverage on ASD diagnosis from birth from all registers), children with congenital malformation, those with either parent having any psychiatric disorders, children of foreign-born parents, children born to mothers with pre-eclampsia, pre-gestational and gestational diabetes, premature rupture of the membranes, placental abruption, children born before 37 weeks of gestation, or children born with a birth weight $< 2,500\text{g}$, as they are often at greater risk of ASD. To further explore the potential spatial over-adjustment, we provided estimates of the association between pollutants and ASD without conditioning on municipality of birth. We also ran a sensitivity

analysis by including multiple births, for which the information on sib order and birth weight was ambiguous (The Swedish National Board of Health and Welfare. 2003).

We used a directed acyclic graph to determine potential confounders of the investigated associations (see Figure S1 in the supplementary materials). A series of models were run step-wise to assess the gradual changes of risk estimates by further adjustment for potential confounders. We present odds ratios (OR) and 95% confidence intervals (CI) from crude models and models adjusted for gender, birth month, sib order, maternal age, paternal age, mother's marital status, parents' birth countries, mother's education, father's education, mother's employment, father's employment, disposable income within household, and neighborhood deprivation. Statistical analyses were conducted using SAS version 9.4 (SAS Institute, NC, USA).

The study protocol was reviewed and approved by the regional ethical review board in Stockholm, Sweden.

Result

Table 1 lists characteristics of the study population. Compared to controls, ASD cases were on average 2 years younger, more likely to be boys, have a parent with less than 12 years of education, being unemployed, or with psychiatric disorders. Maternal characteristics were also slightly different between cases and controls. For example, mothers of cases were less often married or cohabiting, but more often smoked during pregnancy.

Figure 1 shows boxplots of air pollutant levels during the pregnancy and the child's first year of life, respectively. The arithmetic mean levels of NO_x from local traffic were 11.0 µg/m³ during mother's pregnancy, and dropped somewhat to 9.8 µg/m³ during the postnatal period (see Table S3). On the other hand, the yearly arithmetic mean levels of PM₁₀ were relatively

constant (4.2-4.4 $\mu\text{g}/\text{m}^3$). NO_x was closely correlated with PM_{10} ($r^2 \geq 0.7$) over the study period as both have local traffic as the major source.

The associations between air pollution exposure as a continuous measure and the risk of ASD overall, with and without ID are shown in Figure 2. We did not observe any differences in risk of ASD overall, with or without ID by exposure during pregnancy to NO_x or PM_{10} after adjusting for potential confounders (e.g. adjusted OR was 1.02 (95% CI 0.94-1.10) for ASD overall by per 20 $\mu\text{g}/\text{m}^3$ increase of NO_x and 1.06 (95% CI 0.89-1.26) for ASD with ID, and 1.01 (95% CI 0.93-1.10) for ASD without ID). Corresponding ORs for an increment of 10 $\mu\text{g}/\text{m}^3$ of PM_{10} were 1.00 (95% CI 0.86-1.15), 1.03 (95% CI 0.74-1.42) and 0.98 (95% CI 0.84-1.15), respectively. Results were similar for exposure during the child's first year of life (see Figure 2 and Table S4). However, when pollutants were categorized into quartiles, children living in areas within the 3rd and 4th quartiles of air pollution exposure had a slightly lower risk of ASD overall compared to those living in areas within the least polluted quartile, especially in relation to exposure during pregnancy (Table 2). We observed no major confounding by individual or neighborhood covariates on the association between pollutant levels and ASD (see Table 2 and Table S4). However, the adjustments consistently generated somewhat lower ORs for ASD without ID, which constituted the major group among the cases.

We observed a similar pattern of associations in subgroups based on parental education (Figure 3 for ASD overall, Figures S2 and S3 for ASD with and without ID). There was a suggested inverse relation between air pollution exposure and ASD for those in the most deprived neighborhoods, especially for ASD without ID. A statistically significant interaction was seen for residential mobility, where a decreased risk for ASD with air pollution exposure pre- or postnatally was only seen among those changing residential address during pregnancy (all p-interactions < 0.03). Further analyses showed that moving patterns were related to SES

and psychiatric illness among the parents (Table S5). Interactions with air pollution exposure in relation to ASD were also investigated for several other characteristics, including gender, sib order, mother's smoking and marital status, but no statistically significant effect modification was observed (data not shown).

In sensitivity analyses including multiple births (see Table S6) or excluding children born to mothers with changing residential address during pregnancy, cases diagnosed before 2 years of age, children with congenital malformation, children born before 2003, children born to mothers with pre-eclampsia, pre-gestational and gestational diabetes, premature rupture of the membranes, placental abruption, children of foreign-born parents, children born with a <37 weeks of gestation or with a birth weight <2,500 gram, we found similar patterns of results for both pollutants as in our main analysis (Table S7). Furthermore, in analyses not conditioning on municipality of birth, we observed that exposure to NO_x and PM₁₀ during pre- or post-natal period appeared to be associated with a decreased risk of ASD (Table S8).

Discussion

Among children born between 1993-2007 in Stockholm County, we found that pre- and post-natal exposure to either NO_x or PM₁₀ was not associated with ASD overall, with or without ID. There was an inverse association between air pollution exposure and ASD risk for children of mothers who changed residential addresses during pregnancy, which could be partly explained by confounding by SES and psychiatric diseases in the parents.

We previously reported no association between traffic-related air pollution during pre- or post-natal periods and sub-clinical ASD outcomes using a twin cohort from Stockholm, however, risk estimates tended to lie below one with wide confidence intervals (Gong et al. 2014). The current study showed a similar result using a much larger sample but contradict the positive associations reported in several previous studies from the United States (Kalkbrenner et al. 2010; Kalkbrenner et al. 2015; Raz et al. 2014; A. L. Roberts et al. 2013;

Volk et al. 2011; Volk et al. 2013). One possible explanation for the inconsistent results could be the lower levels of air pollution in Stockholm. For example, mean levels of PM₁₀ in California, USA and Taiwan were reported at about 25-36 and 58 µg/m³ (Becerra et al. 2013; Jung et al. 2013; Kalkbrenner et al. 2015; Volk et al. 2013). However, the local traffic related PM₁₀ concentrations in the current study was 4.3 µg/m³ during mother's pregnancy and child's first year of life and the background PM₁₀ level generated from long-distance transportation near Stockholm County has remained at a rather stable level at 10 µg/m³ across the whole study period (Burman and Norman 2013). Previous studies have generally reported on nitrogen dioxide (NO₂) with mean levels in California, US and Taiwan of 32-43 µg/m³ (converted from 17 and 28.8 ppb with temperature at 25 Celsius degrees) (Becerra et al. 2013; Jung et al. 2013; Volk et al. 2013). We used NO_x as a marker since it better reflects the tail pipe emissions but it is less often measured or reported. The average urban background levels of NO₂ during the study period in Stockholm decreased from around 20 to 14 µg/m³ (Durant et al. 2014), and were thus considerably lower than the levels in the areas reported in earlier publications. Furthermore, the air pollution levels in Stockholm County, constituting the catchment area for our cohort and another cohort included in a recent European meta-analysis, are lower than in most (NO_x) or all (PM₁₀) other areas included this meta-analysis (Guxens et al. 2016). It is possible that we may have missed an association that is primarily seen at higher levels of exposure.

Residential mobility during pre- and post-natal life could also contribute to the inconsistent results. Young maternal age, being unmarried, having psychiatric diseases, and low socioeconomic status have been associated with residential mobility during pregnancy (Bell and Belanger 2012; Fell et al. 2004; Miller et al. 2010; Tulloch et al. 2010) and may thereby affect offspring's ASD risk estimates (Rai et al. 2012; Andrea L. Roberts et al. 2013; Thomas et al. 2012). Most previous studies have not investigated the potential impact by residential

mobility (Jung et al. 2013; Kalkbrenner et al. 2015; A. L. Roberts et al. 2013; Volk et al. 2013). One report from the Nurses' Health Study II found that the positive association of PM_{2.5} and ASD was stronger among non-movers, although no interaction test was performed based on moving status (Raz et al. 2014). We observed an inverse association among families that changed their residential addresses during mother's pregnancy. The reason for this is unclear and the influence of moving on the association between air pollution and ASD, which was related to SES and psychiatric disorders among the parents, should be investigated in further studies. Furthermore, misclassification of municipality at delivery among movers could contribute to the observed inverse association as we did not take into account the effect of municipalities where the mother lived prior to delivery.

Another explanation of the inconsistent results could be different proportions of high- and low-functioning ASD in different study settings. For example, one study in USA reported 40% of the ASD cases also report having ID (Kalkbrenner et al. 2015), but there were only 19% of ASD cases with co-existing ID diagnosis in our study. If the positive association could only be seen in this subtype of ASD, the higher proportion of ASD without ID in our study might have resulted in diluted associations.

Residual confounding could also contribute to the inconsistent findings by e.g. socioeconomic indicators. Children from higher SES families were more likely to be diagnosed with ASD in previous studies and the OR estimates for both air pollutants appeared slightly lower after adjusting for confounders including birth year, birth order, parental age, income, education, and ethnicity (Becerra et al. 2013; Kalkbrenner et al. 2015; Raz et al. 2014; A. L. Roberts et al. 2013; Thomas et al. 2012; Volk et al. 2013). Family and contextual SES covariates had relatively small effects on the association between air pollutants and ASD in our study.

Sweden has a rather universal healthcare system compared to the private sector dominated

healthcare system in the US (Anell et al. 2012), which may lead to differences in case ascertainment and bias related to SES.

We could not completely rule out the possibility of an association in certain subgroups, such as in those born to mothers not moving during pregnancy and among children of parents without any psychiatric diagnoses. As noted above residential mobility, SES and psychiatric disorders were related, making it difficult to disentangle their specific influence on the associations between air pollution exposure and ASD. A familial aggregation of ASD and other psychiatric conditions has been documented (Daniels et al. 2008). Our data suggest that the effect of air pollution on ASD could be masked by confounding from psychiatric diagnoses, partly related to differential moving patterns and SES.

Our study has several strengths. Firstly, the linkage of encrypted data from multiple registers enabled us to include cases and controls from the same study base, and to retrieve detailed information on validated outcomes (Idring et al. 2012) and highly reliable measures at individual level. Secondly, the air pollution exposure assessment methodology has been validated and generated positive associations for several outcomes in children of our study area, such as asthma, allergy and lung function disturbances (Melen et al. 2008; Nordling et al. 2008; Schultz et al. 2012). Using register-based information on moving dates for each residential address, time-weighted estimates of pollutant levels based on the duration of stay at the respective address was also taken into consideration. Thirdly, because of the large sample size, we had sufficient statistical power to perform subgroup specific analyses.

Our study also has some limitations. Due to the late establishment of PASTILL, HAB and VAL registers, we did not have a full age coverage of the diagnoses for all children in the study base. However, we found a similar result on the association of pollutant levels and ASD in the sub-population where complete information on ASD diagnoses from all registers was

available among children born since 2003. Furthermore, there was no data on maternal exposure prior to conception or on paternal exposure during the child's early life. The analyses conditioned on municipality (and birth year) to minimize risks for differences in diagnostic practices influencing the results. This may have led to some over-adjustment in relation to the air pollution effects. However, we did not observe an increased risk by exposure to either pollutant in sensitivity analysis not conditioning on municipality.

In conclusion, our results indicate that exposure to NO_x and PM₁₀ during the pre- and post-natal period is not associated with ASD. The absence of positive associations in our study may be related to comparatively low air pollutant levels.

Reference

- Allen JL, Liu X, Pelkowski S, Palmer B, Conrad K, Oberdorster G, et al. 2014. Early postnatal exposure to ultrafine particulate matter air pollution: Persistent ventriculomegaly, neurochemical disruption, and glial activation preferentially in male mice. *Environmental health perspectives* 122:939-945.
- American Psychiatric Association. 2013. Diagnostic and statistical manual of mental disorders : Dsm-5. 5th ed. Washington, D.C.:American Psychiatric Association.
- Anell A, Glenngård AH, Merkur S. 2012. Health systems in transition: Sweden-a health system review. World Health Organization, on behalf of the European Observatory on Health Systems and Policies.
- Bailey A, Le Couteur A, Gottesman I, Bolton P, Simonoff E, Yuzda E, et al. 1995. Autism as a strongly genetic disorder: Evidence from a british twin study. *Psychological medicine* 25:63-77.
- Baird G, Simonoff E, Pickles A, Chandler S, Loucas T, Meldrum D, et al. 2006. Prevalence of disorders of the autism spectrum in a population cohort of children in south thames: The special needs and autism project (snap). *Lancet* 368:210-215.
- Becerra TA, Wilhelm M, Olsen J, Cockburn M, Ritz B. 2013. Ambient air pollution and autism in los angeles county, california. *Environ Health Perspect* 121:380-386.
- Bell ML, Belanger K. 2012. Review of research on residential mobility during pregnancy: Consequences for assessment of prenatal environmental exposures. *Journal of exposure science & environmental epidemiology* 22:429-438.
- Bellander T, Berglind N, Gustavsson P, Jonson T, Nyberg F, Pershagen G, et al. 2001. Using geographic information systems to assess individual historical exposure to air pollution from traffic and house heating in stockholm. *Environmental health perspectives* 109:633-639.
- Braveman PA, Cubbin C, Egerter S, Chideya S, Marchi KS, Metzler M, et al. 2005. Socioeconomic status in health research: One size does not fit all. *JAMA* 294:2879-2888.
- Buchmayer S, Johansson S, Johansson A, Hultman CM, Sparen P, Cnattingius S. 2009. Can association between preterm birth and autism be explained by maternal or neonatal morbidity? *Pediatrics* 124:e817-825.
- Buescher AV, Cidav Z, Knapp M, Mandell DS. 2014. Costs of autism spectrum disorders in the united kingdom and the united states. *JAMA pediatrics* 168:721-728.
- Burman L, Norman M. 2013. The air quality in stockholm: Annual report 2012. Available: http://www.slb.nu/slb/rapporter/pdf8/slb2013_005.pdf [accessed Feb 07 2014].
- Centers for Disease Control and Prevention. 2014. Prevalence of autism spectrum disorder among children aged 8 years — autism and developmental disabilities monitoring network, 11 sites, united states, 2010. *Morbidity and Mortality Weekly Report* 2014; 63 (2).
- Daniels JL, Forssen U, Hultman CM, Cnattingius S, Savitz DA, Feychting M, et al. 2008. Parental psychiatric disorders associated with autism spectrum disorders in the offspring. *Pediatrics* 121:e1357-1362.
- Durant JL, Beelen R, Eeftens M, Meliefste K, Cyrus J, Heinrich J, et al. 2014. Comparison of ambient airborne pm(2).(5), pm(2).(5) absorbance and nitrogen dioxide ratios measured in 1999 and 2009 in three areas in europe. *The Science of the total environment* 487:290-298.
- Eneroth K, Johansson C, Bellander T. 2006. Comparison between measurements and calculations based on dispersion modelling (expose). 2006:12. Stockholm Environment and Health Protection Administration.
- Fell DB, Dodds L, King WD. 2004. Residential mobility during pregnancy. *Paediatric and perinatal epidemiology* 18:408-414.
- Gardener H, Spiegelman D, Buka SL. 2011. Perinatal and neonatal risk factors for autism: A comprehensive meta-analysis. *Pediatrics* 128:344-355.
- Gong T, Almqvist C, Bolte S, Lichtenstein P, Anckarsater H, Lind T, et al. 2014. Exposure to air pollution from traffic and neurodevelopmental disorders in swedish twins. *Twin Res Hum Genet* 17:553-562.

- Gruzieva O, Bellander T, Eneroth K, Kull I, Melen E, Nordling E, et al. 2012. Traffic-related air pollution and development of allergic sensitization in children during the first 8 years of life. *J Allergy Clin Immunol* 129:240-246.
- Guxens M, Ghassabian A, Gong T, Garcia-Esteban R, Porta D, Giorgis-Allemand L, et al. 2016. Air pollution exposure during pregnancy and childhood autistic traits in four european population-based cohort studies: The escape project. *Environ Health Perspect* 124:133-140.
- Hajat A, Diez-Roux AV, Adar SD, Auchincloss AH, Lovasi GS, O'Neill MS, et al. 2013. Air pollution and individual and neighborhood socioeconomic status: Evidence from the multi-ethnic study of atherosclerosis (mesa). *Environ Health Perspect* 121:1325-1333.
- Hoekstra RA, Happe F, Baron-Cohen S, Ronald A. 2009. Association between extreme autistic traits and intellectual disability: Insights from a general population twin study. *The British journal of psychiatry : the journal of mental science* 195:531-536.
- Hoekstra RA, Happe F, Baron-Cohen S, Ronald A. 2010. Limited genetic covariance between autistic traits and intelligence: Findings from a longitudinal twin study. *American journal of medical genetics Part B, Neuropsychiatric genetics : the official publication of the International Society of Psychiatric Genetics* 153b:994-1007.
- Hultman CM, Sparen P, Cnattingius S. 2002. Perinatal risk factors for infantile autism. *Epidemiology (Cambridge, Mass)* 13:417-423.
- Idring S, Rai D, Dal H, Dalman C, Sturm H, Zander E, et al. 2012. Autism spectrum disorders in the stockholm youth cohort: Design, prevalence and validity. *PLoS One* 7:e41280.
- Idring S, Lundberg M, Sturm H, Dalman C, Gumpert C, Rai D, et al. 2015. Changes in prevalence of autism spectrum disorders in 2001–2011: Findings from the stockholm youth cohort. *Journal of autism and developmental disorders* 45:1766-1773.
- Johansson C, Andersson C, Bergström R, Krecl P. 2008. Exposure to particles due to local and non-local sources in stockholm - estimates based on modelling and measurements 1997 - 2006. 175. Department of Applied Environmental Science, Stockholm University, Stockholm Sweden.
- Jung CR, Lin YT, Hwang BF. 2013. Air pollution and newly diagnostic autism spectrum disorders: A population-based cohort study in taiwan. *PLoS One* 8:e75510.
- Kalkbrenner AE, Daniels JL, Chen JC, Poole C, Emch M, Morrissey J. 2010. Perinatal exposure to hazardous air pollutants and autism spectrum disorders at age 8. *Epidemiology* 21:631-641.
- Kalkbrenner AE, Windham GC, Serre ML, Akita Y, Wang X, Hoffman K, et al. 2015. Particulate matter exposure, prenatal and postnatal windows of susceptibility, and autism spectrum disorders. *Epidemiology* 26:30-42.
- Lauritsen MB, Pedersen CB, Mortensen PB. 2004. The incidence and prevalence of pervasive developmental disorders: A danish population-based study. *Psychological medicine* 34:1339-1346.
- Lee BK, Magnusson C, Gardner RM, Blomstrom A, Newschaffer CJ, Burstyn I, et al. 2015. Maternal hospitalization with infection during pregnancy and risk of autism spectrum disorders. *Brain Behav Immun* 44:100-105.
- Lichtenstein P, Carlstrom E, Rastam M, Gillberg C, Anckarsater H. 2010. The genetics of autism spectrum disorders and related neuropsychiatric disorders in childhood. *The American journal of psychiatry* 167:1357-1363.
- Ludvigsson JF, Otterblad-Olausson P, Pettersson BU, Ekbohm A. 2009. The swedish personal identity number: Possibilities and pitfalls in healthcare and medical research. *European journal of epidemiology* 24:659-667.
- Lundstrom S, Reichenberg A, Melke J, Rastam M, Kerekes N, Lichtenstein P, et al. 2015. Autism spectrum disorders and coexisting disorders in a nationwide swedish twin study. *Journal of child psychology and psychiatry, and allied disciplines* 56:702-710.
- Magnusson C, Rai D, Goodman A, Lundberg M, Idring S, Svensson A, et al. 2012. Migration and autism spectrum disorder: Population-based study. *Br J Psychiatry* 201:109-115.
- Melen E, Nyberg F, Lindgren CM, Berglund N, Zucchelli M, Nordling E, et al. 2008. Interactions between glutathione s-transferase p1, tumor necrosis factor, and traffic-related air pollution for development of childhood allergic disease. *Environmental health perspectives* 116:1077-1084.

- Miller A, Siffel C, Correa A. 2010. Residential mobility during pregnancy: Patterns and correlates. *Matern Child Health J* 14:625-634.
- Nassar N, Dixon G, Bourke J, Bower C, Glasson E, de Klerk N, et al. 2009. Autism spectrum disorders in young children: Effect of changes in diagnostic practices. *International journal of epidemiology* 38:1245-1254.
- Nordling E, Berglind N, Melen E, Emenius G, Hallberg J, Nyberg F, et al. 2008. Traffic-related air pollution and childhood respiratory symptoms, function and allergies. *Epidemiology (Cambridge, Mass)* 19:401-408.
- Palmer RF, Blanchard S, Wood R. 2009. Proximity to point sources of environmental mercury release as a predictor of autism prevalence. *Health Place* 15:18-24.
- Rai D, Lewis G, Lundberg M, Araya R, Svensson A, Dalman C, et al. 2012. Parental socioeconomic status and risk of offspring autism spectrum disorders in a swedish population-based study. *J Am Acad Child Adolesc Psychiatry* 51:467-476 e466.
- Rai D, Lee BK, Dalman C, Golding J, Lewis G, Magnusson C. 2013. Parental depression, maternal antidepressant use during pregnancy, and risk of autism spectrum disorders: Population based case-control study. *BMJ* 346:f2059.
- Raz R, Roberts AL, Lyall K, Hart JE, Just AC, Laden F, et al. 2014. Autism spectrum disorder and particulate matter air pollution before, during, and after pregnancy: A nested case-control analysis within the nurses' health study ii cohort. *Environmental health perspectives*.
- Ritvo ER, Jorde LB, Mason-Brothers A, Freeman BJ, Pingree C, Jones MB, et al. 1989. The ucla-university of utah epidemiologic survey of autism: Recurrence risk estimates and genetic counseling. *The American journal of psychiatry* 146:1032-1036.
- Roberts AL, Lyall K, Hart JE, Laden F, Just AC, Bobb JF, et al. 2013. Perinatal air pollutant exposures and autism spectrum disorder in the children of nurses' health study ii participants. *Environmental health perspectives* 121:978-984.
- Roberts AL, Lyall K, Rich-Edwards JW, Ascherio A, Weisskopf MG. 2013. Maternal exposure to childhood abuse is associated with elevated risk of autism. *JAMA psychiatry* 70:508-515.
- Ronald A, Larsson H, Anckarsater H, Lichtenstein P. 2011. A twin study of autism symptoms in sweden. *Molecular psychiatry* 16:1039-1047.
- Sandin S, Lichtenstein P, Kuja-Halkola R, Larsson H, Hultman CM, Reichenberg A. 2014. The familial risk of autism. *JAMA* 311:1770-1777.
- Sariaslan A, Långström N, D'Onofrio B, Hallqvist J, Franck J, Lichtenstein P. 2013. The impact of neighbourhood deprivation on adolescent violent criminality and substance misuse: A longitudinal, quasi-experimental study of the total swedish population. *International Journal of Epidemiology* 42:1057-1066.
- Schultz ES, Gruzieva O, Bellander T, Bottai M, Hallberg J, Kull I, et al. 2012. Traffic-related air pollution and lung function in children at 8 years of age: A birth cohort study. *Am J Respir Crit Care Med* 186:1286-1291.
- The Swedish National Board of Health and Welfare. 2003. The swedish medical birth register-a summary of content and quality. 2003-112-3.
- Thomas P, Zahorodny W, Peng B, Kim S, Jani N, Halperin W, et al. 2012. The association of autism diagnosis with socioeconomic status. *Autism* 16:201-213.
- Tulloch AD, Fearon P, Fahy T, David A. 2010. Residential mobility among individuals with severe mental illness: Cohort study of uk700 participants. *Social psychiatry and psychiatric epidemiology* 45:767-777.
- Windham GC, Zhang L, Gunier R, Croen LA, Grether JK. 2006. Autism spectrum disorders in relation to distribution of hazardous air pollutants in the san francisco bay area. *Environ Health Perspect* 114:1438-1444.
- Volk HE, Hertz-Picciotto I, Delwiche L, Lurmann F, McConnell R. 2011. Residential proximity to freeways and autism in the charge study. *Environmental health perspectives* 119:873-877.
- Volk HE, Lurmann F, Penfold B, Hertz-Picciotto I, McConnell R. 2013. Traffic-related air pollution, particulate matter, and autism. *JAMA Psychiatry* 70:71-77.

Table 1. Characteristics of 5,136 ASD and 18,237 randomly-selected control subjects born in Stockholm, Sweden between 1993 and 2007.

| | Cases (n=5,136) | Controls (n=18,237) | p |
|---|-----------------|---------------------|---------|
| Child characteristics | | | |
| Age at end of follow-up | 13.0±4.1 | 11.5±4.5 | <0.0001 |
| Gender | | | |
| Male | 3,760 (73.2) | 9,408 (51.6) | <0.0001 |
| Female | 1,376 (26.8) | 8,829 (48.4) | |
| Birth year | | | |
| 1993-1995 | 1,321 (25.7) | 3,527 (19.3) | <0.0001 |
| 1996-1998 | 1,184 (23.1) | 3,050 (16.7) | |
| 1999-2001 | 1,088 (21.2) | 3,200 (17.5) | |
| 2002-2004 | 920 (17.9) | 4,014 (22.0) | |
| 2005-2007 | 623 (12.1) | 4,446 (24.4) | |
| Sib order | | | |
| First child | 2,402 (46.8) | 7,785 (42.7) | <0.0001 |
| Not first child | 2,734 (53.2) | 10,452 (57.3) | |
| Birth weight (grams) | | | |
| <2,500 | 259 (5.0) | 487 (2.6) | <0.0001 |
| 2,500-3,000 | 576 (11.2) | 1,777 (9.7) | |
| 3,001-3,500 | 1,507 (29.3) | 5,845 (32.1) | |
| 3,501-4,000 | 1,742 (33.9) | 6,561 (36.0) | |
| >4000 | 1,015 (19.8) | 3,488 (19.1) | |
| Congenital malformations | | | |
| No | 4,851 (94.7) | 16,565 (90.8) | <0.0001 |
| Yes | 274 (5.3) | 530 (2.9) | |
| Intellectual disability | | | |
| No | 4,223 (82.2) | 18,119 (99.4) | <0.0001 |
| Yes | 913 (17.8) | 118 (0.6) | |
| Family characteristics | | | |
| Maternal age at child birth | 30.5±5.3 | 30.9±4.9 | <0.0001 |
| Paternal age at child birth | 33.3±6.6 | 33.6±6.1 | <0.0001 |
| Season of conception | | | |
| Spring (Mar, Apr, May) | 1,300 (25.3) | 5,008 (27.5) | 0.0042 |
| Summer (Jun, Jul, Aug) | 1,340 (26.1) | 4,784 (26.2) | |
| Autumn (Sep, Oct, Nov) | 1,288 (25.1) | 4,230 (23.2) | |
| Winter (Dec, Jan, Feb) | 1,208 (23.5) | 4,215 (23.1) | |
| Residential mobility during pregnancy | | | 0.003 |
| Non-movers | 4,025 (78.6) | 14,646 (80.5) | |
| Movers | 1,095 (21.4) | 3,551 (19.5) | |
| Parents' birth countries | | | |
| Both from Sweden | 3,512 (68.4) | 12,796 (70.2) | 0.0013 |
| One from other countries | 889 (17.3) | 2,780 (15.2) | |
| Both from other countries | 713 (13.9) | 2,610 (14.1) | |
| Disposable income within household at child birth (quintiles) | | | |
| Lowest | 586 (11.4) | 1,860 (10.2) | <0.0001 |
| Lower middle | 1,193 (23.2) | 3,509 (19.2) | |
| Middle | 1,266 (24.7) | 4,229 (23.2) | |
| Upper middle | 1,142 (22.2) | 4,187 (23.0) | |
| Highest | 949 (18.5) | 4,452 (24.4) | |
| Neighborhood deprivation index at child birth | 0.15±1.12 | -0.02±1.12 | |
| Low (-7.78 to -0.55) | 1,463 (28.5) | 6,249 (34.3) | |
| Medium (-0.55 to 0.19) | 1,643 (32.0) | 6,301 (34.6) | <0.0001 |
| High (0.19 to 4.06) | 2,030 (39.5) | 5,687 (31.2) | |
| Mother's education at child birth | | | |
| Low (≤9 years) | 808 (15.7) | 2,230 (12.2) | <0.0001 |
| Medium (10-12 years) | 2,394 (46.6) | 7,879 (43.2) | |
| High (>12 years) | 1,921 (37.4) | 8,076 (44.3) | |

| | | | |
|---|--------------|---------------|---------|
| Father's education at child birth | | | |
| Low (≤ 9 years) | 856 (16.7) | 2,486 (13.6) | <0.0001 |
| Medium (10-12 years) | 2,316 (45.1) | 7,781 (42.7) | |
| High (>12 years) | 1,890 (36.8) | 7,766 (42.6) | |
| Highest education in family at child birth | | | |
| At least one parent having >12 years of education | 2,571 (50.1) | 10,442 (57.3) | <0.0001 |
| Neither parents having 12 years of education | 2,564 (49.9) | 7,789 (42.7) | |
| Mother's employment during pregnancy | | | |
| Employed | 3,512 (68.4) | 13,888 (76.2) | <0.0001 |
| Unemployed with tasks | 602 (11.7) | 1,549 (8.5) | |
| Unemployed without tasks | 1,020 (19.9) | 2,799 (15.4) | |
| Father's employment during pregnancy | | | |
| Employed | 4,200 (81.8) | 15,751 (86.4) | <0.0001 |
| Unemployed with tasks | 391 (7.6) | 1,019 (5.6) | |
| Unemployed without tasks | 436 (8.5) | 1,216 (6.7) | |
| Mother's marital status at child birth | | | |
| Married/Cohabiting | 4,289 (83.5) | 16,072 (88.1) | <0.0001 |
| Single/Other situations | 830 (16.2) | 2,128 (11.7) | |
| Parental psychiatric history (F-diagnoses) | | | |
| Father diagnosed | 578 (11.3) | 1,925 (10.6) | <0.0001 |
| Mother diagnosed | 1,622 (31.6) | 4,266 (23.4) | |
| Both parents diagnosed | 868 (16.9) | 1,552 (8.5) | |
| Parental history of schizophrenia | | | |
| Father diagnosed | 18 (0.4) | 25 (0.1) | 0.0063 |
| Mother diagnosed | 9 (0.2) | 28 (0.2) | |
| Parental history of bipolar disorders | | | |
| Father diagnosed | 43 (0.8) | 90 (0.5) | <0.0001 |
| Mother diagnosed | 98 (1.9) | 170 (0.9) | |
| Parental history of intellectual disability | | | |
| Father diagnosed | 9 (0.2) | 4 (0.0) | <0.0001 |
| Mother diagnosed | 20 (0.4) | 14 (0.1) | |
| Parental history of non-affective psychosis | | | |
| Father diagnosed | 83 (1.6) | 146 (0.8) | <0.0001 |
| Mother diagnosed | 39 (0.8) | 136 (0.8) | |
| Gestational age (weeks) | | | |
| <33 | 79 (1.5) | 129 (0.7) | <0.0001 |
| 33-36 | 270 (5.3) | 661 (3.6) | |
| 37-42 | 4,370 (85.1) | 16,028 (87.9) | |
| >42 | 406 (7.9) | 1,393 (7.6) | |
| Pre-eclampsia | | | |
| No | 4,949 (96.4) | 17,781 (97.5) | <0.0001 |
| Yes | 187 (3.6) | 456 (2.5) | |
| Pre-gestational and gestational diabetes | | | |
| No | 5,037 (98.1) | 18,072 (99.1) | <0.0001 |
| Yes | 99 (1.9) | 165 (0.9) | |
| Placental abruption | | | |
| No | 5,107 (99.4) | 18,172 (99.6) | 0.0373 |
| Yes | 29 (0.6) | 65 (0.4) | |
| Premature rupture of the membranes | | | |
| No | 5,033 (98.0) | 18,009 (98.8) | <0.0001 |
| Yes | 103 (2.0) | 228 (1.3) | |
| Maternal smoking during pregnancy | | | |
| No | 3,910 (76.1) | 14,476 (79.4) | <0.0001 |
| 1-10 cigarettes/day | 383 (7.5) | 1,075 (5.9) | |
| >10 cigarettes/day | 241 (4.7) | 465 (2.6) | |
| Missing | 602 (11.7) | 2,221 (12.2) | |

Note: Missing with less than 2% was not presented.

Table 2. Risk of ASD overall by quartiles of modeled pollutants' levels at different time windows among 5,136 ASD cases and 18,237 controls born in Stockholm, Sweden between 1993 and 2007.

| Time windows and pollutants | no. of cases/controls | Models A [†] OR (95% CI) | Models B [‡] OR (95% CI) | Models C ^l OR (95% CI) |
|-----------------------------|-----------------------|--------------------------------------|--------------------------------------|--------------------------------------|
| Entire pregnancy | 5,112/18,192 | | | |
| NO _x | | | | |
| 2nd quartile | 1,340/4,486 | 1.01 (0.91, 1.13) | 0.98 (0.88, 1.10) | 0.96 (0.85, 1.07) |
| 3rd quartile | 1,285/4,541 | 0.94 (0.83, 1.06) | 0.91 (0.80, 1.03) | 0.88 (0.78, 1.00) |
| 4th quartile | 1,279/4,547 | 0.94 (0.82, 1.09) | 0.91 (0.78, 1.05) | 0.89 (0.77, 1.03) |
| PM ₁₀ | | | | |
| 2nd quartile | 1,399/4,427 | 1.03 (0.92, 1.14) | 1.01 (0.91, 1.12) | 0.98 (0.88, 1.09) |
| 3rd quartile | 1,247/4,579 | 0.92 (0.81, 1.04) | 0.90 (0.79, 1.02) | 0.87 (0.77, 0.99) |
| 4th quartile | 1,140/4,686 | 0.93 (0.81, 1.06) | 0.88 (0.77, 1.02) | 0.86 (0.75, 0.99) |
| First year of life | 5,121/18,225 | | | |
| NO _x | | | | |
| 2nd quartile | 1,321/4,515 | 1.05 (0.94, 1.18) | 1.02 (0.91, 1.15) | 0.98 (0.87, 1.10) |
| 3rd quartile | 1,328/4,509 | 1.04 (0.91, 1.19) | 1.00 (0.87, 1.14) | 0.95 (0.83, 1.09) |
| 4th quartile | 1,297/4,539 | 1.06 (0.90, 1.25) | 1.00 (0.85, 1.18) | 0.96 (0.81, 1.13) |
| PM ₁₀ | | | | |
| 2nd quartile | 1,398/4,439 | 1.10 (0.99, 1.23) | 1.08 (0.97, 1.21) | 1.03 (0.92, 1.16) |
| 3rd quartile | 1,280/4,557 | 1.03 (0.90, 1.17) | 0.98 (0.86, 1.12) | 0.93 (0.82, 1.07) |
| 4th quartile | 1,160/4,676 | 1.05 (0.90, 1.22) | 0.97 (0.83, 1.13) | 0.92 (0.78, 1.07) |

[†] Models were conditioned on calendar year of birth and municipality of birth, as well as adjusted for gender, birth month.

[‡] Models were conditioned on calendar year of birth and municipality of birth, as well as adjusted for gender, birth month, birth order, parents' birth countries, mother's marital status, mother's education, father's education, mother's employment, father's employment, and disposable income within household.

^l Models were conditioned on calendar year of birth and municipality of birth, as well as adjusted for gender, birth month, birth order, parents' birth countries, mother's marital status, mother's education, father's education, mother's employment, father's employment, disposable income within household, and neighborhood deprivation.

Figure legends

Figure 1 Box plot describing the distribution of NO_x (white) and PM₁₀ (grey) concentrations (μg/m³) from local traffic in study population during mother's pregnancy and first year after child birth. The box and whiskers denotes the 5th, 25th, 50th, 75th, 95th percentile and outlier values of pollutants' distributions.

Figure 2 Odds ratios and 95% confidence intervals for ASD overall, ASD with or without ID by residential address-based NO_x (per 10 μg/m³ increase) and PM₁₀ (per 20 μg/m³ increase) levels during mother's pregnancy and child's first year of life. All models were conditioned on calendar year of birth and municipality of birth, as well as adjusted for gender, birth month, sib order, maternal age, paternal age, mother's marital status, parents' birth countries, mother's education, father's education, mother's employment, father's employment, disposable income within household, and neighborhood deprivation.

Figure 3 Odds ratios and 95% confidence intervals ASD overall by residential address-based NO_x (per 10 μg/m³ increase) and PM₁₀ (per 20 μg/m³ increase) levels during mother's pregnancy and child's first year of life in stratified samples. All models were conditioned on calendar year of birth and municipality of birth, as well as adjusted for gender, birth month, sib order, maternal age, paternal age, mother's marital status, parents' birth countries, mother's education, father's education, mother's employment, father's employment, disposable income within household, and neighborhood deprivation as covariates.

Figure 1.




Figure 2.





Figure 3.



Note: The p-value was presented based on the Wald test for the indicated interaction term.