

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

### **Supplemental Material**

#### **Prenatal Exposure to Perfluoroalkyl Substances and Cardiometabolic Risk in Children from the Spanish INMA Birth Cohort Study**

Cyntia B. Manzano-Salgado, Maribel Casas, Maria-Jose Lopez-Espinosa, Ferran Ballester, Carmen Iñiguez, David Martinez, Dora Romaguera, Silvia Fernández-Barrés, Loreto Santa-Marina, Mikel Basterretxea, Thomas Schettgen, Damaskini Valvi, Jesus Vioque, Jordi Sunyer, Martine Vrijheid

#### **Table of Contents**

**Table S1.** Summary of maternal PFAS concentrations and outcomes by region of residence in our study.

**Table S2.** Details of the imputation modeling.

**Table S3.** Example of summary of characteristics in the observed and the imputed datasets corresponding to the 4-year-old population from the region of Sabadell.

**Table S4.** Summary of characteristics in the excluded and the included samples in our study.

**Table S5.** Associations between maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and anthropometric binary outcomes during childhood.

**Table S6.** Unadjusted associations between maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and cardio-metabolic components during childhood.

**Table S7.** Complete case analysis for the associations between maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and cardio-metabolic components during childhood.

**Figure S1.** Associations between quartiles of maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and anthropometric measurements, and blood pressure at age 7 years (n=1086).

**Figure S2.** Adjusted associations between maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and weight gain, and BMI z-scores by breastfeeding duration of the index child.

**Table S8.** Associations between maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and weight gain, and BMI z-scores after adjustment for maternal glomerular filtration rate.

**Table S9.** Associations between maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and weight gain, and BMI z-scores after adjustment for maternal albumin level.

**Table S10.** Associations between maternal PFAS concentrations ( $\log_2$ -transformed, in ng/mL) and continuous anthropometric outcomes and blood pressure in children not born by cesarean section.