

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Long-Term Exposure to Wind Turbine Noise and Risk for Myocardial Infarction and Stroke: A Nationwide Cohort Study

Aslak Harbo Poulsen, Ole Raaschou-Nielsen, Alfredo Peña, Andrea N. Hahmann, Rikke Baastrup Nordsborg, Matthias Ketznel, Jørgen Brandt, and Mette Sørensen

Table of Contents

Table S1. Characteristics of the populations for study of MI and stroke, respectively, at start of follow-up according to residential low frequency exposure to indoor wind turbine noise calculated as mean exposure during the preceding year.

Table S2. Associations between mean 5-year exposure to residential outdoor and indoor low frequency wind turbine noise and risk of ischemic stroke.

Table S3. Analyses of associations between 5-year exposure to outdoor wind turbine noise and risk of stroke in different sub-populations.

Table S4. Analyses of associations between 5-year exposure to indoor low frequency wind turbine noise and risk of stroke in different sub-populations.

Table S5. Analyses of associations between 5-year exposure to outdoor wind turbine noise and risk of MI and stroke according to sex and age.

Table S6. Analyses of associations between 5-year exposure to indoor low frequency wind turbine noise and risk of MI and stroke according to sex and age.