

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Independent and Combined Effects of Heatwaves and PM_{2.5} on Preterm Birth in Guangzhou, China: A Survival Analysis

Qiong Wang, Bing Li, Tarik Benmarhnia, Shakoor Hajat, Meng Ren, Tao Liu, Luke D. Knibbs, Huanhuan Zhang, Junzhe Bao, Yawei Zhang, Qingguo Zhao, and Cunrui Huang

Table of Contents

Table S1. Daily air pollutants and meteorological factors in the warm season (May-October) in Guangzhou, China, 2015-2017.

Table S2. Air pollutants and meteorological factors in the last gestational week before delivery.

Table S3. PM_{2.5} and mean temperature during different gestational time periods and the correlations.

Table S4. PM_{2.5} in the last gestational week before delivery by heatwave exposure.

Table S5. Hazard ratios of preterm birth associated with PM_{2.5} exposure in the last gestational week in Guangzhou, China, 2015-2017.

Table S6. Hazard ratios of overall preterm birth associated with heatwave exposure in the last gestational week before delivery with adjustment for PM_{2.5} exposures during weeks 20-28.

Table S7. Hazard ratios of preterm birth associated with heatwave exposure in the last gestational week before delivery, stratified by delivery method.

Table S8. Hazard ratios of preterm birth associated with PM_{2.5} exposures in the last gestational week before delivery.

Figure S1. Locations of meteorological and air quality monitoring stations.