Environ Health Perspect

DOI: 10.1289/EHP5975

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Early-Life Environmental Exposures and Childhood Obesity: An Exposome-Wide Approach

Martine Vrijheid, Serena Fossati, Léa Maitre, Sandra Márquez, Theano Roumeliotaki, Lydiane Agier, Sandra Andrusaityte, Solène Cadiou, Maribel Casas, Montserrat de Castro, Audrius Dedele, David Donaire-Gonzalez, Regina Grazuleviciene, Line S. Haug, Rosemary McEachan, Helle Margrete Meltzer, Eleni Papadopouplou, Oliver Robinson, Amrit K. Sakhi, Valerie Siroux, Jordi Sunyer, Per E. Schwarze, Ibon Tamayo-Uria, Jose Urquiza, Marina Vafeiadi, Antonia Valentin, Charline Warembourg, John Wright, Mark J. Nieuwenhuijsen, Cathrine Thomsen, Xavier Basagaña, Rémy Slama, and Leda Chatzi

Table of Contents

Table S1. Exposure data sources for urban environment exposures.

Table S2. Summary of the indoor air pollution prediction models. P-values of multivariate and bivariate models for variables included in the final prediction model.

Table S3. Biological matrices used for biomarker analysis of maternal and child samples.

Table S4. Number of maternal samples analysed as part of HELIX and in already existing data collections.

Table S5. Description of the study population by cohort.

Table S6. Childhood exposure confounder-only model for zBMI and overweight/obesity (N=1301).

Table S7. Analysis of the association between prenatal and childhood exposures, and child zBMI score (N=1301). Single exposure (ExWAS) model and multi-exposure (DSA) model. Shown are all exposure variables with uncorrected ExWAS p-value<0.05 or selected in 10% or more of DSA runs; variables are ordered by increasing p-value in the ExWAS. ExWAS results for all exposures are shown in Table S17.

Table S8. Analysis of the association between prenatal and childhood exposures, and overweight/obesity status (vs normal weight. N overweight/obesity=374, N normal weight=927). Single exposure (ExWAS) model and multi-exposure (DSA) model. Shown are all exposure variables with uncorrected ExWAS p-value<0.05 or selected in 10% or more of DSA runs; variables are ordered by increasing p-value in the ExWAS.

Table S9. Analysis of the association between prenatal and childhood exposures, and waist circumference z-score (N=1297). Single exposure (ExWAS) model and multi-exposure (DSA) model. Shown are all exposure variables with uncorrected ExWAS p-value<0.05 or selected in 10% or more of DSA runs; variables are ordered by increasing p-value in the ExWAS.

Table S10. Analysis of the association between prenatal and childhood exposures, and skinfolds z-score (N=1288). Single exposure (ExWAS) model and multi-exposure (DSA) model. Shown are all exposure variables with uncorrected ExWAS p-value<0.05 or selected in 10% or more of DSA runs; variables are ordered by increasing p-value in the ExWAS.

Table S11. Analysis of the association between prenatal and childhood exposures, and percentage fatmass z-score (N=1290). Single exposure (ExWAS) model and multi-exposure (DSA) model. Shown are all exposure variables with uncorrected ExWAS p-value<0.05 or selected in 10% or more of DSA runs; variables are ordered by increasing p-value in the ExWAS.

Table S12. Sensitivity analysis - sex stratification. Association between prenatal and childhood exposures and zBMI score stratified by girls and boys; DSA multi-exposure model.

Table S13. Sensitivity analysis – Maternal education stratification. Association between prenatal and childhood exposures and zBMI score stratified by maternal educational category; DSA multi-exposure model.

Table S14. Sensitivity analysis – adjusting for additional maternal lifestyle factors. Association between maternal smoking during pregnancy and zBMI score.

Table S15. Sensitivity analysis – adjusting for additional childhood social and lifestyle factors. Association between *childhood* exposures and zBMI score; DSA multi-exposure model.

Table S16. Sensitivity analysis – Association between childhood exposures and zBMI score adjusted for maternal smoking during pregnancy; final multi-exposure model. DSA multi-exposure model.

Table S17. All associations between pregnancy and childhood exposures, and child zBMI score (N=1301). Single exposure (ExWAS) model. Variables ordered by exposure group.

Figure S1. Direct Acyclic Graph design to identify potential confounders in the association between the prenatal exposome and child obesity outcomes.

Figure S2. Direct Acyclic Graph design to identify potential confounders in the association between the childhood exposome and child obesity outcomes.

Figure S3. Association between prenatal and childhood exposures and waist circumference z-score in single-exposure ExWAS model - Volcano plot showing significance (p-value) against beta coefficient.

Figure S4. Association between prenatal and childhood exposures and skinfold z-score in single-exposure ExWAS model - Volcano plot showing significance (p-value) against beta coefficient.

Figure S5. Association between prenatal and childhood exposures and percentage fatmass z-score in single-exposure ExWAS model - Volcano plot showing significance (p-value) against beta coefficient.

Figure S6. Forest plots for association between prenatal and childhood exposures and zBMI score in multi-exposure model – country specific estimates and I^2 statistic.

Additional File- Excel Document