

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Bisphenol S and Epidermal Growth Factor Receptor Signaling in Human Placental Cytotrophoblasts

Elvis Ticiani, Jeremy Gingrich, Yong Pu, Mathew Vettathu, Jacquelyn Davis, Denny Martin, Margaret G. Petroff, and Almudena Veiga-Lopez

Table of Contents

Table S1. Primer sequences.

Figure S1. Identification and purity of human cytotrophoblast (hCTB) cells. **(A)** Immunofluorescence stain of hCTB cell positive marker cytokeratin 7 (CKT7), desmoplakin (DSP), and nuclear DAPI counterstain after a 24 h culture. **(B)** Flow cytometry analysis of hCTB isolates measuring the expression of CKT7. hCTBs were targeted with a direct method using an Alexa Fluor 488-conjugated anti-CKT7 antibody. Fluorescence was measured in channel FL1 and plotted in histograms. IgG control primary antibodies that lack specificity to the target were used as negative controls. Percentages of positive cells are corrected for control values. **(C)** hCTB cells immunostained against DSP and counterstained with nuclear DAPI stain (*top*). Middle and bottom panels are two negative controls (*middle*; using normal mouse serum) and (*bottom*; without primary antibody).

Figure S2. **(A)** Cell viability effect of BPS (ng/ml) on human cytotrophoblast (hCTB) using a MTT assay after 96 hours exposure. **(B)** Cell viability effect of BPS (1,000 ng/ml) and EGF (10 ng/ml) on human cytotrophoblasts exposure using an MTT assay after 96 hours exposure. Asterisks denote treatments whose results were significantly different from the control ($P < 0.05$). $N = 5$ technical replicates.

Table S1. Primer sequences

Gene	Primers	Accession	Length (bp)
<i>CCND1</i> - F	CAGATCATCCGCAAACACGC	NM_053056.3	143
<i>CCND1</i> - R	AAGTTGTTGGGGCTCCTCAG		
<i>PTGS2</i> - F	AAGTGCGATTGTACCCGGAC	NM_000963.4	105
<i>PTGS2</i> - R	GTGCACTGTGTTTGGAGTGG		
<i>MYC</i> - F	CACTAACATCCCACGCTCTGA	NM_001354870.1	195
<i>MYC</i> - R	AAATCATCGCAGGCGGAACA		
<i>GAPDH</i> - F	GGAAGCTCACTGGCATGGCCTTCC	NM_001357943.2	119
<i>GAPDH</i> - R	GCCTGCTTCACCACCTTCTTG		




Figure S1. Identification and purity of human cytotrophoblast (hCTB) cells. (A) Immunofluorescence stain of hCTB cell positive marker cyokeratin 7 (CKT7), desmoplakin (DSP), and nuclear DAPI counterstain after a 24 h culture. (B) Flow cytometry analysis of hCTB isolates measuring the expression of CKT7. hCTBs were targeted with a direct method using an Alexa Fluor 488-conjugated anti-CKT7 antibody. Fluorescence was measured in channel FL1 and plotted in histograms. IgG control primary antibodies that lack specificity to the target were used as negative controls. Percentages of positive cells are corrected for control values. (C) hCTB cells immunostained against DSP and counterstained with nuclear DAPI stain (*top*). Middle and bottom panels are two negative controls (*middle*; using normal mouse serum) and (*bottom*; without primary antibody).




Figure S2. (A) Cell viability effect of BPS (ng/ml) on human cytotrophoblast (hCTB) using a MTT assay after 96 hours exposure. (B) Cell viability effect of BPS (1,000 ng/ml) and EGF (10 ng/ml) on human cytotrophoblasts exposure using an MTT assay after 96 hours exposure. Asterisks denote treatments whose results were significantly different from the control ($P < 0.05$). $N = 5$ technical replicates.