

Air Pollution and Suicide: Exploring a Potential Risk Factor

Nate Seltenrich

<https://doi.org/10.1289/EHP3901>

Could air pollution be a trigger for suicide? Researchers first began asking this question less than a decade ago.^{1,2,3} Accumulated evidence from around the world now suggests there may well be a connection,^{4,5,6} although the nature of such a connection is still unknown. The authors of a study in *Environmental Health Perspectives* add to the evidence for this link, drawing upon a robust data set of pollution and suicide figures.⁷

The researchers examined the relationship between daily suicide deaths and daily mean levels of nitrogen dioxide (NO₂), sulfur dioxide (SO₂), and categories of particulate matter (PM₁₀, PM_{10–2.5}, PM_{2.5}) in 10 large Northeast Asian cities. The data covered one to three decades, depending on the country. The team controlled for variables such as hours of daylight, day of week, and ambient temperature, which can potentially affect the risk of suicide.^{8,9,10}


On a city-by-city basis, higher levels of air pollution were not always associated with higher suicide risk; in some cities, the association was even reversed, with increases in air pollution associated with lower risks of suicide. But when up to 30 years of information for PM₁₀, NO₂, and SO₂ was combined across all 10 cities, higher average exposures on the same day and over the previous 1–3 days were associated with a higher daily suicide

risk. Combined estimates for PM₁₀ and PM_{10–2.5} across three cities with two to eight years of data also suggested an increased risk of suicide with higher exposures. However, these estimates were less precise, particularly for PM_{2.5}.

The estimated increases in suicide risk were small but consistent. For example, each 4.3-ppb increase in average daily exposure to SO₂ was associated with a 2.0% increase in estimated suicide risk on the same day, while each increase of 36.4 mg/m³ in PM₁₀ was associated with a 1.6% increase in estimated risk.

“Previous studies have considered [data for] maybe a decade or so, but having up to thirty years is a unique contribution,” says University of Utah professor of psychiatry Amanda Bakian, who was not affiliated with the study. “There’s growing evidence to suggest an association between ambient air pollution and suicide risk in diverse populations from around the world.”

Japan, South Korea, and Taiwan share more than just the waters of the East China Sea. They all have above-average suicide rates, with South Korea ranking fourth worldwide in 2016 with 26.9 deaths per 100,000 people and Japan fourteenth with 18.5.¹¹ Taiwan’s rate of 16 per 100,000 in 2016 also significantly exceeded



Hikers scale Geumjeongsan, a mountain overlooking the city of Busan, South Korea. In 2015, South Korea had the seventh highest average PM_{2.5} levels of all developed countries and the third highest levels in East Asia, behind China and North Korea.¹⁴ Image: © MiriamPolito/iStock.

the global average of 10.6.¹² Worldwide, roughly 800,000 people die from suicide every year.¹³

One major unanswered question is exactly how specific pollutants, or air pollution in general, might influence suicide risk. The young line of inquiry has yet to provide any answers, although some studies have suggested that neuroinflammation may be involved.^{4,6} The authors note that suicide is a complex behavior linked to a number of psychosocial factors. Geographical differences such as cultural backgrounds, socioeconomic factors, and sources and components of air pollution all deserve consideration, they write.

“From my perspective, the broader take-home message relates to how we think about preventing suicide,” says Sunnybrook Research Institute’s Mark Sinyor, a psychiatrist and expert in mood disorders and suicide prevention, who was not affiliated with the study. “Any effort to make an enduring dent in suicide rates must address broader social problems and, as the evidence increasingly suggests, environmental problems such as air pollution as well. That may seem daunting, but at least there is a confluence of agendas—efforts to protect and improve our world are also likely to lead to fewer suicide deaths.”

Nate Seltenrich covers science and the environment from the San Francisco Bay Area. His work on subjects including energy, ecology, and environmental health has appeared in a wide variety of regional, national, and international publications.

References

1. Kim C, Jung SH, Kang DR, Kim HC, Moon KT, Hur NW, et al. 2010. Ambient particulate matter as a risk factor for suicide. *Am J Psychiatry* 167(9):1100–1107, PMID: 20634364, <https://doi.org/10.1176/appi.ajp.2010.09050706>.
2. Szyszkowicz M, Willey JB, Grafstein E, Rowe BH, Colman I. 2010. Air pollution and emergency department visits for suicide attempts in Vancouver, Canada. *Environ Health Insights* 4:79–86, PMID: 21079694, <https://doi.org/10.4137/EHI.S5662>.
3. Biermann T, Stilianakis N, Bleich S, Thürauf N, Kornhuber J, Reulbach U. 2009. The hypothesis of an impact of ozone on the occurrence of completed and attempted suicides. *Med Hypotheses* 72(3):338–341, PMID: 19027246, <https://doi.org/10.1016/j.mehy.2008.09.042>.
4. Bakian AV, Huber RS, Coon H, Gray D, Wilson P, McMahon WM, et al. 2015. Acute air pollution exposure and risk of suicide completion. *Am J Epidemiol* 181(5):295–303, PMID: 25673816, <https://doi.org/10.1093/aje/kwu341>.
5. Lin GZ, Li L, Song YF, Zhou YX, Shen SQ, Ou CQ. 2016. The impact of ambient air pollution on suicide mortality: a case-crossover study in Guangzhou, China. *Environ Health* 15(1):90, PMID: 27576574, <https://doi.org/10.1186/s12940-016-0177-1>.
6. Ng CF, Stickley A, Konishi S, Watanabe C. 2016. Ambient air pollution and suicide in Tokyo, 2001–2011. *J Affect Disord* 201:194–202, PMID: 27240312, <https://doi.org/10.1016/j.jad.2016.05.006>.
7. Kim Y, Ng CFS, Chung Y, Kim H, Honda Y, Guo YL, et al. 2018. Air pollution and suicide in 10 cities in Northeast Asia: a time-stratified case-crossover analysis. *Environ Health Perspect* 126(3):037002, PMID: 29529596, <https://doi.org/10.1289/EHP2223>.
8. Kim Y, Kim H, Honda Y, Guo YL, Chen BY, Woo JM, et al. 2016. Suicide and ambient temperature in East Asian countries: a time-stratified case-crossover analysis. *Environ Health Perspect* 124(1):75–80, PMID: 26069051, <https://doi.org/10.1289/ehp.1409392>.
9. Coimbra DG, Pereira E Silva AC, de Sousa-Rodrigues CF, Barbosa FT, de Siqueira Figueredo D, Araújo Santos JL, et al. 2016. Do suicide attempts occur more frequently in the spring too? A systematic review and rhythmic analysis. *J Affect Disord* 196:125–137, PMID: 26921865, <https://doi.org/10.1016/j.jad.2016.02.036>.
10. Sinyor M, Tse R, Pirkis J. 2017. Global trends in suicide epidemiology. *Curr Opin Psychiatry* 30(1):1–6, PMID: 27845946, <https://doi.org/10.1097/YCO.0000000000000296>.
11. WHO (World Health Organization). World Health Statistics data visualizations dashboard | Suicide [website]. Geneva, Switzerland:World Health Organization. <http://apps.who.int/gho/data/node.sdg.3-4-viz-2?lang=en> [accessed 11 May 2018].
12. Focus Taiwan News Channel. 25 percent of suicides in 2016 in Taiwan involved seniors: center. Taipei, Taiwan:The Central News Agency (23 June 2017). <http://focustaiwan.tw/news/asoc/201706230027.aspx> [accessed 11 May 2018].
13. WHO. Mental Health: Suicide Data [website]. Geneva, Switzerland:World Health Organization. http://www.who.int/mental_health/prevention/suicide/suicideprevent/en/ [accessed 11 May 2018].
14. OECD Statistics. 2017. Population exposure to PM_{2.5} in countries and regions. <https://stats.oecd.org/> [accessed 12 June 2018].