Jack Rowley
GSM Association
London, United Kingdom
E-mail: jrowley@gsm.org

References

Dolan M, Rowley J. 2009. The precautionary principle in the context of mobile phone and base station radiofrequency exposures. Environ Health Perspect 117:1329–1332. .

ICNIRP (International Commission on Non-Ionizing Radiation Protection). 1998. Guidelines for limiting exposure to time-varying electric, magnetic, and electromagnetic fields (up to 300 GHz). Health Phys 74(4):494-522.

ICNIRP (International Commission on Non-Ionizing Radiation Protection). 2009. Guidelines for limiting exposure to time-varying electric, magnetic, and electromagnetic fields (up to 300 GHz). Health Phys 97(3):257–258.

International Agency for Research on Cancer. 1987. Asbestos. IARC Monogr Eval Carcinog Risk Hum 14(suppl 7):106–116. Available: http://monographs.iarc.fr/ENG/Monographs/suppl7/Suppl7-20.pdf [accessed 8 December 2009].

Scientific Committee on Emerging and Newly Identified Health Risks. 2009. Health Effects of Exposure to EMF. Available: http://ec.europa.eu/health/ph_risk/committees/04_scenihr/docs/scenihr_o_022.pdf [accessed 15 September 2009].

Erratum

The authors report an error in Figure 2 of "Air Pollution Exposures and Circulating Biomarkers of Effect in a Susceptible Population: Clues to Potential Causal Component Mixtures and Mechanisms" [Environ Health Perspect 117:1232–1238 (2009)]. In "Materials and Methods," they noted that "four influential high outliers for IL-6 > 10 pg/mL were reset to 10 pg/mL"; however, the analysis shown in Figure 2A for interleukin (IL)-6 included these outliers without them having been reset to 10 pg/mL. In all other figures showing results for IL-6 and in Supplemental Material (doi:10.1289/ehp.0800194.S1 via http://dx.doi.org/), the outliers were correctly reset to 10 pg/mL. The IL-6 outliers are themselves valid measurements, and the results in both cases do not change the finding that the association of IL-6 with $PM_{0.25}$ (particulate matter < 0.25 µm in aerodynamic diameter) is stronger than associations with larger particle size fractions.

Choose from over 100 science and interdisciplinary lessons based on selected articles published by *EHP*, the top monthly journal in public, environmental, and occupational health.



It's healthy, it's smart, and it's free.
www.ehponline.org/science-ed-new/

International Society for Environmental Epidemiology

BEST ENVIRONMENTAL EPIDEMIOLOGY PAPER AWARD 2009

An ISEE annual award for the best environmental epidemiology paper published in a peer-reviewed journal

Award criteria include:

- **Quality (data, analysis, presentation)**
- **Originality**
- **Importance/impact**

- Authors, editors, and readers may make nominations. Only outstanding papers should be nominated.
- The nominated paper should be printed in hard copy in a scientific journal between January 1 and December 31, 2009.
- Submit nomination letters no later than **April 1, 2010** by postal mail or email, accompanied by a copy of the paper and supporting letter (maximum of 2 pages) stating why the paper should receive the award.

Address for sending nominations:

Secretariat, International Society for Environmental Epidemiology
c/o JSI Research and Training Institute, 44 Farnsworth Street, Boston, MA 02210 USA
ATTENTION: Chair, Best Environmental Epidemiology Paper Award Committee
iseepi@jsi.com

For more information visit www.iseepi.org