

**Note to Readers:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Elemental Constituents of Particulate Matter and Newborn's Size in Eight European Cohorts**

Marie Pedersen, Ulrike Gehring, Rob Beelen, Meng Wang, Lise Giorgis-Allemand, Anne-Marie Nybo Andersen, Xavier Basagaña, Claire Bernard, Marta Cirach, Francesco Forastiere, Kees de Hoogh, Regina Gražulevičienė, Olena Gruzieva, Gerard Hoek, Aleksandra Jedynska, Claudia Klümper, Ingeborg M. Kooter, Ursula Krämer, Jaakko Kukkonen, Daniela Porta, Dirkje S. Postma, Ole Raaschou-Nielsen, Lenie van Rossem, Jordi Sunyer, Mette Sørensen, Ming-Yi Tsai, Tanja G. M. Vrijkotte, Michael Wilhelm, Mark J. Nieuwenhuijsen, Göran Pershagen, Bert Brunekreef, Manolis Kogevinas, and Rémy Slama

#### **Table of Contents**

**Table S1.** Mother-child cohort data collection.

**Table S1.** Mother-child cohort data collection – cont.

**Table S2.** Variance of between- vs. within-center variations for the elements.

**Table S3.** Correlations between PM constituents ( $\text{ng}/\text{m}^3$ ).

**Table S4.** Correlations between elemental ( $\text{ng}/\text{m}^3$ ) and total mass concentration of PM ( $\mu\text{g}/\text{m}^3$ ) by cohort.

**Table S5.** Sensitivity and stratified analyses of sulfur on term LBW.

**Table S6.** Sex-stratified associations between exposure to PM constituents and term LBW.

**Table S7.** Associations between exposure to PM constituents and term LBW results from the study population with information on atmospheric pressure and pressure adjusted models.

**Table S8.** Associations between exposure to PM constituents and term LBW results from between centers and within centers models.

**Table S9.** Associations between exposure to PM constituents and birth weight results from the study population with information on atmospheric pressure and pressure adjusted models.

**Table S10.** Associations between exposure to PM constituents and birth weight results from between centers and within centers models.

**Table S11.** Sex-stratified associations between exposure to PM constituents, birth weight and head circumference.

**Table S12.** Associations between exposure to PM constituents and birth head circumference results from the study population with information on atmospheric pressure and temperature and pressure adjusted models.

**Table S13.** PM constituents and birth head circumference results from between centers and within centers models.

**Figure S1.** Distributions of PM<sub>2.5</sub>, PM<sub>10</sub>, maternal height, pre-pregnancy weight, birth weight, birth head circumference and gestational age by cohorts and for the pooled study population. The line in the middle of the box represents the median values, the ends of the box refer to the 25th and 75th percentiles and the ends of the whiskers indicate the variability outside the upper and lower quartiles (i.e., within 1.5 interquartile range of the lower quartile and upper quartile). Outliers are plotted as individual dots.