

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Genome-Wide Analysis of DNA Methylation and Cigarette Smoking in Chinese

Xiaoyan Zhu, Jun Li, Siyun Deng, Kuai Yu, Xuezhen Liu, Qifei Deng, Huizhen Sun, Xiaomin Zhang, Meian He, Huan Guo, Weihong Chen, Jing Yuan, Bing Zhang, Dan Kuang, Xiaosheng He, Yansen Bai, Xu Han, Bing Liu, Xiaoliang Li, Liangle Yang, Haijing Jiang, Yizhi Zhang, Jie Hu, Longxian Cheng, Xiaoting Luo, Wenhua Mei, Zhiming Zhou, Shunchang Sun, Liyun Zhang, Chuanyao Liu, Yanjun Guo, Zhihong Zhang, Frank B. Hu, Liming Liang, and Tangchun Wu

Table of Contents

Table S1. The 318 identified CpGs associated with cigarette smoking in the genome-wide meta-analysis ($FDR < 0.05$).

Table S2. The effect size (s.e.) of smoking on the 318 CpGs with inverse normal transformed methylation values calculated in different groups.

Table S3. The median methylation values (inter quartile range) of the 318 smoking-related CpGs in never, former and current smokers.

Table S4. Methylation-gene expression correlations of the smoking-related CpGs in SY ($n = 144$).

Table S5. Contribution rates of smoking on urinary PAHs metabolites calculated in male individuals from the WHZH cohort ($n = 206$).

Table S6. Associations between the smoking-related CpGs and urinary 2-hydroxynaphthalene levels in males from the WHZH Cohort and the Coke Oven Cohort.

Figure S1. The flowchart of the study.

References