

SUPPLEMENTAL MATERIAL:

**Hyaluronan Activation of the Nlrp3 Inflammasome Contributes to the Development of
Airway Hyperresponsiveness**

Feifei Feng^{1,2}, Zhuowei Li¹, Erin N. Potts-Kant¹, Yiming Wu², W. Michael Foster¹,
Kristi L. Williams^{1,3†}, and John W. Hollingsworth^{1,4†*}

¹Department of Medicine, Division of Pulmonary, Allergy, and Critical Care Medicine, School of Medicine, Duke University Medical Center, Durham, North Carolina, USA

²Department of Occupational and Environmental Health, College of Public Health, Zhengzhou University, Zhengzhou, Henan, China

³School of Nursing, Duke University Medical Center, Durham, North Carolina, USA

⁴Department of Immunology, School of Medicine, Duke University Medical Center, Durham, North Carolina, USA

Running Title: Hyaluronan activation of the inflammasome and environmental airways disease

Acknowledgements: This work was supported by National Institutes of Health grants ES016126, AI081672, ES020350 & ES020426 (to JWH), AI089756 (to K LW), and an unrestricted educational grant from the China Scholarship Council (to FF).

***Corresponding Author:** John W. Hollingsworth M.D., Division of Pulmonary, Allergy, and Critical Care Medicine, Department of Medicine, Duke University School of Medicine, DUMC 103004, 106 Research Drive, Durham, NC 27710, USA; Phone: (919) 684-4588, Fax: (919) 684-5266; E-mail: john.hollingsworth@duke.edu

Table of Contents

Page 3. Figure S1: RNA expression of *IL-1 β* and *IL-18* in whole lung macrophages.

Page 4. Figure S2. Level of non-inflammasome pro-inflammatory factors in BAL at 24 hours after ozone exposure.

Page 5. Figure S3. Biological response to hyaluronan and genes of the Nlrp3-inflammasome.

Page 6. Figure S4. Level of non-inflammasome pro-inflammatory factors in the BAL after HA exposure.

Supplement Figures

Figure S1.




Figure S1: RNA expression of *IL-1β* and *IL-18* in whole lung macrophages. (A/B) Whole lung macrophages were isolated after exposure to ozone and evaluated for mRNA expression of *IL-1β* and *IL-18* by real-time PCR and no differences were detected. **(C/D)** Whole lung macrophages were isolated after exposure to hyaluronan fragments and evaluated for mRNA expression of *IL-1β* and *IL-18* by real-time PCR and no differences were detected. Data presented as mean ± SEM (n=5/group).

Figure S2.




Figure S2. Level of non-inflammasome pro-inflammatory factors in BAL at 24 hours after ozone exposure. BAL IL-1 α , IL-6, monocyte chemoattractant protein-1(MCP-1), tumor necrosis factor- α (TNF- α), keratinocyte-derived chemokine (KC), IL-17, and C3a were increased with ozone exposure, and MCP-1 was totally dependent on the presence of Nlrp3-inflammasome, including caspase1, ASC and Nlrp3; IL-1 α , IL-6, TNF- α and KC were totally dependent on caspase1 and ASC and partially dependent on Nlrp3; IL-17 was partially dependent on ASC; C3a was partially dependent on Nlrp3-inflammasomes after ozone exposure. Data presented as mean \pm SEM and are representatives from two similar experiments(n=5, * p < 0.05).

Figure S3.




Figure S3. Biological response to hyaluronan and genes of the Nlrp3-inflammasome. (A)

The level of total cells in bronchoalveolar lavage (BALF) was independent of challenge to hyaluronan. **(B)** The level of BALF protein was independent of hyaluronan exposure. Data presented as mean \pm SEM (n=5/group).

Figure S4.




Figure S4. Level of non-inflammasome pro-inflammatory factors in the BAL after HA exposure. The release of BAL IL-6, MCP-1, TNF- α and KC were totally dependent on ASC and caspase1 and partially dependent on Nlrp3; IL-1 α was partially dependent on ASC. The level of IL-17 was partially dependent on caspase1 and ASC, but not dependent on Nlrp3; C3a was totally dependent on ASC and caspase1, and partially dependent on Nlrp3 after hyaluronan exposure. Data presented as mean \pm SEM and are representatives from two similar experiments (n=5, *: $p < 0.05$).