

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehponline@niehs.nih.gov](mailto:ehponline@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Geographic Differences in Persistent Organic Pollutant Levels of Yellowfin Tuna**


Sascha C.T. Nicklisch, Lindsay T. Bonito, Stuart Sandin, and Amro Hamdoun

#### **Table of Contents**


**Figure S1.** Capture locations and measured pollutant classes in this study. Shown are the 12 different locations where a total of 117 wild yellowfin tuna were caught. Numbers in parenthesis show the sample size per site. Filled circles indicate the pollutant classes that were measured at each site, color-coded for the 29 OCPs (red), 9 PBDEs (orange), and 209 PCBs (green). The world map file is available in the public domain under the Creative Commons CC0 1.0 Universal Public Domain Dedication license (<https://commons.wikimedia.org/wiki/File%3ABlankMap-World6-Equirectangular.svg>).

**Figure S2.** Ranked POP and TIC levels and concentrations for the 78 individual fish. Total POP levels are displayed in ng/g wet weight (*A*) and in micromolar (*B*). The TIC levels for each individual fish are shown in micromolar (*C*) and in ng/g wet weight (*D*).


**Figure S3.** Concentrations OCPs and PCBs and relative contributions of congeners at four additional sites. Shown are the range of total OCP (*A*) and PCB (*B*) concentrations in ng/g wet weight and the relative contributions of grouped POP congeners to total OCPs (*C*) and total PCBs (*D*) in percent. NA = not available.



**Figure S1. Capture locations and measured pollutant classes in this study.** Shown are the 12 different locations where a total of 117 wild yellowfin tuna were caught. Numbers in parenthesis show the sample size per site. Filled circles indicate the pollutant classes that were measured at each site, color-coded for the 29 OCPs (red), 9 PBDEs (orange), and 209 PCBs (green). The world map file is available in the public domain under the Creative Commons CC0 1.0 Universal Public Domain Dedication license (<https://commons.wikimedia.org/wiki/File%3ABlankMap-World6-Equirectangular.svg>).



**Figure S2. Ranked POP and TIC levels and concentrations for the 78 individual fish.** Total POP levels are displayed in ng/g wet weight (A) and in micromolar (B). The TIC levels for each individual fish are shown in micromolar (C) and in ng/g wet weight (D).



**Figure S3. Concentrations OCPs and PCBs and relative contributions of congeners at four additional sites.** Shown are the range of total OCP (A) and PCB (B) concentrations in ng/g wet weight and the relative contributions of grouped POP congeners to total OCPs (C) and total PCBs (D) in percent. NA = not available.