

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Effects of Radon and UV Exposure on Skin Cancer Mortality in Switzerland

Danielle Vienneau, Kees de Hoogh, Dimitri Hauri, Ana M. Vicedo-Cabrera, Christian Schindler, Anke Huss, and Martin Rössli, for the SNC Study Group

Table of Contents

Figure S1. Exposure distributions for radon in Bq/m³ (top) and UV-erythemal dose in W/m² (bottom)

Table S1. Negative control outcome: Population characteristics and number of deaths due to glioma (ICD10: C71, definitive primary cause of death)

Table S2. Hazard ratios per IQR (and 95%-confidence intervals) for radon (Bq/m³) and UV-erythemal dose (W/m²) exposure and type of skin cancer mortality, age 60

Table S3. Negative control outcome: Hazard ratios (and 95%-confidence intervals) for radon (per 100 Bq/m³) and UV-erythemal dose (per W/m²) exposure and glioma (ICD10: C71, definitive primary cause of death), age 60

Table S4. Sensitivity analyses (S4) stratified according to outdoor occupation with UV exposure, radon and UV-erythemal dose and mortality from all skin cancers, age 60