

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehponline@niehs.nih.gov](mailto:ehponline@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Associations of Prenatal Urinary Bisphenol A Concentrations with Child Behaviors and Cognitive Abilities**

Joseph M. Braun, Gina Muckle, Tye Arbuckle, Maryse Bouchard, William D. Fraser, Emmanuel Ouellet, Jean R. Séguin, Youssef Oulhote, Glenys M. Webster, and Bruce Lanphear

#### **Table of Contents**

**Table S1:** Description of neurobehavioral tests administered to children or parents in the MIREC Study at approximately 3 years of age.

**Table S2:** Baseline maternal covariates among women whose children did not complete follow-up at 3 years of age (n=895), completed neurobehavioral assessments via questionnaire at 3 years of age (n=812), and completed in-person assessments of neurobehavior at 3 years of age (n=544).

**Table S3:** Covariate adjusted mean child BASC-2 and BRIEF-P scores by specific gravity standardized maternal urinary BPA quintile: MIREC Study (n=806-812).

**Table S4:** Covariate adjusted mean child SRS-2 scores by specific gravity standardized maternal urinary BPA quintile: MIREC Study (n=537).

**Figure S1:** Directed acyclic graph describing the relations between prenatal urinary BPA concentrations, child neurobehavior at 3 years of age, and covariates.

**Figure S2:** Adjusted difference in child SRS-2 total scores with 2-fold increase in maternal urinary BPA concentrations during pregnancy: Impact of various covariate adjustments (MIREC Study).

**Figure S3:** Difference in child BASC-2 internalizing scores with 2-fold increase in maternal urinary BPA concentrations during pregnancy: Impact of various covariate adjustments (MIREC Study).