

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

First Trimester Urinary Bisphenol A Concentration in Relation to Anogenital Distance, an Androgen-Sensitive Measure of Reproductive Development, in Infant Girls

Emily S. Barrett, Sheela Sathyanarayana, Omar Mbowe, Sally W. Thurston, J. Bruce Redmon, Ruby H.N. Nguyen, and Shanna H. Swan

Table of Contents

Table S1. TIDES participant demographics by study center.

Table S2. Linear regression models examining the relationship between log(SpG adj. (BPA)) and covariates and log-transformed anogenital distance measures in newborn daughters (n=381).

Figure S1. Scatterplots examining the relationship between AGD-AC and continuous covariates.

Figure S2. Scatterplots examining the relationship between AGD-AF and continuous covariates.

Figure S3. Adjusted generalized additive model plots with the response variable AGD-AC

Figure S4. Adjusted generalized additive model plots with the response variable AGD-AF