

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material to:

Endocrine disruption in human fetal testis explants by individual and combined exposures to selected pharmaceuticals, pesticides and environmental pollutants

Pierre Gaudriault , Séverine Mazaud-Guittot , Vincent Lavoué , Isabelle Coiffec , Laurianne Lesné , Nathalie Dejudcq-Rainsford , Martin Scholze , Andreas Kortenkamp , Bernard Jégou

Table of Contents

TABLE S1. Names of the 27 screened chemicals. The bold names correspond to the chemicals included in our mixtures.....2

Figure S1. Concentration-response data for individual chemicals from the organotypic culture system in human fetal testes. The graphs show the experimental data as mean \pm SEM of at least 3 independent studies Testosterone production is represented as relative to the first day of culture (D0) production and the control level, see text for more details. These datasets represent the 16 screened chemicals not included in our mixtures.....3

Figure S2. Single compound effect on the apoptosis of the human fetal testis treated at highest non-toxic concentration after 96h of culture. Apoptosis positive cells were labeled with an anti- cleaved caspase 3 antibody. The 3,3'-diaminobenzidine tetrahydrochloride staining appears brown in all photos, and sections were counterstained with hematoxylin. Testis cords and interstitial tissue could be identified in all the sections (dashed line represent testis cords). Scale bar corresponds to 20 μ m.....6