

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

A human mixture risk assessment for neurodevelopmental toxicity associated with polybrominated diphenyl ethers used as flame retardants

Olwenn V Martin, Richard M Evans, Michael Faust and Andreas Kortenkamp

Table of Contents

Table S1. PBDE congeners in breast milk samples (ng/g lipid weight)

Table S2. PBDE intakes via food for children (1-3 years), ng/kg bw/d

Table S3. PBDE intakes via food for adults, ng/kg bw/d

Table S4. PBDE congeners in indoor dust samples (ng/g)

Table S5. PBDE congeners in serum and blood samples (ng/g lipid weight)

Table S6. Mixture risk assessment for breast-feeding infants (0-3 months), supplemental information to Table 2, main text

Table S7. Mixture risk assessment for small children (1-3 years), supplemental information to Table 2, main text

Table S8. Mixture risk assessment for adults, supplemental information to Table 2, main text

Table S9. Mixture risk assessment for foetuses (supplemental information to Table 3 of main text)

Table S10. Level 2 mixture risk assessment for small children based on body burdens estimated from PBDE serum levels (supplemental information to Table 3 of main text)

Table S11. Level 3 mixture risk assessment for small children based on body burdens estimated from PBDE serum levels (supplemental information to Table 3 of main text)

Table S12. Level 2 mixture risk assessment for adults based on body burdens estimated from PBDE serum levels (supplemental information to Table 3 of main text) – France

Table S13. Level 3 mixture risk assessment for adults based on body burdens estimated from PBDE serum levels (supplemental information to Table 3 of main text) – France

Table S14. Level 2 mixture risk assessment for adults based on body burdens estimated from PBDE serum levels (supplemental information to Table 3 of main text) – USA

Table S15. Level 3 mixture risk assessment for adults based on body burdens estimated from PBDE serum levels (supplemental information to Table 3 of main text) – USA

References