

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Children's Lead Exposure: A Multimedia Modeling Analysis to Guide Public Health Decision-Making

Valerie Zartarian, Jianping Xue, Rogelio Tornero-Velez, and James Brown

Table of Contents

S1. SHEDS-IEUBK Model Coupling Methodology

S2. Summary of Key Model Input Values

Figure S1. Age-specific relationships between available intake ($\mu\text{g}/\text{day}$) and blood lead ($\mu\text{g}/\text{dL}$)

Figure S2. SHEDS-IEUBK Predicted BLL Evaluation with 2009-2014 NHANES Data, for 2-Day Analysis (children 1 to <6 years)

Figure S3. SHEDS-IEUBK Evaluation for different NHANES sampling times (2-day averaging time analyses), 1 to <6 year-olds

Figure S4. SHEDS-IEUBK Evaluation for different NHANES sampling periods 2009-2014 (30-day averaging time analyses), 1 to <6 year-olds

Figure S5. Estimated Relative Contribution to BLL by Exposure Pathway adjusted for bioavailability of each pathway for U.S. children (0 to 6 months, 1 to <2 year-olds, and 2 to <6 years) using available data (30-day averaging time analyses; BLL on y-axis)

Figure S6. Percent Contribution by Pathway adjusted for bioavailability of each media for NHEXAS Region 5 Study

Figure S7. Impact of using von Lindern et al. (2016) Soil/Dust Ingestion Rate: SHEDS-IEUBK BLL estimates vs. NHANES years 2009-14; 1 to <2 year-olds

Table S1. Age-specific regressions for relationship between GI uptake ($\mu\text{g}/\text{day}$) and blood lead concentration ($\mu\text{g}/\text{dL}$)

Table S2. Comparison of SHEDS-IEUBK and IEUBK batch for a population of 8872 individuals with nominal blood values of 1-5 Pb $\mu\text{g}/\text{dL}$.

Table S3. Summary of Main SHEDS-IEUBK Inputs for Multimedia Pb Analysis

Table S4. Pb Statistics from NHEXAS Region 5 Study (based on Clayton et al., 1999)

Table S5. Correlation Coefficients Inputs for Lead Exposure Assessment based on Clayton et al., 1999 and HUD data

Table S6. SHEDS-IEUBK Evaluation (National Scale, 2-day averaging time analysis)

Table S7. Prediction of Blood Lead Concentration ($\mu\text{g}/\text{dL}$) for NHEXAS Region 5 Study

Table S8. Sensitivity analyses on Dietary Pb Intake and Soil/Dust Ingestion factors (ages 1 to <6 years; NHANES 2009-2014)

Table S9. Impact of Using Alternate Soil/Dust Ingestion Rate from von Lindern et al. 2016

Table S10. Impact of Using Alternate Soil/Dust Ingestion Rate Assuming 80mg/day