

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Plasma Concentrations of Per- and Polyfluoroalkyl Substances at Baseline and Associations with Glycemic Indicators and Diabetes Incidence among High-Risk Adults in the Diabetes Prevention Program Trial

Andres Cardenas, Diane R. Gold, Russ Hauser, Ken P. Kleinman, Marie-France Hivert, Antonia M. Calafat, Xiaoyun Ye, Thomas F. Webster, Edward S. Horton, and Emily Oken

Table of Contents

Figure S1. PFAS geometric means in the Diabetes Prevention Program (red) and six NHANES cycles (blue colors)

Table S1. PFAS geometric means and 95% CIs (ng/mL) in NHANES and in the Diabetes Prevention Program (DPP)

Table S2. Adjusted cross-sectional associations between baseline PFAS plasma concentrations and glycemic outcomes in the Diabetes Prevention Program among isomers of PFOS (n-PFOS, Sm-PFOS, Sm2-PFOS) and PFOA (n-PFOA and Sb-PFOA) and PFOA (n-PFOA, Sb-PFOA)

Table S3. Adjusted longitudinal associations of baseline plasma PFAS concentrations and prospective metabolic and glycemic measurements in the DPP.

Table S4. Adjusted Hazards Ratios (HRs) for time to first fasting hyperglycemia during study period relative to \log_2 -PFAS plasma concentrations in the Diabetes Prevention Program trial

Table S5. Adjusted Hazards Ratios (HRs) for the risk of developing diabetes during study period relative to \log_2 -PFAS plasma concentrations stratified by treatment assignment

Table S6. Adjusted Hazards Ratios (HRs) for the risk of developing diabetes during study period relative to \log_2 -PFAS plasma concentrations stratified by sex

Table S7. Adjusted Hazards Ratios (HRs) for the risk of developing diabetes during study period: multiplicative interaction between baseline \log_2 -PFAS plasma concentrations and BMI at baseline.

Appendix A. Longitudinal Regression Model Equations