

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Environmental Suitability of *Vibrio* Infections in a Warming Climate: An Early Warning System

Jan C. Semenza, Joaquin Trinanes, Wolfgang, Lohr, Bertrand Sudre, Margareta Löfdahl, Jaime Martinez-Urtaza, Gordon L. Nichols, and Joacim Rocklöv

Table of Contents

Figure S1. Isotherms for the differences between 2016 and projections for 2050 under RCP4.5 and RCP8.5, for June, July, August, September.

Figure S2. Number of months per year with risk of transmission (%) of *Vibrio* infections associated with climate change scenario RCP 4.5, 21th century.

Figure S3. Number of months per year with risk of transmission (%) of *Vibrio* infections associated with climate change scenario RCP 8.5, 21th century.

Figure S4. *Vibrio* suitability in New York in May–July 2012 during a *Vibrio parahaemolyticus* outbreak associated with shellfish from Oyster Bay Harbor in New York.

Figure S5. Suitability for *Vibrio* during an *V. parahaemolyticus* outbreak in British Columbia in 2015.

Figure S6. Trends for British Columbia based only on sea surface temperature for RCP 4.5 and RCP 8.5, from 2010 till 2060, by month.