

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Cadmium Body Burden and Gestational Diabetes Mellitus: A Prospective Study

Wenyu Liu, Bin Zhang, Zheng Huang, Xinyun Pan, Xiaomei Chen, Chen Hu, Hongxiu Liu, Yangqian Jiang, Xiaojie Sun, Yang Peng, Wei Xia, Shunqing Xu, and Yuanyuan Li

Table of Contents

Table S1. Associations between maternal urinary Cd levels and GDM, stratified by parity.

Table S2. Risk ratios from GEE model evaluating the interaction between Cd and As.

Table S3. Association between maternal urinary cadmium levels and gestational diabetes among women without passive smoking during pregnancy (n =1450).

Figure S1. Adjusted β -Coefficients and 95% CIs for urinary Cd levels (log₁₀ transformed SG-corrected Cd levels ($\mu\text{g/L SG}$) and plasma glucose (mmol/L) at different time points during oral glucose tolerance test (A: fasting plasma glucose; B: 1-hour plasma glucose; C: 2-hour plasma glucose; D: sum of plasma glucose z-score). Squares represent total women, diamonds and triangles represent women with male and female fetuses, respectively.

Figure S2. Adjusted β -Coefficients and 95% CIs for urinary Cd levels (log₁₀ transformed SG-corrected Cd levels ($\mu\text{g/L SG}$) and plasma glucose (mmol/L) at different time points during oral glucose tolerance test (A: fasting plasma glucose; B: 1-hour plasma glucose; C: 2-hour plasma glucose; D: sum of plasma glucose z-score). Squares represent normal weight women and diamonds represent overweight/obese women.