

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Prenatal Exposure to Perfluoroalkyl Substances and IQ Scores at Age 5; a Study in the Danish National Birth Cohort

Zeyan Liew, Beate Ritz, Cathrine Carlsen Bach, Robert F. Asarnow, Bodil Hammer Bech, Ellen A. Nohr, Rossana Bossi, Tine Brink Henriksen, Eva Cecilie Bonefeld-Jørgensen, and Jørn Olsen

Table of Contents

Figure S1. Selection of the LDPS study population from the DNBC.

Figure S2. Adjusted difference in IQ scores in boys and girls at age five according to prenatal PFAS levels (per natural-log ng/ml increase).

Table S1. Comparison of the characteristics for those invited and those who participated in the LDPS-DNBC cohort.

Table S2. PFAS concentrations in the LDPS cohort compared with data from a previous DNBC sample and the U.S. NHANES data reported as median concentration (25th-75th percentiles) in ng/ml.

Table S3. Distributions of selected PFASs by study characteristics in the LDPS-DNBC cohort (N=1,592).

Table S4. Adjusted difference of child IQ scores at age five according to prenatal PFAS (per natural-log ng/ml increase), stratified by parity.

Table S5. Adjusted difference of child IQ scores at age five according to prenatal PFAS (per natural-log ng/ml increase), stratified by maternal socio-economic status (SES).

Table S6. Mean differences in IQ scores in 1,592 children at age five according to prenatal PFAS level quartiles.

Table S7. Adjusted difference of child IQ scores according to prenatal PFAS (per natural-log ng/ml increase), comparing models accounting for selection factors.

Table S8. Adjusted difference of child IQ scores at age five according to prenatal PFAS (per natural-log ng/ml increase), additionally adjusting for birth year.

Table S9. Adjusted difference of child IQ scores at age five according to prenatal PFAS (per natural-log ng/ml increase), excluding blood samples that were subjected to processing delay.

Reference