



ENVIRONMENTAL HEALTH PERSPECTIVES

<http://www.ehponline.org>

Urinary Phthalate Metabolite Concentrations and Reproductive Outcomes among Women Undergoing *in Vitro* Fertilization: Results from the EARTH Study

Russ Hauser, Audrey J. Gaskins, Irene Souter, Kristen W. Smith, Laura E. Dodge, Shelley Ehrlich, John D. Meeker, Antonia M. Calafat, and Paige L. Williams, for the Earth Study Team

<http://dx.doi.org/10.1289/ehp.1509760>

Received: 27 January 2015

Accepted: 30 October 2015

Advance Publication: 6 November 2015

Note to readers with disabilities: *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.



Urinary Phthalate Metabolite Concentrations and Reproductive Outcomes among Women Undergoing *in Vitro* Fertilization: Results from the EARTH Study

Russ Hauser^{1,2,3}, Audrey J. Gaskins^{2,4}, Irene Souter³, Kristen W. Smith¹, Laura E. Dodge², Shelley Ehrlich⁵, John D. Meeker⁶, Antonia M. Calafat⁷, and Paige L. Williams^{2,8}, for the Earth Study Team

¹Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; ²Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; ³Department of Obstetrics and Gynecology, Harvard Medical School/Massachusetts General Hospital Fertility Center, Boston, Massachusetts, USA; ⁴Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; ⁵Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA; ⁶Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, Michigan, USA; ⁷National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA; ⁸Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA. *Please note that Russ Hauser and Audrey Gaskins are considered joint first authors.

Address correspondence to Russ Hauser, Environmental and Occupational Medicine and Epidemiology Program, Department of Environmental Health, Building I, Room 1409, 665 Huntington Avenue, Boston, MA 02115 USA. Telephone: 617-432-3326. E-mail:

رهاuser@hsph.harvard.edu

Running title: Phthalates and reproductive outcomes

Acknowledgments and grant information: Work supported by grants ES009718, ES022955, ES000002, and T32ES007069 from the National Institute of Environmental Health Sciences (NIEHS) and grant T32HD060454 from the National Institute of Child Health and Human Development (NICHD). The authors gratefully acknowledge Manori Silva, Ella Samandar, Jim Preau, and Tao Jia (CDC, Atlanta, GA) for measuring the urinary concentrations of the phthalate metabolites. We also acknowledge all members of the EARTH study team, specifically the Harvard T. H. Chan School of Public Health research nurses Jennifer B. Ford and Myra G. Keller, research staff Ramace Dadd and Patricia Morey, physicians and staff at Massachusetts General Hospital fertility center and a special thanks to all the study participants.

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Competing financial interests: The authors declare they have no actual or potential competing financial interests.

Abstract

Background: Evidence from both animal and human studies suggests that exposure to phthalates may be associated with adverse female reproductive outcomes.

Objective: We evaluated the associations between urinary concentrations of phthalate metabolites and outcomes of assisted reproductive technologies (ART).

Methods: This analysis included 256 women enrolled in the Environment and Reproductive Health (EARTH) prospective cohort study (2004-2012) who provided 1-2 urine samples per cycle prior to oocyte retrieval. We measured 11 urinary phthalate metabolites [mono-2-ethylhexyl phthalate (MEHP), mono-2-ethyl-5-hydroxyhexyl phthalate (MEHHP), mono-2-ethyl-5-oxohexyl phthalate (MEOHP), mono-2-ethyl-5-carboxypentyl phthalate (MECPP), mono-isobutyl phthalate (MiBP), mono-n-butyl phthalate (MBP), monobenzyl phthalate (MBzP), monoethyl phthalate (MEP), monocarboxyisooctyl phthalate (MCOP), and monocarboxyisononyl phthalate (MCNP), and mono(3-carboxypropyl) phthalate (MCP)]]. We used generalized linear mixed models to evaluate the association of urinary phthalate metabolites with IVF outcomes, accounting for multiple IVF cycles per woman.

Results: In multivariate models, women in the highest as compared to lowest quartile of MEHP, MEHHP, MEOHP, MECPP, \sum DEHP (MEHP+MEHHP+MEOHP+MECPP) and MCNP had lower oocyte yield. Similarly, the number of mature (MII) oocytes retrieved was lower in the highest versus lowest quartile for these same phthalate metabolites. The adjusted difference (95% CI) in proportion of cycles resulting in clinical pregnancy and live birth between women in the fourth vs. first quartile of \sum DEHP were -0.19 (-0.29, -0.08) and -0.19 (-0.28, -0.08), respectively, and there was also a lower proportion of cycles resulting in clinical pregnancy and live birth for individual DEHP metabolites.

Conclusions: Urinary concentrations of DEHP metabolites were inversely associated with oocyte yield, clinical pregnancy and live birth following ART.

INTRODUCTION

Infertility, a disease characterized by the inability to have a child, affects one in six couples and will likely rise as the postponement of childbearing increases in developed regions of the world (Chandra et al. 2005; Evers 2002; Pinnelli and Di Cesare 2005). Even among fertile women, 22% of pregnancies fail before they can be clinically recognized (Wilcox et al. 1988). The associated health care cost of infertility is in the billions of dollars per year and does not include the tremendous physical and psychological burden placed on the couple. This highlights the public health importance of understanding risk factors that may impair the ability to have a child.

It is well appreciated that environmental exposures are potential risk factors for infertility and adverse pregnancy outcomes. Human and experimental animal studies identify several classes of chemicals that adversely impact fertility and pregnancy. Recently, the American College of Obstetricians (ACOG) and the American Society for Reproductive Medicine (ASRM) issued a Joint Committee Opinion on ‘Exposure to Toxic Environmental Agents’ which emphasized that exposure to environmental chemicals was ubiquitous and preconception and prenatal exposure can have a profound effect on reproductive health (ACOG 2013). The Joint Committee Opinion concluded that ACOG and ASRM, along with scientists and other clinical practitioners, noted the need for timely action to identify and reduce exposure to environmental agents while addressing the consequences of exposure.

One class of environmental chemicals for which there is concern related to its risk of adverse reproductive and developmental effects are the ortho-phthalates (herein referred to as phthalates), a family of multifunctional chemicals widely used in personal care and consumer products.

Human exposure to phthalates is widespread and occurs through multiple routes, including ingestion, inhalation, dermal contact, and parenteral exposure from medical devices containing phthalates (CDC 2013; Hauser and Calafat 2005). The low molecular weight phthalates, such as diethyl phthalate (DEP), di-n-butyl phthalate (DnBP), and di-iso-butyl phthalate (DiBP) are principally used in personal care products (e.g., body lotions, cosmetics, shampoos, deodorants), as solvents and plasticizers for cellulose acetate, varnishes and coatings, including coatings used for time release in some orally administered medications (ATSDR 2002; Hauser and Calafat 2005). High-molecular-weight phthalates (e.g., di(2-ethylhexyl) phthalate (DEHP) and di-isononyl phthalate (DiNP)) are used primarily as plasticizers in the manufacture of flexible vinyl, which, in turn, is used in consumer products, flooring and wall coverings, food contact applications, and medical devices (ATSDR 2002; Hauser and Calafat 2005). As these phthalate plasticizers are not chemically bound to the polymer chain, they leach, migrate, or off gas, thus leading to human exposure (Koch et al. 2006). Phthalates have a short half-life (i.e., hours) and are rapidly metabolized and excreted in urine and feces (Hauser and Calafat 2005; Silva et al. 2006).

Some phthalates are anti-androgenic and adversely affect the development of the male fetus and male fertility following gestational exposure in the rat to high doses of phthalates (Fisher 2004). However, the impact of phthalates on female reproductive health, fertility, and early pregnancy outcomes is much less studied (Ema et al. 2000; Gray et al. 2006; Lovekamp and Davis 2001; Svechnikova et al. 2007). In Gray et al., oral administration of DnBP to female Long Evans hooded rats, from weaning through puberty, mating and gestation, induced mid-pregnancy abortions with up to a 90% reduction in the percentage of females delivering live pups at 1000

mg/kg/day (Gray et al. 2006). The decreased litter size was associated with reduced serum progesterone levels and ovarian progesterone production on gestational day 13. Other studies have shown a reduction in serum levels of progesterone and estradiol in prepubertal female rats dosed with DEHP for 10 days (Svechnikova et al. 2007) and increased post-implantation embryo loss was reported in rats treated with DnBP (Ema et al. 2000).

A limited number of epidemiologic studies have reported associations between measures of exposure to some phthalates with decreased rates of pregnancy and risk of miscarriage, as well as complications such as anemia, toxemia, and preeclampsia in women (Heudorf et al. 2007). A recent human study reported an association between higher urinary concentrations of mono-2-ethylhexyl phthalate (MEHP), a metabolite of DEHP, and early pregnancy loss (Toft et al. 2012). However, as far as we are aware, no human studies have explored the association of phthalate exposure with very early pregnancy outcomes, such as oocyte fertilization and embryo implantation. The present analysis uses data from our ongoing prospective cohort study, the Environment and Reproductive Health study, using assisted reproductive technologies (ART) as a model of human reproduction to explore the relationship between environmental exposures and both male and female reproductive health outcomes.

METHODS

Participants

Study participants were recruited into the EARTH Study between November 2004 and April 2012 from patients undergoing *in vitro* fertilization (IVF) at the Massachusetts General Hospital (MGH) Fertility Center. All women over 18 years of age and less than 46 years (at enrollment)

were eligible to participate (approximately 60% of those contacted by the research nurses participated in the study). Women must have contributed their own oocytes and at least one urine sample for the measurement of phthalate metabolites during an IVF cycle to be included in the present analysis. From 429 eligible IVF cycles, we excluded IVF cycles for which women used an egg donor (N=18), cryo-thaw cycles (N=35), and those missing information on BMI (N=1). This left us with 375 IVF cycles with complete information on exposure, outcome, and all covariates. The EARTH study was approved by the Human Studies Institutional Review Boards of the MGH, Harvard School of Public Health (HSPH), and the Centers for Disease Control and Prevention (CDC). Participants signed an informed consent after the study procedures were explained by a research nurse and all questions were answered.

Clinical data

Clinical information at entry into the study and after each IVF cycle was abstracted from the patient's electronic medical record by the research nurse. At study entry and each IVF cycle, a blood sample was drawn on the third day of the menstrual cycle and the serum was analyzed for follicle stimulating hormone (FSH) with an automated electrochemiluminescence immunoassay at the MGH Core Laboratory as previously described (Mok-Lin et al. 2010). At entry into the EARTH Study and subsequent to an infertility evaluation, each patient was given an infertility diagnosis by a physician at the MGH Fertility Center according to the Society for Assisted Reproductive Technology (SART) definitions as previously described (Mok-Lin et al. 2010; SART 2013). At entry, the participant's date of birth was collected and weight and height were measured by the nurse. Body mass index (BMI) was calculated as weight (in kilograms) per height (in meters) squared.

At each IVF cycle, depending on clinical indications and factors such as age and infertility diagnosis, women underwent one of three IVF treatment protocols : (1) luteal phase gonadotropin releasing hormone (GnRH) agonist (low-, regular-, or high-dose leuprolide acetate, Lupron), (2) follicular phase GnRH-agonist/Flare stimulation, and (3) GnRH-antagonist. For each cycle, on the day of ovulation trigger with HCG, serum peak estradiol was measured with an automated electrochemiluminescence immunoassay at the MGH Core Laboratory (Mok-Lin et al. 2010). Couples underwent assisted reproduction with conventional IVF or intracytoplasmic sperm injection (ICSI) (ASRM 2015; Bhattacharya et al. 2001). The ICSI technique was originally developed to treat cases of severe male factor infertility (one or more severely decreased semen parameters) but is now used for certain other indications (ASRM 2012). After egg retrieval, embryologists classified oocytes as germinal vesicle, metaphase I, metaphase II (MII) or degenerated. Embryologists determined fertilization 17-20 hours after insemination. Fertilization was confirmed by the presence of a fertilized oocyte with two pronuclei. Embryos were monitored for cell number and morphological quality (1 (best) to 5 (worst)) on day 2 and 3. For analysis we classified embryos as best quality if they had 4 cells on day 2, 8 cells on day 3, and a morphologic quality score of 1 or 2 on days 2 and 3 (Veeck and Zaninovic 2003). An overall score of 1 or 2 was considered high quality, 3 was considered intermediate quality and 4 or 5 indicated poor quality embryos. If a cycle lacked information on day 2 or day 3 embryo quality (e.g. failed fertilization, day 2 transfer) they were classified as having no best quality embryos. We defined implantation as a serum β -hCG level > 6 mIU/mL typically measured 17 days (range 15–20 days) after egg retrieval, clinical pregnancy as the presence of an intrauterine pregnancy confirmed by ultrasound at approximately 6 weeks

gestation, and live birth as the birth of a neonate on or after 24 weeks gestation. Prior to the start of the EARTH study, we determined that the infertility medications do not contain phthalates (Kelley et al. 2012) and we tested the IVF equipment and medical supplies for phthalates (data not shown). We did not identify these as potential sources of exposure.

Urinary phthalate metabolite measurements

Two urine samples were collected during each IVF cycle (between day three and nine of the gonadotropin phase and on the day of the oocyte retrieval). The median time between the two urine samples collected per cycle was 8 days (interquartile range: 6, 9), Urine samples were collected between November 2004 and April 2012. Urine was collected in a sterile polypropylene cup. After measuring specific gravity (SG) using a handheld refractometer (National Instrument Company, Inc.), the urine was divided into aliquots and frozen at -80°C. Samples were shipped on dry ice overnight to the CDC (Atlanta, GA, USA) for the quantification of concentrations of MEHP, mono-2-ethyl-5-hydroxyhexyl phthalate (MEHHP), mono-2-ethyl-5-oxohexyl phthalate (MEOHP), mono-2-ethyl-5-carboxypentyl phthalate (MECPP), mono(3-carboxypropyl) phthalate (MCPP), monocarboxyisooctyl phthalate (MCOP), monocarboxyisononyl phthalate (MCNP), monobenzyl phthalate (MBzP), monoethyl phthalate (MEP), mono-isobutyl phthalate (MiBP) and mono-n-butyl phthalate (MBP). The analytical approach, based on solid phase extraction coupled with high performance liquid chromatography-isotope dilution tandem mass spectrometry, followed standard QA/QC procedures as previously described (Silva et al. 2007). The limits of detection (LOD) were 0.5-1.2 µg/L (MEHP), 0.2-0.7 µg/L (MEHHP and MEOHP), 0.2-0.6 µg/L (MECPP), 0.1-0.2 µg/L (MCPP), 0.2-0.7 µg/L (MCOP), 0.2-0.6 µg/L (MCNP), 0.2-0.3 µg/L (MBzP), 0.4-0.8 µg/L

(MEP), and 0.4-0.6 $\mu\text{g/L}$ (MBP). We calculated the molar sum of DEHP metabolites (ΣDEHP) by dividing each metabolite concentration by its molecular weight and then summing:
$$[(\text{MEHP} \cdot (1/278.34)) + (\text{MEHHP} \cdot (1/294.34)) + (\text{MEOHP} \cdot (1/292.33)) + (\text{MECPP} \cdot (1/308.33))].$$
MCPP is a metabolite of di-n-octyl phthalate (DnOP) and a nonspecific metabolite of high molecular weight phthalates, MCOP is a metabolite of DiNP, and MCNP is a metabolite of DiDP.

Statistical analysis

Demographic characteristics of the study participants and clinical characteristics (cycle- and embryo-level) were reported using mean \pm SD or percentages. Cycle specific urinary concentrations of phthalate metabolites were calculated using the geometric mean (GM) of the two urinary phthalate concentrations from each IVF cycle. Urinary phthalate metabolite concentrations below the LOD were replaced with a value equal to the $\text{LOD}/\sqrt{2}$ (Hornung and Reed 1990). Although there are newer methods to replace values below the LOD (Cole et al. 2009; Nie et al. 2010), our use of quartiles minimized the impact of samples less than LOD; essentially all women who were below the LOD were classified into the first quartile given that the lowest detection limit was 75.5% (for MEHP). For all other metabolites, the lowest quartile included women with concentrations below the LOD and others with low concentrations.

To adjust for urinary dilution, the following formula was used: $P_c = P[(1.015 - 1)/SG - 1]$, where P_c is the SG-corrected phthalate metabolite concentration ($\mu\text{g/L}$), P is the measured phthalate metabolite concentration ($\mu\text{g/L}$), and 1.015 is the mean (and median) SG level in the study

population (Smith et al. 2012). We used SG-corrected phthalate metabolite concentrations in all analyses.

We fit multivariate generalized linear mixed models with random intercepts to evaluate the association between urinary phthalate metabolites and IVF outcomes. These models allow for the use of multiple outcome observations per individual while accounting for within-person correlations in outcomes. These models are also appropriate and can provide unbiased estimates in the presence of an unbalanced design (e.g., different number of cycles contributed per woman) when imbalance in the number of IVF cycles is not completely random, and the lack of balance can be accurately predicted by all measured covariates in the adjusted model. A linear distribution and identity link function was specified for peak estradiol, a Poisson distribution and log link function were specified for oocyte counts, and a binomial distribution and logit link function were specified for fertilization, embryo quality, and clinical outcomes. Tests for trend were conducted across quartiles using the median urinary phthalate metabolite concentration in each quartile as a continuous variable in the regression models. All results are presented as population marginal means, adjusted for covariates.

Confounding was evaluated using prior knowledge and descriptive statistics from our cohort through the use of directed acyclic graphs. The following covariates were considered for inclusion in the final model: maternal age (continuous), BMI (continuous), smoking status (ever smoked and never smoked), year of treatment (2004-2006, 2007-2009, 2010-2012), day 3 FSH (continuous), treatment protocol type (Luteal phase or Flare/GnRH antagonist), and primary infertility diagnosis (female factor, male factor, and unexplained). Variables were included in the

final model if they were associated with phthalate exposure in our population, were suspected to be associated with phthalate exposure based on previous research, or were strong predictors of the outcome. To test whether the associations of urinary phthalate metabolite concentrations with IVF outcomes were modified by ICSI, a product of quartiles of phthalate and a binary variable representing the presence or absence of ICSI was entered into the models. Given our limited power, a suggestion of interaction was considered if the p-value for this interaction term was <0.10 . As a sensitivity analysis we evaluated the association between urinary phthalate metabolite concentrations and probability of implantation, clinical pregnancy, and live birth per embryo transfer to assess whether any association between urinary phthalates and clinical outcomes remained after excluding early failures. We conducted all statistical analyses using SAS version 9.2 (SAS Institute Inc., Cary, NC) and considered two-sided significance levels less than 0.05 as statistically significant.

RESULTS

Our analysis included 256 women who were on average 35.3 years of age; 72% had never smoked and 82% were Caucasian (Table 1). The primary SART diagnosis was 37% male factor, followed by 33% unexplained infertility and 30% female factor infertility. The women underwent a total of 375 IVF cycles, with 179 women (69%) contributing 1 cycle, 47 women (18%) contributing 2 cycles, 21 women (8%) contributing 3 cycles, 7 women (3%) contributing 4 cycles, 1 woman ($<1\%$) contributing 5 cycles, and 1 woman ($<1\%$) contributing 6 cycles. The luteal phase treatment protocol was used for 67% of IVF cycles and ICSI (fertilization method) was used for 55% of IVF cycles (Table 2). Among cycles with male factor as the primary infertility diagnosis, 88% used ICSI as compared to 40% and 31% using ICSI with female factor

or unexplained infertility diagnoses, respectively. A total of 673 urine samples were collected from the 256 women, 79% of the IVF cycles had 2 urine samples and 21% IVF cycles had only 1 urine sample per cycle. The distribution of each urinary phthalate metabolite in our population is shown in Table 3. An overview of the 375 IVF cycles is shown in Figure 1. In brief, 337 cycles out of the initial 375 cycles underwent embryo transfer (90%). Out of the cycles that underwent embryo transfer, the percent resulting in implantation, clinical pregnancy, and live birth were 60%, 53%, and 44%, respectively.

In multivariable models adjusted for age, BMI, smoking status and infertility diagnosis, only MCNP was associated with peak estradiol (p -trend=0.009). The adjusted mean peak estradiol was 2225 $\mu\text{g/L}$ (95% CI 2036, 2414) in quartile 1 compared to 1902 $\mu\text{g/L}$ (95% CI 1710, 2094) in quartile 4 (p -value for Q4 vs. Q1 = 0.02) (results not shown). In adjusted multivariable models, there were statistically significant associations of ΣDEHP , MEHP, MEHHP, MEOHP, MECPP and MCNP with reduced total oocyte yield for the 4th compared to the 1st quartile (Figure 2). Consistent with the decrease in oocyte yield, there were decreased numbers of MII oocytes retrieved in quartile 4 as compared to 1 of ΣDEHP , MEHP, MEHHP, MEOHP, MECPP, and MCNP (Figure 2). The total number of fertilized oocytes was significantly lower in quartile 4 as compared to quartile 1 for MCOP and MCNP, but not the other phthalate metabolites (Figure 2). There was no interaction between use of ICSI and the association of phthalates with the number of fertilized oocytes (data not shown, p -value for interaction > 0.10). In multivariable models there were no statistically significant associations of any phthalate metabolite with percent of cycles with 1 or more high quality embryos (p , trend > 0.05 for all metabolites and no point estimate comparisons were significant) (data not shown).

After controlling for age, BMI, smoking status, primary infertility diagnosis, increased quartiles of Σ DEHP were associated with a reduced probability of clinical pregnancy and live birth such that there appeared to be monotonic downward trend (p , trend=0.04 and 0.01, respectively) (Table 4). The adjusted difference (95% CI) in proportion of cycles resulting in clinical pregnancy between women in the highest DEHP quartile (4th) versus lowest quartile (1st) was -0.19 (-0.29, -0.08), representing a decrease in the probability of clinical pregnancy from 0.57 to 0.38 (p , trend=0.04). Similarly, women in this highest Σ DEHP quartile vs the lowest had a mean decrease in proportion of cycles resulting in live birth of 0.19 (-0.28, -0.08), representing a decrease from 0.47 to 0.28, respectively (p , trend=0.01). There was a similar monotonic downward trend in proportion of cycles resulting in clinical pregnancy and live birth across increasing quartiles of individual DEHP metabolites, MEHP, MEHHP, MEOHP, MECPP (Table 4). The monotonic downward trends between DEHP metabolites and implantation, clinical pregnancy and live birth were similar when analyses were restricted to cycles with embryo transfer ($n=337$ cycles) (data not shown). Although concentrations of the other phthalate metabolites measured (MEP, MBP, MIBP, MBzP, MCPP, MCOP and MCNP) were not significantly related to clinical outcomes following IVF (Table 4), seventeen of the eighteen adjusted models for probability of implantation, clinical pregnancy and live birth showed reduced proportions for quartile 4 as compared to quartile 1. Finally, across all the urinary phthalate metabolites, there were no interactions between the use of ICSI and associations of phthalates with live birth (data not shown).

It is well appreciated that during the course of the study, urinary concentrations of some of the phthalate metabolites have decreased in the general population (Zota et al. 2014) and there may also be a temporal trend in IVF success. In our dataset, urinary DEHP metabolite concentrations tended to be higher in earlier years but calendar year was not significantly associated with IVF outcome. When we explored adjusting for calendar year in the multivariable models, the effect estimates remained similar and the conclusions remained the same (data not shown). In order to keep the most parsimonious model possible, we opted to exclude year in the final multivariable model.

DISCUSSION

The current study utilized the model of IVF to investigate human reproduction and pregnancy outcomes, ranging chronologically from the number and maturity of oocytes at retrieval, number of fertilized embryos, and embryo implantation to live birth. Urinary metabolites of DEHP, and DiDP were associated with decreased oocyte yield and number of MII oocytes at retrieval, whereas only metabolites of DiNP and DiDP were associated with reduced fertilization rate. There is evidence that a decreased oocyte yield predicts poorer IVF outcomes (Ji et al. 2013; Sunkara et al. 2011). Urinary concentrations of DEHP metabolites were associated with reduced probability of clinical pregnancy and live birth. The magnitude of this reduction was clinically relevant as the adjusted proportion of live births in the highest quartile of \sum DEHP was 0.28 as compared to 0.47 in the lowest quartile. Although urinary concentrations of other phthalate metabolites (i.e., MEP, MBP, MIBP, MBzP, MCPP, and MCNP) were not significantly associated with decreased probability of implantation, clinical pregnancy or live birth, nearly all

of the models showed small to moderate reduced likelihood of these outcomes when the highest quartile was compared with the lowest quartile.

There are few epidemiologic studies on associations of phthalates with rates of pregnancy and miscarriage (Heudorf et al. 2007). A recent publication on time to pregnancy in a U.S. prospective cohort (LIFE study) did not find associations between maternal urinary concentrations of 14 phthalate metabolites and fecundity (Buck Louis et al. 2014). They enrolled 501 couples from 2005-2009 who discontinued contraception and attempted to become pregnant. They used time to pregnancy to assess fecundity and to determine the number of menstrual cycles required for pregnancy confirmed by human chorionic gonadotropin (hCG). Possible explanations for differences in results between our study which found associations between some urinary phthalate metabolites and reduced probability of pregnancy and live birth and the LIFE study which did not find associations of phthalate metabolites with reduced fecundity, include the choice of study population, differences in co-exposures and the potential for unmeasured confounding. Our study included couples from a fertility clinic whereas the LIFE study enrolled couples discontinuing contraception and attempting to become pregnant, excluding any couples with physician diagnosed infertility/sterility. The differences in results across studies may suggest that couples from an infertility clinic may represent a sensitive sub-population to environmental chemicals, specifically phthalates. Other explanations for differences in results may be related to analytical methods used to measure phthalate metabolites which varied across the two studies and potential co-exposures to other chemicals in the two study populations.

Another recent publication including 430 couples enrolled in a Danish prospective cohort study (from 1992 to 1994) reported an association between urinary concentrations of phthalate metabolites and pregnancy loss (Toft et al. 2012). They explored associations of urinary phthalate metabolite concentrations with early pregnancy loss endpoints comparable to those measured in our study. In the Danish study, 128 women were enrolled after discontinuation of birth control and followed prospectively until a clinical pregnancy or for six menstrual cycles if there was no clinical pregnancy. Subclinical embryonic loss (referred to as early pregnancy loss) was determined through measurements of hCG in urine samples collected on the first 10 days of each menstrual cycle. By phone interview with the women, data were also collected on self-reported clinical spontaneous abortion. Pregnancy loss was increased among women in the upper tertile of urinary MEHP concentrations (adjusted odds ratio = 2.87, 95% CI: 1.09, 7.57) as compared to the lowest tertile. When early and late pregnancy loss were analyzed in separate models, the odds ratio for urinary MEHP concentrations and early pregnancy loss was 40.67 (95% CI: 4.48, 369.5) for tertile 3 compared to tertile 1, whereas late pregnancy loss was negatively associated with urinary MEHP (OR = 0.25, 95% CI: 0.05, 1.8 for the third tertile compared to the first tertile). For the other phthalate metabolites measured (MEP, MBP, MBzP, and oxidative metabolites of DEHP, MEOHP and MEHHP) there were also elevated, non-significant, adjusted odds of early pregnancy loss (ranging from 1.13 (95% CI: 0.36, 3.59) for MEP to 1.64 (95% CI: 0.52, 5.20) for MBP and 3.11 (95% CI: 0.87, 11.09) for MBzP). There were no statistically significant associations between the other measured urinary phthalate metabolites and odds for spontaneous abortion.

Although the results of the Danish study and our study for DEHP metabolites are consistent, it should be cautioned that the early pregnancy loss endpoint in Danish women conceiving naturally was not the same as our endpoints of implantation, clinical pregnancy and live birth. It is important to point out that the authors of the Danish study urge caution when interpreting their results for MEHP because the incidence of early pregnancy loss in the first tertile was only 3%, thus they note that the very high OR for early pregnancy loss may be partially attributable to chance. Although both studies measured urinary phthalate metabolites in the concurrent conception cycle (measured on days 3 and 9 in our study, and day 10 in the Danish study), there were differences in analytical methods for the measurement of phthalate metabolites that may contribute to differences in results across studies.

Although implantation, clinical pregnancy, and live birth measured in women are not strictly comparable with pregnancy loss endpoints measured in experimental animals, studies in rats, albeit at doses much higher than those experienced by our study population (Heudorf et al. 2007), support associations between exposure to phthalates and pregnancy loss. Further support for the biological plausibility of the associations between early pregnancy loss in the epidemiologic studies were studies in rats that showed that exposure to DEHP led to reductions in aromatase mRNA and protein levels leading to a decreased conversion of testosterone to estradiol (Lovekamp-Swan and Davis 2003) and a decrease in progesterone and estradiol in prepubertal female rats dosed with DEHP (Svechnikova et al. 2007). Human studies have shown that low pregnancy levels of estradiol and progesterone are associated with fetal loss (Schindler 2004) and thus DEHP may be operating through this hormonal pathway to increase the risk of

early pregnancy loss. Our results raise concerns primarily about the potential of DEHP to result in lower probability of clinical pregnancy and live birth following ART.

One potential limitation of the present study is the generalizability of our findings to couples that conceive without medical assistance. However, couples with infertility represent 10-15% of US couples and the number of couples undergoing infertility treatment is increasing. The number of ART births in the U.S. more than tripled from 1996 to 2009 (ACOG 1998, CDC 2010). Based on this trend, the number of U.S. children born through ART over the next 10 years is expected to be well over 1 million (CDC 2010). We also have indirect evidence to support the generalizability of these results based on previous work among this cohort of couples undergoing IVF that have either been consistent with, or since been corroborated by, other studies of non-IVF populations. Some examples include associations between phthalates and lower male steroid hormones (Meeker et al. 2009; Mendiola et al. 2011; Pan et al. 2006); pesticide exposure and semen quality (Meeker et al. 2008; Perry et al. 2007), thyroid hormone levels (Lacasana et al. 2010; Meeker et al. 2006), and sperm DNA damage (Meeker et al. 2004; Xia et al. 2005); and self-reported maternal exposure to secondhand tobacco smoke and early pregnancy loss (Meeker et al. 2007a; Meeker et al. 2007b; Peppone et al. 2009). Of interest, our decision to study the fertility clinic population was based on utilizing an efficient design with sufficient power to investigate environmental influences on clinically relevant, yet previously unobservable, outcomes (e.g. fertilization rate and implantation) in a potentially vulnerable sub-population. Thus, in summary, we believe that any potential limitations in generalizability in the proposed study are outweighed by its strengths because other study designs do not allow for the depth and breadth of exploration into these reproductive health effects in such a cost-efficient

manner. In addition, other strengths include state-of-the-art exposure biomarker measures at the CDC, clinical outcomes from electronic medical records, and collection of data on potential confounders.

Another potential limitation of the study is misclassification of exposure based on urinary concentrations of phthalate metabolites. Single measures of urinary biomarkers of exposure to these short-lived compounds may not represent exposure during critical windows that may span weeks (i.e. implantation) or months (i.e. miscarriage, live birth). We have conducted a study (Braun et al. 2012), as have others (Hoppin et al. 2002; Peck et al. 2010; Preau et al. 2010), that demonstrate low to moderate reproducibility (intraclass correlations (ICC) ranged from 0.15 to 0.65) of urinary levels of phthalate metabolites over time frames of weeks to a few months.

Although it is well recognized that environmental exposures in close temporal proximity to the outcome are considered highly relevant, it is also important to consider that exposures distant from the outcome may also be relevant. For example, exposure very early in pregnancy may impact miscarriage later in pregnancy through effects on embryo quality and placentation. To reduce exposure misclassification in the present study, we collected two urine samples from each woman during each IVF cycle and used the geometric mean of the two cycle-specific urine concentrations as the exposure metric for the peri-conception exposure window.

In conclusion, using IVF as a model to investigate human reproduction and pregnancy outcomes, we found that concentrations of urinary metabolites of DEHP and DiDP were inversely associated with oocyte yield and number of MII oocytes at retrieval, whereas only metabolites of DiNP and DiDP were associated with reduced fertilization rate. Urinary concentrations of DEHP

metabolites were negatively associated with likelihood of clinical pregnancy and live birth following IVF. These results highlight the potential reproductive effects of low level exposure (i.e., background exposure levels of the general population) to phthalates and adverse IVF outcomes.

References

- American Congress of Obstetricians and Gynecologists (ACOG) Committee. 1998. Assisted reproductive technology in the United States and Canada: 1995 results generated from the American Society for Reproductive Medicine/Society for Assisted Reproductive Technology Registry. *Fertil Steril* 69(3): 389-398.
- American Congress of Obstetricians and Gynecologists (ACOG) Committee. 2010. National Center for Chronic Disease Prevention and Health Promotion's 2010 Assisted Reproductive Technology Success Rates Reports.
- American Congress of Obstetricians and Gynecologists (ACOG) Committee. 2012. Intracytoplasmic sperm injection (ICSI) for non-male factor infertility: a committee opinion. *Fertil Steril* 98(6): 1395-1399.
- American Congress of Obstetricians and Gynecologists (ACOG) Committee. 2013. Exposure to toxic environmental agents. *Fertil Steril* 100(4): 931-934.
- American Society for Reproductive Medicine (ASRM). 1999-2015. ART: Step-by-Step Guide. Available: <https://www.asrm.org/detail.aspx?id=1903> [accessed October 26 2015].
- Agency for Toxic Substances and Disease Registry (ATSDR). 2002. Available: <http://www.atsdr.cdc.gov/>.
- Bhattacharya S, Hamilton MP, Shaaban M, Khalaf Y, Seddler M, Ghobara T, et al. 2001. Conventional in-vitro fertilisation versus intracytoplasmic sperm injection for the treatment of non-male-factor infertility: a randomised controlled trial. *Lancet* 357(9274): 2075-2079.
- Braun JM, Smith KW, Williams PL, Calafat AM, Berry K, Ehrlich S, et al. 2012. Variability of urinary phthalate metabolite and bisphenol A concentrations before and during pregnancy. *Environ Health Perspect* 120(5): 739-745.
- Buck Louis GM, Sundaram R, Sweeney AM, Schisterman EF, Maisog J, Kannan K. 2014. Urinary bisphenol A, phthalates, and couple fecundity: the Longitudinal Investigation of Fertility and the Environment (LIFE) Study. *Fertil Steril*.
- Centers for Disease Control and Prevention (CDC). 2010. Assisted Reproductive Technology Reports. Atlanta, GA:US Centers for Disease Control and Prevention; 2010 [cited 2012 September 20]; Available from: <http://www.cdc.gov/ART/index.htm>.

- Centers for Disease Control and Prevention (CDC). 2013. Fourth National Report on Human Exposure to Environmental Chemicals, Updated Tables.
- Chandra A, Martinez GM, Mosher WD, Abma JC, Jones J. 2005. Fertility, family planning, and reproductive health of U.S. women: data from the 2002 National Survey of Family Growth. Vital and health statistics Series 23, Data from the National Survey of Family Growth(25): 1-160.
- Cole SR, Chu H, Nie L, Schisterman EF. 2009. Estimating the odds ratio when exposure has a limit of detection. *Int J Epidemiol* 38(6): 1674-1680.
- Ema M, Miyawaki E, Kawashima K. 2000. Effects of dibutyl phthalate on reproductive function in pregnant and pseudopregnant rats. *Reprod Toxicol* 14(1): 13-19.
- Evers JL. 2002. Female subfertility. *Lancet* 360(9327): 151-159.
- Fisher JS. 2004. Environmental anti-androgens and male reproductive health: focus on phthalates and testicular dysgenesis syndrome. *Reproduction* 127(3): 305-315.
- Gray LE, Jr., Laskey J, Ostby J. 2006. Chronic di-n-butyl phthalate exposure in rats reduces fertility and alters ovarian function during pregnancy in female Long Evans hooded rats. *Toxicol Sci* 93(1): 189-195.
- Hauser R, Calafat AM. 2005. Phthalates and human health. *Occup Environ Med* 62(11): 806-818.
- Heudorf U, Mersch-Sundermann V, Angerer J. 2007. Phthalates: toxicology and exposure. *Int J Hyg Environ Health* 210(5): 623-634.
- Hoppin JA, Brock JW, Davis BJ, Baird DD. 2002. Reproducibility of urinary phthalate metabolites in first morning urine samples. *Environ Health Perspect* 110(5): 515-518.
- Hornung R, Reed L. 1990. Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg* 5: 46-51.
- Ji J, Liu Y, Tong XH, Luo L, Ma J, Chen Z. 2013. The optimum number of oocytes in IVF treatment: an analysis of 2455 cycles in China. *Hum Reprod* 28(10): 2728-2734.
- Kelley KE, Hernandez-Diaz S, Chaplin EL, Hauser R, Mitchell AA. 2012. Identification of phthalates in medications and dietary supplement formulations in the United States and Canada. *Environ Health Perspect* 120(3): 379-384.

- Koch HM, Preuss R, Angerer J. 2006. Di(2-ethylhexyl)phthalate (DEHP): human metabolism and internal exposure-- an update and latest results. *Int J Androl* 29(1): 155-165; discussion 181-155.
- Lacasana M, Lopez-Flores I, Rodriguez-Barranco M, Aguilar-Garduno C, Blanco-Munoz J, Perez-Mendez O, et al. 2010. Association between organophosphate pesticides exposure and thyroid hormones in floriculture workers. *Toxicol Appl Pharmacol* 243(1): 19-26.
- Lovekamp-Swan T, Davis BJ. 2003. Mechanisms of phthalate ester toxicity in the female reproductive system. *Environ Health Perspect* 111(2): 139-145.
- Lovekamp TN, Davis BJ. 2001. Mono-(2-ethylhexyl) phthalate suppresses aromatase transcript levels and estradiol production in cultured rat granulosa cells. *Toxicol Appl Pharmacol* 172(3): 217-224.
- Meeker JD, Barr DB, Hauser R. 2006. Thyroid hormones in relation to urinary metabolites of non-persistent insecticides in men of reproductive age. *Reprod Toxicol* 22(3): 437-442.
- Meeker JD, Barr DB, Hauser R. 2008. Human semen quality and sperm DNA damage in relation to urinary metabolites of pyrethroid insecticides. *Hum Reprod* 23(8): 1932-1940.
- Meeker JD, Calafat AM, Hauser R. 2009. Urinary metabolites of di(2-ethylhexyl) phthalate are associated with decreased steroid hormone levels in adult men. *J Androl* 30(3): 287-297.
- Meeker JD, Missmer SA, Cramer DW, Hauser R. 2007a. Maternal exposure to second-hand tobacco smoke and pregnancy outcome among couples undergoing assisted reproduction. *Hum Reprod* 22(2): 337-345.
- Meeker JD, Missmer SA, Vitonis AF, Cramer DW, Hauser R. 2007b. Risk of spontaneous abortion in women with childhood exposure to parental cigarette smoke. *Am J Epidemiol* 166(5): 571-575.
- Meeker JD, Singh NP, Ryan L, Duty SM, Barr DB, Herrick RF, et al. 2004. Urinary levels of insecticide metabolites and DNA damage in human sperm. *Hum Reprod* 19(11): 2573-2580.
- Mendiola J, Jorgensen N, Andersson AM, Calafat AM, Silva MJ, Redmon JB, et al. 2011. Associations between urinary metabolites of di(2-ethylhexyl) phthalate and reproductive hormones in fertile men. *Int J Androl* 34(4pt1): 369-378.
- Mok-Lin E, Ehrlich S, Williams PL, Petrozza J, Wright DL, Calafat AM, et al. 2010. Urinary bisphenol A concentrations and ovarian response among women undergoing IVF. *Int J Androl* 33(2): 385-393.

- Nie L, Chu H, Liu C, Cole SR, Vexler A, Schisterman EF. 2010. Linear regression with an independent variable subject to a detection limit. *Epidemiology* 21 Suppl 4: S17-24.
- Pan G, Hanaoka T, Yoshimura M, Zhang S, Wang P, Tsukino H, et al. 2006. Decreased serum free testosterone in workers exposed to high levels of di-n-butyl phthalate (DBP) and di-2-ethylhexyl phthalate (DEHP): a cross-sectional study in China. *Environ Health Perspect* 114(11): 1643-1648.
- Peck JD, Sweeney AM, Symanski E, Gardiner J, Silva MJ, Calafat AM, et al. 2010. Intra- and inter-individual variability of urinary phthalate metabolite concentrations in Hmong women of reproductive age. *J Expo Sci Environ Epidemiol* 20(1): 90-100.
- Peppone LJ, Piazza KM, Mahoney MC, Morrow GR, Mustian KM, Palesh OG, et al. 2009. Associations between adult and childhood secondhand smoke exposures and fecundity and fetal loss among women who visited a cancer hospital. *Tob Control* 18(2): 115-120.
- Perry MJ, Venners SA, Barr DB, Xu X. 2007. Environmental pyrethroid and organophosphorus insecticide exposures and sperm concentration. *Reprod Toxicol* 23(1): 113-118.
- Pinnelli A, Di Cesare M. 2005. Human fertility: sociodemographic aspects. *Contraception* 72(4): 303-307.
- Preau JL, Jr., Wong LY, Silva MJ, Needham LL, Calafat AM. 2010. Variability over 1 week in the urinary concentrations of metabolites of diethyl phthalate and di(2-ethylhexyl) phthalate among eight adults: an observational study. *Environ Health Perspect* 118(12): 1748-1754.
- Society for Assisted Reproductive Technologies (SART). 2013. Clinical Summary Report: All SART Member Clinics. Available: https://www.sartcorsonline.com/rptCSR_PublicMultYear.aspx?ClinicPKID=0 [accessed October 26 2015].
- Schindler AE. 2004. First trimester endocrinology: consequences for diagnosis and treatment of pregnancy failure. *Gynecol Endocrinol* 18(1): 51-57.
- Silva MJ, Reidy JA, Preau JL, Samandar E, Needham LL, Calafat AM. 2006. Measurement of eight urinary metabolites of di(2-ethylhexyl) phthalate as biomarkers for human exposure assessment. *Biomarkers* 11(1): 1-13.
- Silva MJ, Samandar E, Preau JL, Jr., Reidy JA, Needham LL, Calafat AM. 2007. Quantification of 22 phthalate metabolites in human urine. *J Chromatogr B Analyt Technol Biomed Life Sci* 860(1): 106-112.

- Smith KW, Braun JM, Williams PL, Ehrlich S, Correia KF, Calafat AM, et al. 2012. Predictors and variability of urinary paraben concentrations in men and women, including before and during pregnancy. *Environ Health Perspect* 120(11): 1538-1543.
- Sunkara SK, Rittenberg V, Raine-Fenning N, Bhattacharya S, Zamora J, Coomarasamy A. 2011. Association between the number of eggs and live birth in IVF treatment: an analysis of 400 135 treatment cycles. *Hum Reprod* 26(7): 1768-1774.
- Svechnikova I, Svechnikov K, Soder O. 2007. The influence of di-(2-ethylhexyl) phthalate on steroidogenesis by the ovarian granulosa cells of immature female rats. *J Endocrinol* 194(3): 603-609.
- Toft G, Jonsson BA, Lindh CH, Jensen TK, Hjollund NH, Vested A, et al. 2012. Association between pregnancy loss and urinary phthalate levels around the time of conception. *Environ Health Perspect* 120(3): 458-463.
- Veeck LL, Zaninovic N. 2003. *An atlas of human blastocysts*. Boca Raton, FL Parthenon Publishing Group.
- Wilcox AJ, Weinberg CR, O'Connor JF, Baird DD, Schlatterer JP, Canfield RE, et al. 1988. Incidence of early loss of pregnancy. *N Engl J Med* 319(4): 189-194.
- Xia Y, Cheng S, Bian Q, Xu L, Collins MD, Chang HC, et al. 2005. Genotoxic effects on spermatozoa of carbaryl-exposed workers. *Toxicol Sci* 85(1): 615-623.
- Zota AR, Calafat AM, Woodruff TJ. 2014. Temporal trends in phthalate exposures: findings from the National Health and Nutrition Examination Survey, 2001-2010. *Environ Health Perspect* 122(3): 235-241.

Table 1. Demographic characteristics and primary SART diagnosis among 256 women in the Environment and Reproductive Health Study enrolled between 2004 and 2012.

Characteristic	N (%)
Age at study entry (years)	
Mean \pm SD	35.3 \pm 3.93
Range	21 - 43
Age \geq 37	95 (37%)
BMI (kg/m ²)	
Mean \pm SD	24.1 \pm 4.34
Range	16.1 – 42.4
Underweight or normal (< 25)	180 (70%)
Overweight or obese (\geq 25)	76 (30%)
Smoking	
Never smoked	185 (72%)
Ever smoked	
Current smoker	6 (2%)
Former smoker	65 (26%)
Race	
Caucasian	211 (82%)
Black/African American	6 (2%)
Asian	21 (8%)
Other	18 (7%)
Primary SART diagnosis at study entry	
Female factor	77 (30%)
Diminished ovarian reserve	18 (7%)
Ovulation disorders	22 (9%)
Endometriosis	17 (7%)
Uterine disorders	2 (1%)
Tubal factor	18 (7%)
Male infertility	95 (37%)
Unexplained	84 (33%)
Year at study entry	
2004-2006	49 (19%)
2007-2009	115 (45%)
2010-2012	92 (36%)

Abbreviations: BMI, body mass index; SD, standard deviation; SART, Society for Assisted Reproductive Technology (SART 2013);

Table 2. Cycle-specific clinical characteristics from 375 IVF cycles among 256 women in the Environment and Reproductive Health Study enrolled between 2004 and 2012.

Cycle-Specific Characteristics (N=375 IVF cycles)	N (%) or mean ± SD (range)
Treatment Protocol	
Luteal Phase	251 (67%)
Flare	77 (20%)
Antagonist	47 (13%)
Number of Embryos Transferred	
No embryos transferred	38 (10%)
1 embryo	46 (12%)
2 embryos	206 (55%)
3+ embryos	85 (23%)
Embryo Transfer Day	
No embryos transferred	38 (10%)
Day 2	18 (5%)
Day 3	215 (57%)
Day 5	104 (28%)
Fertilization Protocol (N=357)	
ICSI	196 (55%)
Traditional IVF	161 (45%)
Controlled Ovarian Hyperstimulation Outcomes (N = 357 IVF cycles^a)	
Day 3 FSH (IU/L)	7.15 ± 2.17 (0.2 – 15.2)
Peak Estradiol (pg/mL)	2071 ± 853 (551 – 5263)
Total Number of Oocytes Retrieved	10.91 ± 5.41 (1 – 32)
Mature (MII) Oocytes retrieved	9.20 ± 4.65 (0 – 27)
Maturation Rate (MII oocytes/total oocytes retrieved)	0.86 ± 0.16 (0 – 1)
Normal (2PN) Fertilized Oocytes	6.37 ± 3.66 (0 – 21)
Fertilization Rate (2PN/MII oocytes)	0.69 ± 0.24 (0 – 1)
Total Embryos	5.99 ± 3.75 (0 – 21)
Best Embryos	1.48 ± 1.90 (0 – 13)
Pregnancy Outcome (N = 375 IVF cycles)	
No oocytes retrieved	18 (4.8%)
Fertilization failure	9 (2.4%)
Arrested embryo development or all embryos frozen	11 (2.9%)
Implantation failure ^b	135 (36.0%)
Chemical pregnancy ^c	21 (5.6%)
Ectopic pregnancy	4 (1.1%)
Spontaneous abortion	27 (7.2%)
Therapeutic abortion	1 (0.3%)
Stillbirth	2 (0.5%)
Live Birth	147 (39.2%)

Abbreviations: FSH, follicle stimulating hormone; ICSI, intra-cytoplasmic sperm injection; IVF, in vitro fertilization; MII, mature oocytes;

^aFresh IVF cycles with successful egg retrieval (missing cycles include cycles failing prior to egg retrieval).

^bImplantation failure was defined as a negative pregnancy test (β hCG < 6 mIU/ml) 17 days following embryo transfer or insemination.

^cChemical pregnancy was defined as implantation with no subsequent clinical pregnancy.

Table 3. Distribution of urinary phthalate metabolite concentrations (metabolite or molar sum) measured among 256 women in the Environment and Reproductive Health Study enrolled between 2004 and 2012.

Phthalate	Units	N ^b	LOD	% Detect ^c	SG-Adjusted			Unadjusted ^a		
					Min	Median (IQR)	Max	Min	Median (IQR)	Max
∑DEHP metabolites	μmol/L	375	--	--	0.020	0.19 (0.10, 0.42)	5.60	0.009	0.20 (0.09, 0.44)	7.52
MEHP	μg/L	375	0.5-1.2	77.1%	0.28	2.88 (1.37, 6.87)	99.3	0.50	2.72 (1.30, 6.91)	130
MEHHP	μg/L	375	0.2-0.7	99.7%	0.62	15.7 (7.75, 35.0)	561	0.47	16.3 (7.07, 38.2)	582
MEOHP	μg/L	375	0.2-0.7	99.6%	0.90	10.5 (5.48, 25.4)	306	0.30	11.3 (4.92, 25.7)	387
MECPP	μg/L	375	0.2-0.6	96.7%	3.51	26.3 (14.6, 57.2)	761	1.40	29.2 (12.9, 62.9)	1190
MEP	μg/L	375	0.4-0.8	100%	2.40	49.3 (21.5, 129)	3481	0.80	52.2 (20.8, 133)	2537
MBP	μg/L	375	0.4-0.6	97.0%	1.10	12.9 (7.32, 20.8)	435	0.40	12.6 (6.40, 26.3)	250
MCPP	μg/L	375	0.1-0.2	96.7%	0.31	3.22 (1.80, 6.64)	202	0.18	3.63 (1.77, 7.76)	179
MiBP	μg/L	375	0.2-0.3	96.7%	0.44	6.66 (3.50, 12.0)	49.7	0.20	7.03 (3.14, 14.6)	76.3
MBzP	μg/L	375	0.2-0.3	95.5%	0.27	3.44 (1.75, 7.35)	121	0.19	3.99 (1.57, 7.60)	189
MCOP	μg/L	337	0.2-0.7	98.6%	1.28	22.8 (9.10, 57.7)	1350	0.69	22.7 (8.00, 66.2)	2250
MCNP	μg/L	337	0.2-0.6	96.6%	0.79	4.58 (2.51, 7.97)	281	0.40	4.75 (2.41, 9.18)	281

Abbreviations: IQR, interquartile range; LOD, limit of detection; min, minimum; max, maximum; SG, specific gravity; N, number of IVF cycles.

^aUnadjusted concentrations presented to facilitate comparison with other studies.

^bGeometric mean of up to 2 urinary phthalate concentrations from each IVF cycle was used.

^cPercent of phthalate metabolite concentrations above the detection limits (N=673 total urine samples except for MCOP and MCNP which have 646 urine samples); Total samples <LOD by analyte: MEHP=154, MEHHP=2; MEOHP=3, MECPP=22, MEP=0, MBP=20, MCPP=22, MiBP=22, MBzP=30, MCOP=9, MCNP=22.

Table 4. Unadjusted and adjusted mean proportion of cycles resulting in implantation, clinical pregnancy, and live birth by quartile of urinary DEHP metabolites and Σ DEHP concentrations among 256 women undergoing 375 IVF cycles.

Phthalate Measure Quartiles	Mean Proportion (95% Confidence Intervals)					
	Implantation		Clinical Pregnancy		Live Birth	
	Unadjusted	Adjusted	Unadjusted	Adjusted	Unadjusted	Adjusted
ΣDEHP Metabolites (μmol/L)						
Q1(0.02-0.10)	0.63 (0.52, 0.72)	0.62 (0.50, 0.72)	0.59 (0.48, 0.69)	0.57 (0.45, 0.67)	0.50 (0.39, 0.60)	0.47 (0.36, 0.58)
Q2 (0.11-0.19)	0.54 (0.44, 0.65)	0.55 (0.44, 0.65)	0.46 (0.36, 0.57)	0.46 (0.36, 0.57)	0.44 (0.33, 0.54)	0.43 (0.32, 0.54)
Q3 (0.20-0.42)	0.51 (0.41, 0.62)	0.52 (0.41, 0.63)	0.48 (0.38, 0.59)	0.49 (0.38, 0.59)	0.39 (0.29, 0.50)	0.39 (0.29, 0.50)
Q4 (0.43-5.60)	0.48 (0.38, 0.59)*	0.49 (0.39, 0.60)	0.38 (0.28, 0.48)*	0.38 (0.28, 0.49)*	0.28 (0.20, 0.39)*	0.28 (0.19, 0.39)*
P-trend ^a	0.09	0.18	0.02	0.04	0.005	0.01
MEHP (μg/L)						
Q1 (0.28-1.36)	0.61 (0.50, 0.70)	0.60 (0.49, 0.70)	0.55 (0.45, 0.65)	0.54 (0.43, 0.64)	0.47 (0.37, 0.58)	0.45 (0.35, 0.56)
Q2 (1.37-2.88)	0.57 (0.46, 0.66)	0.56 (0.45, 0.66)	0.53 (0.42, 0.63)	0.52 (0.41, 0.62)	0.46 (0.35, 0.56)	0.44 (0.34, 0.55)
Q3 (2.89-6.80)	0.47 (0.37, 0.57)	0.48 (0.37, 0.58)	0.43 (0.33, 0.53)	0.43 (0.33, 0.54)	0.37 (0.27, 0.48)	0.36 (0.27, 0.47)
Q4 (6.87-99.3)	0.52 (0.42, 0.63)	0.54 (0.43, 0.65)	0.40 (0.30, 0.50)*	0.41 (0.31, 0.52)	0.30 (0.22, 0.41)*	0.30 (0.21, 0.41)
P-trend	0.43	0.68	0.04	0.09	0.02	0.04
MEHHP(μg/L)						
Q1 (0.62-7.74)	0.61 (0.50, 0.70)	0.58 (0.47, 0.69)	0.58 (0.47, 0.68)	0.55 (0.44, 0.66)	0.58 (0.47, 0.68)	0.45 (0.34, 0.56)
Q2 (7.75-15.4)	0.57 (0.46, 0.66)	0.57 (0.47, 0.67)	0.49 (0.39, 0.60)	0.50 (0.39, 0.60)	0.49 (0.39, 0.60)	0.46 (0.36, 0.57)
Q3 (15.5-34.5)	0.51 (0.41, 0.62)	0.52 (0.42, 0.63)	0.46 (0.36, 0.57)	0.47 (0.36, 0.58)	0.46 (0.36, 0.57)	0.35 (0.25, 0.46)
Q4 (34.6-561.3)	0.48 (0.38, 0.58)	0.49 (0.39, 0.60)	0.38 (0.28, 0.48)*	0.38 (0.28, 0.49)*	0.38 (0.28, 0.48)*	0.30 (0.21, 0.41)
P-trend	0.11	0.22	0.01	0.04	0.01	0.03
MEOHP(μg/L)						
Q1 (0.90-5.44)	0.60 (0.49, 0.69)	0.58 (0.47, 0.68)	0.58 (0.47, 0.68)	0.56 (0.44, 0.66)	0.49 (0.39, 0.60)	0.47 (0.36, 0.58)
Q2 (5.45-10.4)	0.56 (0.45, 0.66)	0.57 (0.46, 0.67)	0.47 (0.37, 0.58)	0.48 (0.37, 0.59)	0.42 (0.32, 0.53)	0.42 (0.32, 0.53)
Q3 (10.5-24.7)	0.54 (0.43, 0.64)	0.54 (0.43, 0.64)	0.47 (0.37, 0.58)	0.47 (0.36, 0.58)	0.37 (0.27, 0.48)	0.36 (0.26, 0.47)
Q4 (24.8-306.0)	0.48 (0.38, 0.58)	0.50 (0.39, 0.60)	0.39 (0.29, 0.49)*	0.40 (0.30, 0.51)*	0.32 (0.23, 0.42)*	0.32 (0.22, 0.43)
P-trend	0.14	0.26	0.03	0.07	0.03	0.07
MECPP(μg/L)						

Q1 (3.51-14.5)	0.61 (0.50, 0.70)	0.59 (0.48, 0.69)	0.55 (0.45, 0.66)	0.54 (0.43, 0.64)	0.46 (0.36, 0.57)	0.43 (0.33, 0.54)
Q2 (14.6-26.2)	0.55 (0.44, 0.65)	0.55 (0.44, 0.66)	0.47 (0.37, 0.58)	0.48 (0.37, 0.58)	0.44 (0.34, 0.55)	0.43 (0.33, 0.54)
Q3 (26.3-56.6)	0.51 (0.41, 0.62)	0.52 (0.41, 0.62)	0.47 (0.37, 0.58)	0.47 (0.37, 0.58)	0.38 (0.28, 0.49)	0.37 (0.27, 0.48)
Q4 (56.7-760.5)	0.50 (0.40, 0.61)	0.51 (0.41, 0.62)	0.41 (0.31, 0.51)	0.42 (0.31, 0.52)	0.33 (0.24, 0.43)	0.32 (0.23, 0.43)
P-trend	0.25	0.40	0.09	0.17	0.06	0.12
MEP (µg/L)						
Q1 (2.40-21.4)	0.62 (0.52, 0.72)	0.60 (0.49, 0.70)	0.58 (0.48, 0.68)	0.55 (0.44, 0.66)	0.44 (0.34, 0.55)	0.40 (0.30, 0.51)
Q2 (21.5-49.2)	0.52 (0.42, 0.63)	0.53 (0.43, 0.64)	0.45 (0.35, 0.56)	0.46 (0.35, 0.56)	0.38 (0.29, 0.49)	0.38 (0.28, 0.49)
Q3 (49.3-128.1)	0.51 (0.41, 0.61)	0.52 (0.41, 0.62)	0.44 (0.34, 0.54)	0.44 (0.34, 0.55)	0.37 (0.27, 0.47)	0.36 (0.26, 0.47)
Q4 (128.2-3481)	0.50 (0.40, 0.60)	0.52 (0.42, 0.63)	0.43 (0.33, 0.53)*	0.45 (0.34, 0.55)	0.40 (0.30, 0.51)	0.42 (0.31, 0.53)
P-trend	0.25	0.55	0.17	0.40	0.87	0.65
MBP (µg/L)						
Q1 (1.10-7.30)	0.63 (0.52, 0.72)	0.61 (0.50, 0.71)	0.55 (0.45, 0.65)	0.53 (0.42, 0.63)	0.48 (0.37, 0.58)	0.44 (0.34, 0.55)
Q2 (7.31-12.8)	0.57 (0.46, 0.66)	0.57 (0.46, 0.67)	0.50 (0.40, 0.60)	0.50 (0.40, 0.61)	0.41 (0.31, 0.52)	0.41 (0.31, 0.51)
Q3 (12.9-20.8)	0.47 (0.37, 0.57)*	0.48 (0.38, 0.59)	0.40 (0.30, 0.50)*	0.41 (0.31, 0.51)	0.36 (0.27, 0.47)	0.36 (0.27, 0.47)
Q4 (20.9-435.0)	0.50 (0.40, 0.60)	0.51 (0.41, 0.62)	0.45 (0.35, 0.55)	0.46 (0.36, 0.57)	0.35 (0.25, 0.45)	0.35 (0.25, 0.46)
P-trend	0.10	0.22	0.17	0.36	0.09	0.19
MiBP (µg/L)						
Q1 (0.44-3.50)	0.63 (0.52, 0.72)	0.62 (0.51, 0.72)	0.54 (0.44, 0.64)	0.53 (0.42, 0.63)	0.47 (0.36, 0.57)	0.44 (0.34, 0.55)
Q2 (3.51-6.63)	0.53 (0.43, 0.63)	0.53 (0.42, 0.63)	0.49 (0.39, 0.60)	0.48 (0.38, 0.59)	0.40 (0.30, 0.51)	0.39 (0.29, 0.49)
Q3 (6.64-12.0)	0.49 (0.39, 0.59)	0.51 (0.40, 0.62)	0.43 (0.33, 0.53)	0.44 (0.34, 0.55)	0.36 (0.27, 0.47)	0.37 (0.27, 0.48)
Q4 (12.1-49.7)	0.51 (0.41, 0.62)	0.52 (0.41, 0.63)	0.44 (0.34, 0.55)	0.44 (0.34, 0.55)	0.37 (0.27, 0.48)	0.36 (0.26, 0.47)
P-trend	0.20	0.30	0.18	0.27	0.23	0.30
MBzP (µg/L)						
Q1(0.27-1.74)	0.55 (0.45, 0.65)	0.54 (0.43, 0.65)	0.50 (0.39, 0.60)	0.48 (0.38, 0.59)	0.41 (0.32, 0.52)	0.39 (0.29, 0.50)
Q2 (1.75-3.43)	0.58 (0.47, 0.68)	0.58 (0.47, 0.68)	0.48 (0.38, 0.59)	0.47 (0.37, 0.58)	0.37 (0.28, 0.48)	0.35 (0.25, 0.46)
Q3 (3.44-7.334)	0.54 (0.43, 0.64)	0.55 (0.45, 0.66)	0.48 (0.38, 0.59)	0.50 (0.39, 0.61)	0.46 (0.35, 0.56)	0.47 (0.37, 0.58)
Q4 (7.35-120.9)	0.50 (0.40, 0.61)	0.50 (0.40, 0.61)	0.44 (0.34, 0.55)	0.44 (0.33, 0.55)	0.36 (0.26, 0.47)	0.34 (0.25, 0.45)
P-trend	0.37	0.44	0.44	0.51	0.53	0.59
MCPP (µg/L)						
Q1 (0.31-1.80)	0.60 (0.50, 0.70)	0.60 (0.49, 0.70)	0.53 (0.43, 0.63)	0.52 (0.41, 0.63)	0.45 (0.34, 0.55)	0.43 (0.33, 0.54)
Q2 (1.81-3.19)	0.50 (0.40, 0.61)	0.50 (0.40, 0.61)	0.43 (0.33, 0.53)	0.42 (0.32, 0.53)	0.40 (0.30, 0.51)	0.39 (0.29, 0.50)
Q3 (3.20-6.58)	0.53 (0.42, 0.63)	0.53 (0.42, 0.63)	0.47 (0.37, 0.58)	0.47 (0.36, 0.57)	0.42 (0.32, 0.53)	0.41 (0.31, 0.52)

Q4 (6.59-202.2)	0.53 (0.43, 0.63)	0.55 (0.44, 0.65)	0.47 (0.37, 0.58)	0.48 (0.38, 0.59)	0.33 (0.24, 0.43)	0.32 (0.23, 0.43)
P-trend	0.67	0.87	0.79	0.98	0.11	0.16
MCOP (µg/L)^b						
Q1 (1.28-8.93)	0.54 (0.43, 0.64)	0.54 (0.43, 0.65)	0.49 (0.38, 0.60)	0.50 (0.39, 0.61)	0.46 (0.35, 0.57)	0.47 (0.36, 0.58)
Q2 (8.94-22.2)	0.55 (0.44, 0.65)	0.54 (0.43, 0.65)	0.48 (0.37, 0.59)	0.47 (0.36, 0.58)	0.40 (0.30, 0.51)	0.37 (0.27, 0.49)
Q3 (22.3-57.7)	0.58 (0.47, 0.68)	0.57 (0.46, 0.68)	0.48 (0.38, 0.59)	0.47 (0.36, 0.58)	0.40 (0.29, 0.51)	0.38 (0.27, 0.49)
Q4 (57.8-1350.0)	0.52 (0.42, 0.63)	0.53 (0.42, 0.64)	0.47 (0.36, 0.57)	0.47 (0.36, 0.58)	0.35 (0.25, 0.47)	0.34 (0.24, 0.46)
P-trend	0.79	0.85	0.78	0.83	0.23	0.25
MCNP (µg/L)^b						
Q1 (0.79-2.51)	0.52 (0.42, 0.63)	0.52 (0.41, 0.63)	0.44 (0.34, 0.55)	0.43 (0.33, 0.55)	0.42 (0.31, 0.53)	0.41 (0.30, 0.52)
Q2 (2.52-4.52)	0.56 (0.45, 0.66)	0.56 (0.45, 0.67)	0.50 (0.39, 0.61)	0.50 (0.39, 0.61)	0.41 (0.31, 0.53)	0.40 (0.29, 0.51)
Q3 (4.53-7.97)	0.58 (0.47, 0.68)	0.59 (0.47, 0.69)	0.51 (0.40, 0.61)	0.51 (0.40, 0.62)	0.40 (0.29, 0.51)	0.39 (0.28, 0.50)
Q4 (7.98-281.3)	0.52 (0.42, 0.63)	0.52 (0.41, 0.63)	0.47 (0.36, 0.57)	0.46 (0.35, 0.57)	0.39 (0.28, 0.50)	0.37 (0.27, 0.49)
P-trend	0.83	0.81	0.98	0.98	0.68	0.70

Adjusted models control for maternal age (continuous), body mass index (continuous), smoking status (never, ever), and primary SART infertility diagnosis at study entry (female, male, unexplained).

^aTests for linear trend were performed using the median level of urinary phthalate metabolite in each quartile as a continuous variable in the model.

^b38 missing values for MCOP and MCNP

*p-value for comparison for a specific quartile against Q1 < 0.05

Figure Legends

Figure 1. Overview of *in vitro* fertilization (IVF) outcomes of 256 women (375 cycles) in the Environment and Reproductive Health Study enrolled between 2004 and 2012. Abbreviations: IUI, intrauterine insemination; IVF, *in vitro* fertilization; TAB, therapeutic abortion; SAB, spontaneous abortion; SB, stillbirth.

Figure 2. Adjusted mean (95% CI) total, mature (MII), and fertilized oocyte count by quartile of urinary phthalate metabolites and Σ DEHP concentrations among 247 women undergoing 357 IVF cycles with successful egg retrieval. Adjusted models control for maternal age (continuous), body mass index (continuous), smoking status (never, ever), and primary SART infertility diagnosis at study entry (female, male, unexplained). An asterisk represents a p-value for comparison against Q1 < 0.05. Panel A) Σ DEHP, Panel B) MEHP, Panel C) MEHHP, Panel D) MEOHP, Panel E) MECPP, Panel F) MCOP, Panel G) MCNP

Figure 1.







Figure 2.



E.



F.



G.

