



ENVIRONMENTAL HEALTH PERSPECTIVES

<http://www.ehponline.org>

Maternal Occupational Exposure to Noise during Pregnancy and Hearing Dysfunction in Children: A Nationwide Prospective Cohort Study in Sweden

Jenny Selander, Maria Albin, Ulf Rosenhall, Lars Rylander,
Marie Lewné, and Per Gustavsson

<http://dx.doi.org/10.1289/ehp.1509874>

Received: 24 February 2015

Accepted: 23 November 2015

Advance Publication: 8 December 2015

Note to readers with disabilities: *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.



Maternal Occupational Exposure to Noise during Pregnancy and Hearing Dysfunction in Children: A Nationwide Prospective Cohort Study in Sweden

Jenny Selander¹, Maria Albin^{1,2,3}, Ulf Rosenhall⁴, Lars Rylander³, Marie Lewné^{1,2}, and Per Gustavsson^{1,2}

¹Institute of Environmental Medicine, Unit of Occupational Medicine, Karolinska Institutet, Stockholm, Sweden; ²Centre for Occupational and Environmental Medicine at Stockholm County Council, Stockholm, Sweden; ³Division of Occupational and Environmental Medicine, Lund University, Lund, Sweden; ⁴Department of Audiology and Neurotology, Karolinska University Hospital, Stockholm, Sweden

Address correspondence to Jenny Selander, Institute of Environmental Medicine, Unit of Occupational Medicine, Karolinska Institutet, Solnavägen 4, 113 65 Stockholm, Sweden.

Telephone: 0046-8-52480029, 0046-73-9795476. E-mail: Jenny.Selander@ki.se

Running title: Noise during pregnancy & hearing dysfunction in children

Acknowledgment: We express our gratitude for the work by Gun Nise (deceased) on the noise exposure matrix. In addition, we thank the National Board of Health and Welfare, the Swedish Social Insurance Agency, and Statistics Sweden for valuable collaboration during the data collection and The Swedish Research Council for Health, Working Life and Welfare (FORTE) for funding this study (grant no 2009-0801).

Competing financial interests: The authors declare they have no actual or potential competing financial interests.

Abstract

Background: Many women of childbearing age are occupationally active which leads to a large number of pregnancies potentially exposed to occupational exposures. Occupational noise has been identified as a risk factor for hearing impairment in adults. However, very few studies have assessed the effect of occupational noise on the fetus.

Objectives: The aim of this study was to investigate if occupational exposure to noise during pregnancy is associated with hearing dysfunction in children.

Methods: This population based cohort study included 1 422 333 single births in Sweden 1986 - 2008. Data on mothers' occupation, smoking habits, age, ethnicity, BMI, leave of absence and socio-economic factors were obtained from interviews performed by prenatal care unit staff at approximately 10 weeks of gestation, and from national registers. Occupational noise exposure was classified by a job-exposure-matrix as <75 , 75-84, or ≥ 85 $\text{dB}_{\text{LAeq},8\text{h}}$. Diagnosed cases of hearing dysfunction (ICD10 H90.3-7, 91.0, 91.2-3, 91.8, 93.1-2) were identified from a register of specialized medical care. Cox proportional hazards models were used to estimate associations.

Results: In the full sample, containing a mixture of part-time and full time workers during pregnancy, the adjusted HR for hearing dysfunction associated with maternal occupational noise exposure ≥ 85 vs <75 $\text{dB}_{\text{LAeq},8\text{h}}$ was 1.27 (95% CI: 0.99, 1.64; 60 exposed cases). When restricted to children whose mothers worked full-time, and had <20 days leave of absence during pregnancy, the corresponding HR was 1.82 (95% CI 1.08, 3.08; 14 exposed cases).

Conclusions: This study showed an association between occupational noise exposure during pregnancy and hearing dysfunction in children. In view of mechanistic evidence and earlier indicative epidemiological and experimental findings, the results support that pregnant women should not be exposed to high levels of noise at work.

Introduction

A majority of women of child-bearing age in the industrialized part of the world are occupationally active today. In total, 73.7% of Swedish women of working age were employed in 2014 (OECD 2014). This leads to a high number of pregnancies potentially exposed to various occupational hazards. In Sweden, 15% of employed women report exposure to noise during at least $\frac{1}{4}$ of the working day, so loud that they could not have a normal conversation (Arbetsmiljöverket - the Swedish Work Environment Authority 2012). Many studies have shown that occupational noise exposure causes hearing impairment in adults (Verbeek et al. 2012). However, very little is known about the association between occupational noise exposure during fetal life and hearing impairment in the child.

Sound is transmitted from the air over the abdominal wall and the uterus to the fetal head during pregnancy. The noise stimulates the inner ear through a soft tissue conduction route and could potentially affect the hearing of the fetus by damaging inner and outer hair cells within the cochlea, especially since the maturing cochlea is more sensitive to ototraumatic factors than the adult one (Brundin et al. 1989; Chordekar et al. 2012; Gerhardt and Abrams 2000). Both animal and human experimental studies show that the attenuation of noise through the passage of the abdominal wall and the uterus is strongly dependent on frequency. While the fetus is well protected from high-frequency noise, low-frequency noise can even be amplified during the passage of the abdominal wall and the amniotic fluid (Chordekar et al. 2012; Gerhardt and Abrams 2000; Saunders and Chen 1982). In addition, animal experimental studies indicate that intense and sustained noise exposure during pregnancy can induce hearing dysfunction in the offspring, in guinea pigs (Cook et al. 1982), and in sheep (Gerhardt and Abrams 2000; Griffiths et al. 1994; Huang et al. 1997). Three previous small epidemiological cross-sectional studies

have investigated occupational noise exposure during pregnancy and hearing impairment in children, including a study of 131 4-10 year old children in Quebec, a study of 75 children 10-14 years old in France and a study of 80 children 0-6 months old in Brazil (Daniel and Laciak 1982; Lalande et al. 1986; Rocha et al. 2007). Although the studies conducted in Canada and France reported evidence supporting the hypothesis of an association between occupational noise exposures during pregnancy and hearing loss in children, the Brazilian study did not report an association.. However, the studies were all based on small samples and had thereby low precision. The Brazilian study presented unspecific inclusion criteria ,the French study lacked a comparison group and the Canadian study had only two cases in the comparison group.

The aim of this study was to investigate if occupational exposure to noise during pregnancy is associated with an increased risk of hearing dysfunction in children in a population-based nationwide study with individual data on occupation, hearing dysfunction, and potential confounding factors.

Methods

The FENIX study (fetal noise exposure) is a nationwide prospective cohort study based on births in Sweden between 1986 and 2008.

Data sources

Information on job title and if the woman was working full-time, part-time, or not working at all was collected at the registration interview for prenatal care (gestational week 10) together with background characteristics such as maternal age, smoking habits, family structure, and nationality. It was entered in the Medical Birth Register, together with information from records from the maternity wards regarding parity, gender, and date of birth. The Swedish Medical Birth

Register includes 98-99% of all children born in Sweden (Socialstyrelsen - the Swedish National Board of Health and Welfare 2002).

Information on hearing dysfunction was retrieved from the patient register, which includes diagnoses from all out-patient clinics for specialist care in Sweden from 2003 and onwards. The overall coverage in the patient register is approximately 80% (Socialstyrelsen - the Swedish National Board of Health and Welfare 2013), but has not been specifically evaluated for hearing dysfunction.

Data on the mothers highest completed education at the year of the child's birth was gathered through the population based nationwide register (LISA) holding information on all Swedish citizens at the age of 16 or more. . (Statistiska centralbyrån - Statistics Sweden 2011).

Individual information on leave of absence during each pregnancy was retrieved from the Social Insurance MIDAS database and includes information on days of sick-leave and parental leave for the time period 1986-2008. The register has a complete coverage of long-term leave of absence, since every citizen needs to register sick-leave or parental leave in order to receive payment from the Social Insurance Agency, but it does not cover short term sick-leave (≤ 14 consecutive days).

However, parental leave and special sick-leave related to the pregnancy is reported from day one (Försäkringskassan - the Swedish Social Insurance Agency 2011).

The registers were matched with the personal identification number, unique to every Swedish citizen.

Exposure

Job titles were recorded at the registration interview held at the prenatal care facilities in Sweden. The occupation was recorded as free text (not coded) in the Medical Birth Register. Out of the 2,348,250 births eligible for the study (children born between 1986 and 2008 in Sweden), 1,957,189 had data on occupation. The entries were processed down into 64,398 unique occupational titles by removing other characters than letters, by harmonizing abbreviations, and correcting spelling errors. There were approximately 44,142 unique titles held by only one mother in the cohort, they were excluded from the study for reasons of limited resources for coding of occupations, leaving 20,256 job titles held by at least two mothers and covering 98% of the occupational titles. These were coded manually by an occupational hygienist and a safety engineer into a 6-digit AMSYK/SSYK-code based on ISCO-88 (International Labour Office (ILO) 1990) (Figure 1). Occupations that were difficult to assess were discussed among the two assessors until consensus was reached. However, 156,549 entries could not be assigned an occupational code since the information was too unspecific. In addition, 293,117 entries contained non-working groups such as students, unemployed, house wives, refugees (AMSYK-code X33 and X21). Since this study focus on occupational exposures, these mothers were excluded. , leaving 1,463,381 births in the study classified into three noise categories. The study was further restricted to single births and the final sample consisted of 1,422,333 births.

The validated job-exposure matrix has been constructed prior to this study and is presented in detail elsewhere (Sjostrom et al. 2013). Briefly, the job-exposure-matrix contains 321 job families and specifies the annual average of the daily 8h noise exposure levels in three exposures categories (<75, 75-84, ≥85 dBA) in 5-year calendar bands from 1970 to 2004. The noise exposure information used for the job-exposure-matrix derives from measurement reports

collected from occupational medicine clinics, occupational health services, and large companies all over Sweden. The highest occupational noise category $>85\text{dBA}$ is based on the occupational exposure limit for noise in Sweden, the lower interval $<75\text{dBA}$ includes occupations without a dominating noise source.

The occupational codes in this study were linked to the job-exposure-matrix that classified the 8 hour average exposure (L_{Aeq8h}) of each occupation into three categories: $<75\text{dBA}$, $75\text{-}84\text{dBA}$, $\geq 85\text{dBA}$ (figure 1). The job-exposure-matrix was matched in 5-year calendar bands to each occupational code so that the exposure was applied to births occurring within that time frame. It was not possible to classify exposure to low-frequency noise (dBC) separately due to lack of frequency-specific exposure data.

Outcome

We retrieved information on every child born during the period 1986-2008 that had been diagnosed in 2003-2008 with types of hearing dysfunctions that can be related to noise exposure during pregnancy in the patient register (ICD-10): Other diseases of inner ear, (H83.3, H83.8), Conductive mixed with sensorineural hearing loss and Sensorineural hearing loss, (H90.3-7), Other hearing loss, (H91.0, H91.2-3, H91.8), and other disorders of the ear, not elsewhere classified (including Tinnitus), (H93.1, H93.2). This selection is based on the well-established concept that noise-induced hearing loss is of the sensorineural type, caused primarily by a cochlear damage (Hackney and Furness 2007). Diagnoses indicating middle ear problems or conductive hearing loss have not been included, since they are not related to noise exposure. The total sample of all selected hearing disorders was analyzed together as one group. In addition, the two main diagnosis types: Sensorineural hearing loss and Tinnitus that accounted for most of the

cases were large enough to be analyzed separately. In these two sub groups, the same child could be included as a case in both the Sensorineural hearing loss analysis and the Tinnitus analysis if it had been diagnosed with both disorders. In, the main analysis of all cases combined, only the first diagnosis was counted.

Data analysis

Information on diagnoses was available only for the time period 2003 to 2008, while the cohort included all births from 1986. Thus, the chance of having a hearing dysfunction diagnosis varies with year of birth, and all analyses were therefore adjusted for birth year (22 categories).

All analyses were performed using Cox-regression with child's age as the underlying time scale. Person years for each child were calculated from the child's birth to the age at first diagnosis or age at the end of follow-up (31 December 2008), whichever came first. Potential confounders were identified through a review of previous studies on pregnancy outcomes and occupational exposures, by observing the effect each potential confounder had on the association between exposure and outcome and by placing the potential confounders in a causal web. All variables initially assessed: mothers age (quartiles), smoking (three categories: not smoking, 1-9 cigarettes per day, ≥ 10 cigarettes per day), Highest completed education (four categories: pre-high school, high school, university < 3years, university ≥ 3 years), nationality (four categories: Swedish, other European country, outside Europe, unknown), family structure (dichotomous: married/living together with the father, single mother), child's gender (dichotomous: male/female), birth year (quartiles) and parity (three categories: child 1, 2 or ≥ 3) were selected as covariates and are presented in table 1. Complete case analyses were performed and observations

with missing data for any model covariates were thereby excluded when estimating adjusted Hazard Ratios.

All analyses were restricted to single births and to employed women by excluding women who at the beginning of pregnancy (interview gestational week 10) said that they were students, unemployed, house wives, refugees, or in other non-working categories (AMSYK-code X33 and X21) during pregnancy. Women that in gestational week 10, stated an occupation that could be coded into occupational codes were included in the study. The presence at the work place was then evaluated and certain analyses were restricted regarding work participation by combining the answers to the question on current degree (work full time, part time or not at all right now) of occupational activity at beginning of pregnancy (interview at gestational week 10) with the registry data on days of absence (sick-leave, parental leave, pregnancy benefit) during the whole pregnancy. This was done by dividing the cohort into three categories: absent from work, part-time work and full-time work. Absent from work included mothers that stated an occupation but also stated that they did not work at the registration interview in gestational week 10 or mothers that stated an occupation but had more than 153 days (90th percentile) of leave of absence from work during pregnancy. Part-time workers were defined as mothers that stated that they worked part-time at the registration interview or had between 20 (median) and 153 days (90th percentile) of leave of absence during pregnancy. Full-time workers were defined as mothers stating that they worked full-time at the registration interview at the beginning of pregnancy and had less than 20 days (median) of sick-leave during pregnancy.

Statistical significance was determined by 95% confidence intervals. All analyses were performed with STATA SE 12 and the study was approved by the regional ethics committee in Stockholm.

Results

The study included 1,422,333 single births. Of these, 1,320,195 had complete data for the covariates of the adjusted model. In total, 12,668 cases of hearing dysfunction were identified and included in the study; Other diseases of inner ear, 133 cases (ICD10: H83.3, H83.8), Conductive mixed with sensorineural hearing loss and Sensorineural hearing loss, 8,696 cases (H90.3-7), Other hearing loss, 917 cases (H91.0, H91.2-3, H91.8), and other disorders of the ear, not elsewhere classified (including Tinnitus), 3,637 cases (H93.1, H93.2).

Background characteristics are presented in table 1. Compared with less exposed mothers (< 75 dBA), highly exposed mothers (≥ 85 dBA) tended to be younger (25% vs. 16% ≤ 24 years), less educated (21% vs. 8.7% < high school), and more likely to smoke during pregnancy (23% vs. 14%). The percentage of missing data in the background characteristics was low (0-10%). The three most frequent jobs in the high exposure group (≥ 85 dBA) were musicians (15%), carpenters and wood workers (15%), and butchers (12%).

An increased risk of hearing dysfunction among children was indicated after exposure to occupational noise during pregnancy, table 2. For all cases combined, adjusted HRs for 75–84 dBA and ≥ 85 dBA compared with < 75 dBA were 1.05 (95% CI: 1.00, 1.10) and 1.27 (95% CI: 0.99, 1.64), respectively. Corresponding estimates were similar for sensorineural hearing loss (HR = 1.08; 95% CI: 1.03, 1.15 and HR = 1.26; 95% CI: 0.93, 1.70). For Tinnitus, moderate noise was not associated with an increased risk (HR = 0.99; 95% CI: 0.90, 1.08), but the HR for high noise ≥ 85 dBA was similar to the other outcomes (HR = 1.25; 95% CI: 0.74, 2.12). There were 60, 42, and 14 highly exposed cases for all hearing dysfunction, sensorineural hearing loss, and tinnitus, respectively.

The risk of hearing dysfunction among children in relation to occupational noise during pregnancy, subdivided by presence at work during pregnancy is presented in table 3. The association was strongest when analyses were restricted to mothers who worked full-time during pregnancy with < 20 days of absence (362,572 total, 343,712 with complete data). The adjusted HR for occupational noise exposure ≥ 85 vs. < 75 dBA was 1.82 (95% CI: 1.08, 3.08) based on 14 exposed cases and 2,222 cases with low exposure. In contrast, corresponding HRs were 1.25 (95% CI: 0.91, 1.71) for high exposure among mothers classified as working part-time (37 exposed and 5,243 low-exposed cases) and 0.74 (95% CI: 0.35, 1.56) for women who had more than 153 days (90th percentile) of absence from work during pregnancy or who were not working at the time of the registration interview (7 exposed and 1,116 unexposed cases).

Discussion

An association between maternal occupational noise exposure > 85 dBA during pregnancy and hearing dysfunction among children was indicated. The association was more pronounced when restricting the study to full-time working mothers with less than 20 days leave of absence during pregnancy. The results are in line with two previous epidemiological cross-sectional studies on occupational noise exposure and hearing impairment that reported an increased risk of hearing impairment after exposure to high occupational noise ≥ 85 dBA during pregnancy (Daniel and Laciak 1982; Lalande et al. 1986). In addition, animal studies have shown an association between noise exposure during pregnancy and hearing loss in offspring (Cook et al. 1982; Griffiths et al. 1994).

In the human fetus as well as in precocial mammals, the auditory system is functional well before birth (Armitage et al. 1980). The peripheral auditory system, including the cochlea, has an

adult anatomical appearance by the 20th gestational week (Pujol et al. 1990) and the fetus responds to auditory stimulation at that time (Hepper and Shahidullah 1994). However, the maturing cochlea is more sensitive to ototraumatic factors than the fully developed cochlea (Saunders and Chen 1982). The sound transmission mechanisms after birth are not applicable to the fetal cochlea. Instead, the sound stimulates the inner ear through a soft tissue conduction route (Chordekar et al. 2012). The sound passes through the abdominal wall, the uterine wall and the amniotic fluid before reaching the fetal head, where a bone conduction route through the cartilaginous fetal skull bones is activated. Experimental studies both in sheep and humans show that mid- and high-frequency sounds are attenuated by the abdominal soft tissues (>20 dB reduction) (Armitage et al. 1980; Gerhardt et al. 1990; Gerhardt et al. 2000), while low-frequency sounds are not (0 to 5 dB reduction) (Gerhardt et al. 1990; Querleu et al. 1989; Sohmer and Freeman 2001), there is even one report showing that low frequency sound can be amplified (Richards et al. 1992). Impulse noise of 169 dB peak SPL (Sound Pressure Level) in air was attenuated fairly modestly to 161 dB peak SPL near the head of a fetal sheep, indicating very little protection for the fetus against high level impulse sounds (Gerhardt et al. 2000).

The fetus is exposed to sounds not only from external sources, but also from the mothers own body. However, the natural intrauterine sound level consist mainly of very low-frequency noise (Benzaquen et al. 1990) and since sounds consists of a mixture of frequencies (between 20 Hz to 20 000 Hz) the total noise exposure in utero is to a large extent dependent on the sound exposure outside the abdomen.

It was not possible to assess the risk associated with noise exposure at different stages in pregnancy in this study. Experimental studies on pregnant guinea pigs and sheep that also have a fully developed auditory system in utero, show a risk of adverse effects of noise on the fetal

hearing during the later gestational ages, corresponding to the last trimester of pregnancy in humans (Cook et al. 1982; Gerhardt et al. 1999). However, these studies do not provide any information about the risk at different stages of pregnancy. Even though the auditory system is not fully developed until the 20th gestational week in humans, there is a possibility of adverse influence of noise on the hearing even during the embryonic period in the first trimester, when the auditory system is developing. The otocyst embryonic stem cells are able to produce hair-cell like cells (Woods et al. 2004) and it can be speculated that these progenitor cells might be able to react in the event of loud sounds. Isolated adult mammalian cochlear outer hair cells respond with a change in length when subjected to sound stimulation (Canlon et al. 1988) and each cell has a sharply toned frequency response (Brundin et al. 1989). If the progenitor hair cell possesses the same capacity there is a possibility of influence of loud sounds even in earlier stages of pregnancy. Since studies on noise effects on the fetus during early pregnancy are lacking it is impossible to clearly state at what time during pregnancy noise can be harmful to the fetus.

The main limitations of the study concern the exposure classification. Noise exposure was classified through a job-exposure-matrix, thereby introducing some misclassification of exposure. While the matrix accounts for variation in exposure between occupations, it does not account for variation within the occupation (Kromhout and Vermeulen 2001). Neither was it possible to specifically study exposure to low-frequency noise. While all these factors contribute to misclassification of exposure, it is not likely that this misclassification is differential. In addition, an equal non-differential misclassification of exposure in each exposure group will mainly lead to an attenuation of the risk in the middle and high exposure group, since the low exposure group is so large (80% of the full sample) compared with the middle exposed (19% of the full sample) and the highly exposed group (1% of the full sample) and even a minor

misclassification of the large low exposure group will lead to a lot of low exposure mothers to change group to the middle and high exposure group, leading to an attenuation of the association in these groups. The increased risk in the high exposure group >85dBA shown in this study exists despite of this misclassification and the true risk estimate might therefore be higher.

To limit the misclassification introduced by the fact that mothers stating the same occupation may be present at the workplace to a various degree, the cohort was subdivided into full-time working mothers, part-time working mothers and mothers absent from work during pregnancy. . Among full-time working mothers, high exposure to occupational noise was associated with an increased risk of hearing dysfunction. An increase in risk was also indicated in the part-time working mothers. There was no increased risk of hearing dysfunction in children whose mothers reported an exposed occupation in the beginning of the pregnancy, but were absent from work during pregnancy. The fact that the risk increased with presence at work, supports the hypothesis that occupational noise during pregnancy, and not the occupational title as such, is associated with an increased risk of hearing dysfunction.

A statistically significant increase of 5-8% was seen in the intermediate exposure group (75-84dBA) in the main analysis. The interpretation of this small increase is complex since the class is wide, with a range in exposure of 9dBA. Most occupations are likely to be found at the lower end of the interval (75-79 dBA), and an effect in the higher end (80-84 dBA) will have a small effect on the estimated risk for the whole intermediate group. Thus, the results for the intermediate group should not be interpreted as a null-result and that 75-84 dBA is safe level for pregnant women to work in. More studies are needed before the highest safe level can be established.

A wide range of hearing dysfunction diagnoses related to noise exposure was selected, mainly due to the fact that noise induced hearing dysfunctions in children are difficult to classify and thereby are sometimes classified into the selected “other” categories until a more precise diagnosis can be made. An inclusion of such a mixed group might also contribute to an inclusion of a few cases of hearing dysfunction that are unrelated to noise exposure. However, these “other” diagnoses (not including Tinnitus) contributed with very few cases (<10%) and were only included in the dichotomous analyses of all hearing dysfunctions and should therefore only introduce a minor misclassification of disease. In addition, the misclassification of hearing dysfunction in this study is most likely non-differential, since occupational exposures to noise during pregnancy is not an established risk factor for hearing dysfunction in children and it should therefore not make parents more prone to investigate their child’s hearing later in life.

In this study, only hearing dysfunction that was sufficient to prompt an investigation by a specialist could be studied. Thereby the analyses are based on few cases but the low numbers should not be taken as evidence that the problem should be of minor public health relevance. Rather is it possible that a larger number of cases of mild hearing dysfunction, not identified during childhood or not requiring specialist care, could be induced by occupational noise exposure. This was evident in the earlier three epidemiological studies on occupational noise during pregnancy and hearing dysfunction in children that measured hearing loss for each child by hearing thresholds instead of diagnoses of hearing dysfunction and thereby also could detect mild hearing loss even in a small sample (Daniel and Laciak 1982; Lalande et al. 1986; Rocha et al. 2007). Even mild hearing loss in children has been associated with increased social and emotional dysfunction among school children (Bess et al. 1998).

Future studies should investigate the association in different noise intervals in more detail and connect these with outcome measures of milder hearing dysfunction. The advice given today to pregnant women regarding occupational noise differs substantially between clinics and countries, due to the fact that research on this topic is lacking. This study will therefore provide important information to clinicians and policymakers and will contribute to more accurate advice and guidelines.

Conclusions

This nationwide population based study supports that occupational noise exposure during pregnancy is associated with future hearing dysfunction in children. Taken together with previous epidemiological and experimental studies as well as mechanistic data, the available data indicate that pregnant women should not be exposed to high levels of noise at work.

These results need confirmation in further studies. In addition, our results indicate a need to further study the effects of intermediate levels of occupational noise, peak values and leisure-time exposure, e.g. rock concerts. Although leisure-time activities are of much shorter duration, the exposure intensity may be very high.

References

Arbetsmiljöverket - the Swedish Work Environment Authority. 2012. Arbetsmiljön 2011 - the work environment 2011 (in swedish). Stockholm.

Armitage SE, Baldwin BA, Vince MA. 1980. The fetal sound environment of sheep. *Science* 208:1173-1174.

Benzaquen S, Gagnon R, Hunse C, Foreman J. 1990. The intrauterine sound environment of the human fetus during labor. *Am J Obstet Gynecol* 163:484-490.

Bess FH, Dodd-Murphy J, Parker RA. 1998. Children with minimal sensorineural hearing loss: Prevalence, educational performance, and functional status. *Ear Hear* 19:339-354.

Brundin L, Flock A, Canlon B. 1989. Sound-induced motility of isolated cochlear outer hair cells is frequency-specific. *Nature* 342:814-816.

Canlon B, Brundin L, Flock A. 1988. Acoustic stimulation causes tonotopic alterations in the length of isolated outer hair cells from guinea pig hearing organ. *Proc Natl Acad Sci U S A* 85:7033-7035.

Chordekar S, Kriksunov L, Kishon-Rabin L, Adelman C, Sohmer H. 2012. Mutual cancellation between tones presented by air conduction, by bone conduction and by non-osseous (soft tissue) bone conduction. *Hear Res* 283:180-184.

Cook RO, Konishi T, Salt AN, Hamm CW, Lebetkin EH, Koo J. 1982. Brainstem-evoked responses of guinea pigs exposed to high noise levels in utero. *Dev Psychobiol* 15:95-104.

Daniel T, Laciak J. 1982. [clinical observations and experiments concerning the condition of the cochleovestibular apparatus of subjects exposed to noise in fetal life]. *Rev Laryngol Otol Rhinol (Bord)* 103:313-318.

Försäkringskassan - the Swedish Social Insurance Agency. 2011. Midas sjukpenning och rehabilitering - sickleave and rehabilitation (in swedish). Stockholm.

Gerhardt KJ, Abrams RM, Oliver CC. 1990. Sound environment of the fetal sheep. *Am J Obstet Gynecol* 162:282-287.

Gerhardt KJ, Pierson LL, Huang X, Abrams RM, Rarey KE. 1999. Effects of intense noise exposure on fetal sheep auditory brain stem response and inner ear histology. *Ear Hear* 20:21-32.

Gerhardt KJ, Abrams RM. 2000. Fetal exposures to sound and vibroacoustic stimulation. *J Perinatol* 20:S21-30.

Gerhardt KJ, Abrams RM, Huang X, Griffiths SK, Peters AJ. 2000. Intra-abdominal sound pressure levels during impulse noise exposure in sheep. *Mil Med* 165:153-156.

Griffiths SK, Pierson LL, Gerhardt KJ, Abrams RM, Peters AJ. 1994. Noise induced hearing loss in fetal sheep. *Hear Res* 74:221-230.

Hackney C, Furness D. 2007. Noise and its effects, section: Cochlear pathophysiology in response to hazardous noise: John Wiley & Sons. Ltd.

Hepper PG, Shahidullah BS. 1994. Development of fetal hearing. *Arch Dis Child* 71:F81-87.

Huang X, Gerhardt KJ, Abrams RM, Antonelli PJ. 1997. Temporary threshold shifts induced by low-pass and high-pass filtered noises in fetal sheep in utero. *Hear Res* 113:173-181.

International Labour Office (ILO). 1990. International standard classification of occupations (isco-88). Geneva.

Kromhout H, Vermeulen R. 2001. Temporal, personal and spatial variability in dermal exposure. *Ann Occup Hyg* 45:257-273.

Lalande NM, Hetu R, Lambert J. 1986. Is occupational noise exposure during pregnancy a risk factor of damage to the auditory system of the fetus? *Am J Ind Med* 10:427-435.

OECD. 2014. Employment rate of working age population, sweden, women, 2014 q3, <http://data.Oecd.Org/emp/employment-rate.Htm#indicator-chart> date accessed 2015-02-13.

Pujol R, Lavigne-Rebillard M, Uziel A. 1990. Physiological correlates of development of the human cochlea. *Semin Perinatol* 14:275-280.

Querleu D, Renard X, Boutteville C, Crepin G. 1989. Hearing by the human fetus? *Semin Perinatol* 13:409-420.

Richards DS, Frentzen B, Gerhardt KJ, McCann ME, Abrams RM. 1992. Sound levels in the human uterus. *Obstet Gynecol* 80:186-190.

Rocha EB, Frasson de Azevedo M, Ximenes Filho JA. 2007. Study of the hearing in children born from pregnant women exposed to occupational noise: Assessment by distortion product otoacoustic emissions. *Braz J Otorhinolaryngol* 73:359-369.

Saunders JC, Chen CS. 1982. Sensitive periods of susceptibility to auditory trauma in mammals. *Environ Health Perspect* 44:63-66.

Sjostrom M, Lewne M, Alderling M, Willix P, Berg P, Gustavsson P, et al. 2013. A job-exposure matrix for occupational noise: Development and validation. *Ann Occup Hyg* 57:774-783.

Socialstyrelsen - the Swedish National Board of Health and Welfare. 2002. Utvärdering av det medicinska födelseregistret - evaluation of the swedish medical birth register (in swedish). Stockholm.

Socialstyrelsen - the Swedish National Board of Health and Welfare. 2013. Drg-statistik 2011 - en beskrivning av vårdkonsumtion i sverige (in swedish). Stockholm.

Sohmer H, Freeman S. 2001. The pathway for the transmission of external sounds into the fetal inner ear. *J Basic Clin Physiol Pharmacol* 12:91-99.

Statistiska centralbyrån - Statistics Sweden. 2011. Longitudinell integrationsdatabas för sjukförsäkrings- och arbetsmarknadsstudier (lisa), integrated database for labor market research 1990-2009 (in swedish). Stockholm.

Verbeek JH, Kateman E, Morata TC, Dreschler WA, Mischke C. 2012. Interventions to prevent occupational noise-induced hearing loss. *Cochrane Database Syst Rev*.

Woods C, Montcouquiol M, Kelley MW. 2004. Math1 regulates development of the sensory epithelium in the mammalian cochlea. *Nat Neurosci* 7:1310-1318.

Table 1. Baseline characteristics of the study participants and maternal occupational noise exposure during pregnancy.

	Occupational noise exposure ^a		
	<75dBA n (%)	75-84dBA n (%)	≥85dBA n (%)
<i>n</i> total sample (1,422,333) =	1,126,356	290,071	5,906
Mother's characteristics			
Age (quartiles)			
≤ 24 years	175,994 (15.6)	59,519 (20.5)	1,498 (25.4)
25-28 years	311,072 (27.6)	86,938 (30.0)	1,671 (28.3)
29-32 years	336,334 (29.9)	79,124 (27.3)	1,430 (24.2)
≥ 33 years	302,956 (26.9)	64,490 (22.2)	1,307 (22.1)
n (%) missing:	0 (0)	0 (0)	0 (0)
Smoking			
Non-smokers	938,088 (85.9)	225,991 (80.5)	4,434 (77.3)
Smokers, 1-9 cig per day	105,229 (9.6)	34,062 (12.1)	756 (13.2)
Smokers, ≥ 10 cig per day	49,190 (4.5)	20,549 (7.3)	545 (9.5)
n (%) missing:	33,849 (3.0)	9,469 (3.3)	171 (2.9)
Highest completed educational level			
Pre-high school	96,232 (8.7)	52,666 (18.7)	1,222 (21.3)
High school	610,581 (54.9)	133,051 (47.2)	3,593 (62.7)
University < 3 years	151,663 (13.6)	70,206 (24.9)	323 (5.6)
University ≥ 3 years	254,089 (22.8)	25,840 (9.2)	593 (10.4)
n (%) missing:	13,791 (1.2)	8,308 (2.9)	175 (3.0)
Nationality			
Swedish	1,047,984 (93.1)	257,149 (88.7)	5,348 (90.6)
Other European country	39,513 (3.5)	15,089 (5.2)	314 (5.3)
Outside Europe	16,682 (1.5)	9,445 (3.3)	99 (1.7)
Unknown	22,030 (2.0)	8,306 (2.9)	144 (2.4)
n (%) missing:	147 (0)	82 (0)	1 (0)
Working at beginning of pregnancy^b			
No	59,881 (5.9)	14,312 (5.5)	292 (5.4)
Part-time	377,291 (37.2)	91,079 (35.2)	1,251 (23.3)
Full-time	576,487 (56.9)	153,186 (59.2)	3,824 (71.3)
n (%) missing:	112,697 (10.0)	31,494 (10.9)	539 (9.1)
Leave of absence^c			
<20 days	515,358 (45.8)	106,403 (36.7)	1,845 (31.2)
20-153 days	513,628 (45.6)	151,814 (52.3)	3,214 (54.4)
>153 days	97,370 (8.6)	31,854 (11.0)	847 (14.3)
n (%) missing:	0 (0)	0 (0)	0 (0)
Family structure			
Married/living together with the father	1,052,832 (96.4)	266,783 (95.1)	5,490 (96.0)
n (%) missing:	34,455 (3.1)	9,563 (3.3)	186 (3.1)

Child's characteristics

Gender

Male	578,886 (51.4)	149,087 (51.4)	2,986 (50.6)
n (%) missing:	0 (0)	0 (0)	0 (0)

Parity

First child	500,459 (44.4)	119,747 (41.3)	2,596 (44.0)
Second child	413,151 (36.7)	103,068 (35.5)	2,035 (34.5)
Third child or more	212,746 (18.9)	67,256 (23.2)	1,275 (21.6)
n (%) missing:	0 (0)	0 (0)	0 (0)

Birth year

1986-1990	258,479 (23.0)	86,280 (29.7)	1,439 (24.4)
1991-1996	318,963 (28.3)	87,659 (30.2)	1,871 (31.7)
1997-2002	245,798 (21.8)	57,062 (19.7)	1,315 (22.3)
2003-2008	303,116 (26.9)	59,070 (20.4)	1,281 (21.7)
n (%) missing:	0 (0)	0 (0)	0 (0)

a Occupational noise exposure estimated through a job exposure matrix based on measurements at several work sites, subdividing the mother's occupation registered at the beginning of the pregnancy into three noise categories.

b Includes mothers who had reported an occupation at the beginning of pregnancy (gestational week 10), and divides them according to their answer on the separate presence-at-work-question (Do you work right now?: no/part-time/full-time).

c Number of days of leave of absence due to sick-leave or parental-leave reported to the Social Insurance Agency register during pregnancy. Including a summation of all days between the conception date (birthdate-days of gestation=conception date) and the birth date. Cut-off at the median level 20 days and at 153 days of absence, equivalent to the 90th percentile.

Table 2. Hazard ratios (HR) for maternal occupational noise exposure during pregnancy and hearing dysfunction in 1,422,333 children (1,320,195 children in the adjusted analyses)

Occupational noise exposure ^d	Hearing dysfunction, all ^a			Sensorineural hearing loss ^b			Tinnitus ^c		
	n cases	Crude ^e HR (CI 95%)	Adjusted ^f HR (CI 95%)	n cases	Crude ^e HR (CI 95%)	Adjusted ^f HR (CI 95%)	n cases	Crude ^e HR (CI 95%)	Adjusted ^f HR (CI 95%)
<75 dBA	9,813/ 9,001	1.00	1.00	6,688/ 6,176	1.00	1.00	2,548/ 2,285	1.00	1.00
75-84 dBA	2,790/ 2,545	0.96 (0.92, 1.00)	1.05 (1.00, 1.10)	1,962/ 1,782	1.01 (0.96, 1.06)	1.08 (1.03, 1.15)	706/ 645	0.89 (0.82, 0.97)	0.99 (0.90, 1.08)
≥85 dBA	65/ 60	1.20 (0.94, 1.53)	1.27 (0.99, 1.64)	46/ 42	1.26 (0.94, 1.68)	1.26 (0.93, 1.70)	16/ 14	1.10 (0.67, 1.80)	1.25 (0.74, 2.12)

a Hearing dysfunction all selected diagnoses, includes other diseases of inner ear (H83.3, H83.8), Sensorineural hearing loss and sensorineural mixed with conductive hearing loss (H90.3-7), Other hearing loss (H91.0, H91.2-3, H91.8) and other disorders of ear, not elsewhere classified, (H93.1, H93.2).

b Sensorineural hearing loss includes sensorineural hearing loss and mixed conductive and sensorineural ICD10: H90.3-7

c Tinnitus includes ICD10:H93.1

d Occupational noise exposure estimated through a job exposure matrix based on measurements at several work sites, classifying the mother's occupation registered at the beginning of the pregnancy (gestational week 10) into one of three noise categories.

e Crude analyses were restricted to all single births between 1986 and 2008 and excluding mothers in non-working categories (students, house wives, unemployed, refugees)

f Analyses adjusted for mother's age, smoking, education, family structure, nationality and child's, gender, birth year and parity. Including all single births between 1986 and 2008 and excluding mothers in non-working categories (students, house wives, refugees).

Table 3. HR in three separate analyses for maternal occupational noise exposure during pregnancy and hearing dysfunction^a, dividing employed mothers by work participation during pregnancy

Occupational noise exposure ^b dBA	Cases	Absent from work ^c		Cases	Working part-time ^d		Cases	Working full-time ^e	
		Crude n=181,170	Adjusted ^f n=164,598		Crude n=819,489	Adjusted ^f n=760,049		Crude n=362,572	Adjusted ^f n=343,712
		Crude/ adjust.	HR (CI 95%)		HR (CI 95%)	Crude/ adjust.		HR (CI 95%)	HR (CI 95%)
<75	1,223/ 1,116	1.00	1.00	5,749/ 5,243	1.00	1.00	2,363/ 2,222	1.00	1.00
75-84	410/ 368	0.89 (0.79, 1.00)	0.99 (0.88, 1.12)	1,804/ 1,645	1.00 (0.95, 1.06)	1.08 (1.02, 1.14)	447/ 450	0.97 (0.87, 1.07)	1.04 (0.93, 1.16)
≥ 85	7/ 7	0.60 (0.29, 1.26)	0.74 (0.35, 1.56)	40/ 37	1.26 (0.92, 1.71)	1.25 (0.91, 1.71)	16/ 14	1.91 (1.17, 3.13)	1.82 (1.08, 3.08)

a Hearing dysfunction (all selected diagnoses), includes other diseases of inner ear (H83.3, H83.8), Sensorineural hearing loss and sensorineural mixed with conductive hearing loss (H90.3-7), Other hearing loss (H91.0, H91.2-3, H91.8) and other disorders of ear, not elsewhere classified, (H93.1, H93.2).

b Occupational noise exposure estimated through a job exposure matrix based on measurements at several work sites, dividing the mother's occupation registered at the beginning of the pregnancy (gestational week 10) into three noise categories.

c Employed mothers with an absence of >153 days (90th percentile) during pregnancy or who reported "not working at the moment" during the registration interview at the prenatal care service in the beginning of pregnancy (gestational week 10).

d Mothers that reported that they worked part-time at the registration interview at the prenatal care service at the beginning of pregnancy (gestational week 10) or had more than 20 days (50th percentile) and less than 153 days (90th percentile) absence from work during pregnancy.

e Mothers that reported that they worked full-time at the beginning of pregnancy (gestational week 10) and that had less than 20 days (50th percentile) absence from work during pregnancy.

f Analyses adjusted for mothers age, smoking, education, family structure, nationality and child's, gender, birth year and parity.

Including all single births between 1986 and 2008 and excluding mothers in non-working categories (students, house wives, refugees).

Figure 1. Schematic figure showing the process of classifying an occupational free text variable into one out of three noise categories $<75\text{dBA}$, $75\text{-}84\text{dBA}$, $\geq 85\text{dBA}$.

Figure 1.

