



ENVIRONMENTAL HEALTH PERSPECTIVES

<http://www.ehponline.org>

Occupational Lead Exposure and Associations with Selected Cancers: The Shanghai Men's and Women's Health Study Cohorts

Linda M. Liao, Melissa C. Friesen, Yong-Bing Xiang, Hui Cai,
Dong-Hee Koh, Bu-Tian Ji, Gong Yang, Hong-Lan Li,
Sarah J. Locke, Nathaniel Rothman, Wei Zheng, Yu-Tang Gao,
Xiao-Ou Shu, and Mark P. Purdue

<http://dx.doi.org/10.1289/ehp.1408171>

Received: 24 January 2014

Accepted: 17 June 2015

Advance Publication: 19 June 2015

This article will be available in a 508-conformant form upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.



Occupational Lead Exposure and Associations with Selected Cancers: The Shanghai Men's and Women's Health Study Cohorts

Linda M. Liao,¹ Melissa C. Friesen,¹ Yong-Bing Xiang,² Hui Cai,³ Dong-Hee Koh,⁴ Bu-Tian Ji,¹
Gong Yang,³ Hong-Lan Li,² Sarah J. Locke,¹ Nathaniel Rothman,¹ Wei Zheng,³ Yu-Tang Gao,²
Xiao-Ou Shu,³ and Mark P. Purdue¹

¹Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland, USA; ²Department of Epidemiology, Shanghai Cancer Institute, Shanghai, China; ³Division of Epidemiology, Department of Medicine, Vanderbilt University, School of Medicine, Nashville, Tennessee, USA; ⁴National Cancer Control Institute, National Cancer Center, Goyang, Korea

Address correspondence to Linda M. Liao, Occupational and Environmental Epidemiology Branch, National Cancer Institute, 9609 Medical Center Dr, Rm 6-E632, MSC 9771, Bethesda, MD 20892 USA. Telephone: 240-276-7288. E-mail: Linda.Liao@nih.gov

Running title: Occupational lead exposure and cancer risk

Acknowledgments: The authors express their appreciation to the Shanghai residents who participated in the study and thank the research staff of the Shanghai Women's Health Study and the Shanghai Men's Health Study for their dedication and contributions to the study. This project was supported by grants from the US National Institutes of Health (R37 CA070867 to W. Zheng and R01 CA082729 and UM1 CA173640 to X.-O. Shu) and the Intramural Research Program contract N02 CP1101066.

Competing financial interests: The authors declare that they have no actual or potential competing financial interests.

Abstract

Background: Epidemiologic studies of occupational lead exposure have suggested increased risks of cancers of the stomach, lung, kidney, brain, and meninges; however, the totality of the evidence is inconsistent.

Objective: We investigated the relationship between occupational lead exposure and cancer incidence of these five sites in two prospective cohorts in Shanghai, China.

Methods: Annual job/industry-specific estimates of lead fume and lead dust exposure, derived from a statistical model combining expert lead intensity ratings with inspection measurements, were applied to the lifetime work histories of participants from the Shanghai Women's Health Study (SWHS; n=73,363) and the Shanghai Men's Health Study (SMHS; n=61,379) to estimate cumulative exposure to lead fume and lead dust. These metrics were then combined into an overall occupational lead exposure variable. Cohort-specific relative hazard rate ratios (RRs) and 95% confidence intervals (CI) comparing exposed to unexposed participants were estimated using Cox proportional hazards regression and combined by meta-analysis.

Results: The proportion of SWHS and SMHS participants with estimated occupational lead exposure was 8.9% and 6.9%, respectively. Lead exposure was positively associated with meningioma risk in women only (n= 38 unexposed and 9 exposed cases, RR = 2.4; 95% CI: 1.1, 5.0), particularly with above-median cumulative exposure (RR = 3.1; 95% CI: 1.3, 7.4). However, all 12 meningioma cases among men were classified as unexposed to lead. We also observed non-significant associations with lead exposure for cancers of the kidney (n= 157 unexposed and 17 ever exposed cases; RR = 1.4; 95% CI: 0.9, 2.3) and brain (n= 67 unexposed and 10 ever exposed cases; RR = 1.8; 95% CI: 0.7, 4.8) overall.

Conclusions: Our findings, though limited by small numbers of cases, suggest that lead is associated with risk of several cancers in women and men.

Introduction

Lead is a metal commonly used in many industrial settings worldwide, and an important environmental pollutant. The occurrence of lead in the environment has decreased greatly in recent decades because of the elimination of most leaded gasoline; however, occupational exposures continue primarily via lead in the storage battery industry and lead pigments in paints (IARC 2006). Occupations that have had frequent high exposure include battery-production workers, battery-recycling workers, foundry workers, lead chemical workers, lead smelter and refinery workers, leaded-glass workers, pigment workers, construction workers, and radiator-repair workers. In most developed countries, stricter controls have reduced environmental and occupational exposures to lead; however, lead exposure continues to be an issue in developing countries with rapid industrialization, such as China (Gottesfeld and Pokhrel 2011; IARC 2006). Lead as a gasoline additive is a large contributor to environmental lead exposure and developing countries, such as China, have generally lagged behind developed countries in banning lead in gasoline. Leaded gasoline was eventually banned in 1999 in China, and gradually phased out over the 2000's, however, other environmental sources continue to contribute to lead exposure in China. The occupational exposure limit (OEL) for lead and inorganic compounds of lead in China was set in 1979 based on maximum allowable concentrations of 0.05 mg/m^3 for lead dust and 0.03 mg/m^3 for lead fumes (Liang et al. 1995). The OELs remained at the same levels but were based on time-weighted averages from 2002 on, which is similar to the exposure standards for lead in the United States.

High lead exposure is known to be harmful, particularly for children; established health effects include damage to the brain and nervous system, gastrointestinal problems, anemia, liver and

kidney damage, fertility problems and developmental delays (Abadin et al. 2007). Lead is also a suspected carcinogen, with inorganic lead compounds currently designated by the International Agency for Research on Cancer (IARC) as probably carcinogenic (Group 2A), based on limited evidence in humans and sufficient evidence in animals (IARC 2006). Organic lead compounds were designated by IARC as not classifiable as to carcinogenicity (Group 3) due to inadequate evidence. Epidemiologic evidence for carcinogenicity in workers exposed to inorganic lead suggest associations with cancers of the stomach, lung, kidney, brain, and meninges, although the totality of the evidence is inconsistent (IARC 2006; Rousseau et al. 2007; Steenland and Boffetta 2000). Very few of the past studies have evaluated occupational lead exposure among women, although gender differences in lead exposure and metabolism have been observed (Vahter et al. 2007). There is thus a need for additional well-designed epidemiologic studies including men and women to resolve the question of whether lead is a carcinogen (Ward et al. 2010). To that end, we investigated the association between occupational lead exposure and risk of cancers of the stomach, lung, kidney, brain, and meninges in two large prospective cohort studies of women and men in Shanghai, China.

Methods

Study Population

The Shanghai Women's Health Study (SWHS) and the Shanghai Men's Health Study (SMHS) are two population-based prospective cohort studies based in Shanghai, China. The rationale, design and methods of both studies have been described in detail previously (Shu et al. 2015; Zheng et al. 2005). Briefly, using a roster provided by the community office, 81,170 permanent female residents aged 40-70 years were approached for the SWHS study between 1996 and 2000

(Zheng et al. 2005); and 83,033 permanent male residents aged 40-74 years were approached for the SMHS between 2002 to 2006 (Shu et al. 2015). Of the 81,170 eligible women, 75,221 participated in the SWHS study for an overall response rate of 92.7%. It was determined afterwards that 279 of these women did not meet the age eligibility requirements and were excluded, resulting in a cohort of 74,941 women. Of the 83,033 eligible men, 61,480 participated in the SMHS study for an overall response rate of 74.0%. An additional 14 men were lost to follow-up, resulting in a cohort of 61,466 men. An additional exclusion of prevalent cancers at baseline was also applied (n=1578 women and n=0 men; as having a prior history of cancer was a part of the exclusion criteria for participation in the SMHS), leaving 73,363 women and 61,466 men in this analysis. In-person interviews were administered at baseline to obtain information on demographics, lifestyle and dietary habits, medical history, and other characteristics, including lifetime occupational history. All study participants provided written informed consent prior to interview and the study protocols were approved by the institutional review boards of all participating institutions (National Cancer Institute, Vanderbilt University and the Shanghai Cancer Institute).

Cohort members are followed for cancer occurrence through in-person follow-up surveys administered every 2–3 years and annual record linkage with the Shanghai Cancer Registry and Vital Statistics Unit. For the SWHS, the response rates for follow-up (i.e. = number of responders/ number of surviving cohort members) for the first (2000–2002), second (2002–2004), third (2004–2007), and fourth (2008–2011) in-person follow-up surveys were 99.8%, 98.7%, 96.7%, and 92.0%, respectively. For the SMHS, the response rates for the first (2004–2008) and second (2008–2011) follow-up surveys were 97.6% and 93.6%, respectively. Cohort

members known to have permanently moved out of Shanghai or who cannot be contacted in three consecutive follow-ups are considered lost to follow-up. All cancer diagnoses are verified through home visits and medical chart review to ensure pathological confirmation. The cancer sites of interest for this project were stomach cancer (ICD-9 codes 151.0-151.9), lung cancer (ICD-9 codes 162.0-162.9), kidney cancer (ICD-9 code 189.0), brain cancer (ICD-9 code 191), and meningioma (ICD-9 codes 192.1, 192.3 and 225.2; ICDO-3 codes 9530-9539). First-incident cancers of each site of interest were identified through December 31, 2009 in SWHS and December 31, 2010 in SMHS.

Lead Exposure Assessment

Study participants provided a lifetime occupational history, which included all jobs held for at least one year, with specific details on job title, type of business, factory name, description of the work tasks and the employment dates. Occupational history records were then assigned job and industry codes based on the Standard Chinese Classification of Industries and Occupations for the Third National Population Census of 1982 (CSA and CSICSC 1989).

Lead fume and lead dust were estimated and evaluated separately because they can vary in their particle size and composition and thus may have differing bioavailability and health effects.

Lead fume is created by high temperature processes that form fine particulate through condensation of airborne lead vapor, whereas lead dust is formed from mechanical processes that develop both fine and large particles (Needleman 1992). Because both are measured using the same sampling and analytical processes, they are distinguished here by using industrial hygienists' judgment of the expected form of lead exposure based on the work activities and lead

source. The lead fume and lead dust measures described below do not differentiate lead exposures based on the chemical form (metallic, inorganic, organic) or solubility, which will vary by lead source.

Details on the development of job/industry-specific estimates of exposure to lead fumes and lead dust have been previously reported for SWHS and were used for both cohorts in this analysis (Koh et al. 2012). In brief, population-based job exposure matrices (JEMs) were developed to provide expert-based estimates of the probability and intensity of exposure to lead fumes and lead dust for the job and industry codes reported in the study participants' occupational histories. A mixed-effects model was developed separately for lead fume and lead dust to combine the expert ratings of the respective lead intensity metric with its associated inspection measurements (20,084 lead fume measurements; 5383 lead dust measurements) collected by the Shanghai Center for Disease Control and Prevention between 1954 and 2000. Annual job/industry-specific estimates of lead fume and lead dust exposure were calculated from the fixed-effects terms for the JEM intensity ratings and calendar year, and from the random-effects terms for job and industry from their respective mixed models. Job/industry-specific estimates were calculated only for job/industry combinations that met a strict exposure definition based on the JEM probability ratings (job probability =high *or* industry probability =high and job probability = low, medium or high); all other job/industry combinations were assigned 0 exposure. We chose this exposure definition to emphasize specificity over sensitivity, as recommended by Kromhout and Vermeulen for rare exposures to minimize bias from exposure misclassification (Kromhout and Vermeulen 2001). The lead fume and lead dust models were applied to both cohorts to estimate annual occupational lead fume and lead dust exposure for each study participant; the

annual estimates were then summed over each participant's working life to obtain separate cumulative exposure estimates for lead fume and for lead dust. Median values for cumulative lead fumes ($0.33 \text{ mg/m}^3\text{-years}$) and lead dust ($1.32 \text{ mg/m}^3\text{-years}$) were determined by the combined distribution of the exposed participants across both cohorts. Due to small numbers, the cumulative estimates of occupational lead fume and lead dust exposure were also combined into an overall lead exposure variable. Subjects were further categorized as "never" when no cumulative lead dust or lead fume exposure was assigned, "low" when cumulative exposures for either lead dust and/or lead fume \leq median and neither lead dust and/or lead fume was $>$ median, and "high" when either cumulative exposure $>$ median for lead dust or lead fume (Table 1). Cumulative lead estimates incorporating 10- and 20-year lags were also constructed. Using the same mixed-effect models, calibrated JEM estimates that used the fixed-effects terms but not the random-effects terms were also calculated and applied to both cohorts; however, it was determined that this alternate estimate of cumulative exposure was essentially collinear with the job/industry-specific estimates (Pearson correlation = 0.94 for lead fume and Pearson correlation = 0.99 for lead dust). Thus, we chose to present the more refined job/industry-specific estimates of lead exposure (Koh et al. 2012). We refer the readers to the paper by Koh et al. for a more detailed discussion of the models and a review of the sensitivity analyses conducted.

Statistical Analyses

Cox proportional hazards regression, with age as the time scale, was used to estimate cohort-specific hazard rate ratios (RRs) and 95% confidence intervals (95% CI) for the association between lead exposure and risk of each cancer site, with adjustment for potential confounders: education level (elementary school or less, middle school, high school, and professional

education/college or higher), family income level (study-specific, see Table 2 for definition), lifetime pack-years of cigarette use (study-specific- SWHS: never smoker, former smoker ≤ 7.4 pack-years, former smoker > 7.4 pack-years, current smoker ≤ 7.4 pack-years, current smoker > 7.4 pack-years; SMHS: never smoker, former smoker ≤ 22.2 pack-years, former smoker > 22.2 pack-years, current smoker ≤ 22.5 pack-years, current smoker > 22.5 pack-years) and menopause status (defined as absence of menstruation for ≥ 12 months; SWHS only). All confounders were baseline characteristics. Models additionally adjusted for body mass index (continuous) and alcohol consumption (continuous) yielded virtually identical results are not presented. Study participants with missing data (no occupational history provided) were treated as a separate category in the analysis. We then calculated summary RR estimates from cohort-specific results through meta-analysis using a random effects model. We tested for potential RR heterogeneity between cohorts using Cochran's Q statistic. Cohort-specific analyses were conducted using SAS, version 9.3 (SAS Institute Inc., Cary, NC) and the meta-analysis was conducted using STATA, version 13.0 (StataCorp LP, College Station, TX).

Results

The SWHS and SMHS analytic cohorts included 73,363 women (mean follow-up: 10.8 years) and 61,466 men (6.4 years) respectively. SMHS participants tended to report a higher level of education than SWHS participants, but reported lower household income (Table 2). Smoking and alcohol consumption was much more common in the male cohort than the female cohort. In both cohorts, around half of women (50.4%) and men (51.5%) reported working as manual laborers as their longest occupation during their occupational history at baseline. Overall, the

proportion of study participants (women and men combined) identified as ever exposed to lead fumes was 7.4 %, and ever exposed to lead dust was 3.1 %. Lead exposure concentrations in these cohorts have decreased considerably from 1965 to 2000 (Table 2) (Koh et al. 2012). The exposure prevalence was slightly higher in the female cohort (lead fumes, 7.9%; lead dust, 3.7%) than the male cohort (2.3%; 6.7%). The median first year of lead fume or dust exposure in the female cohort was slightly earlier than that of the male cohort (1972 vs. 1975).

The three most commonly reported jobs (based on number of exposed person-years) that were exposed to lead fume or dust in the female cohort (“Install/assemble electric/electronic equipment”, 42.1% of exposed person-years; “Welders”, 11.4%; “Other electric/Electronic equipment install/maintenance”, 7.9%) were a different pattern than those reported in the SMHS cohort (“Install/assemble electric/electronic equipment”, 14.2%; “Rolling Mill & Machinery Operators”, 12.0%; “Smelters”, 8.4%) (Supplemental Material, Table S1). Five of the top ten jobs exposed to lead fume or dust were the same in both cohorts. Males, however, appeared to be exposed to lead through a wider range of occupations than females, as the ten most commonly-exposed occupations reported by males account for only 68% of exposed person-years vs. 83% among females.

A total of 1,918 incident first-primary cancers of interest were identified during follow-up in the two cohorts, including 634 stomach cancers, 974 lung cancers, 174 kidney cancers, 77 brain cancers, and 59 meningiomas (47 females and 12 males). We observed non-significant associations with ever exposure to lead dust or fumes for cancers of the kidney (n = 157 unexposed and 17 ever exposed cases; RR = 1.4; 95% CI: 0.9, 2.3) and brain (n = 67 unexposed

and 10 ever exposed cases; RR = 1.8; 95% CI: 0.7, 4.8) (Table 3). A further elevated risk of kidney cancer was observed among those with high cumulative exposure to lead based on the combined lead metric (n = 12 exposed cases; RR = 1.8; 95% CI: 1.3, 7.4). There was no clear dose-response association with brain cancer, as the association was null among those with high exposure. Similar results for kidney and brain cancer were observed for separate analyses of lead dust and fume. In our cohort of females, an elevated risk with ever exposure to lead was observed for meningioma (n = 38 unexposed cases and 9 ever exposed cases; RR = 2.4; 95% CI: 1.1, 5.0; Table 3). This association was particularly strong for high cumulative lead exposure based on the combined lead metric (n = 38 unexposed cases and 6 exposed cases; RR = 3.1; 95% CI: 1.3, 7.4) and was also present in separate analyses of both lead dust and lead fume. We could not compute an association with meningioma in the male cohort as none of the 12 cases was assessed as having lead exposure. Although a hazard ratio for meningioma was not estimable for the male cohort, given the lack of exposed male cases it is unlikely that lead exposure was positively associated with risk in this study population. (see Supplemental Material, Tables S2 and S3).

We observed null findings for cancers of the lung and stomach (Table 3), however, there was evidence of heterogeneity between cohorts which could be partly due to the lack of an association in the female cohort. When the cohorts were analyzed separately (Table 4), high cumulative lead exposure based on the combined lead metric was non-significantly associated with risk of lung cancer in the male cohort (n = 460 unexposed cases and 35 exposed cases; RR = 1.4; 95% CI: 0.98, 2.0) but not the female cohort (n = 440 unexposed cases and 17 exposed cases; RR = 0.8; 95% CI: 0.5, 1.3; high exposure heterogeneity p-value = 0.06). Similarly, a

suggestive elevated risk of stomach cancer was associated with high lead exposure based on the combined lead metric (n = 293 unexposed cases and 23 exposed cases; RR = 1.6; 95% CI: 1.0, 2.4) in the male cohort that was not observed among the female cohort (n = 292 unexposed cases and 12 exposed cases; RR = 0.8; 95% CI: 0.5, 1.5; high exposure heterogeneity p-value = 0.07). Associations of lung and stomach cancer with high (> median) exposure in the male cohort were stronger for lead fumes than lead dust (see Supplemental Material, Table S3). Also note that males were more likely to be exposed to lead fumes than lead dust based upon the number of exposed cases.

Evaluation of lead exposure with 10- and 20-year time lags were generally consistent with our risk estimates and did not change our findings (see Supplemental Material, Tables S2 and S3). The lack of change in our risk estimates is consistent with the trend described by Koh et al. of lead fume and lead dust concentrations declining over time in this population (Koh et al. 2012). Thus, excluding recent lead exposure had little impact on risk estimates and further demonstrates that most lead exposure occurred in the distant past in these cohorts.

Discussion

We found evidence of an association between exposure to lead dust or fumes and an increased subsequent risk of meningioma in the female cohort, with higher cumulative exposure associated with higher risk. We were unable to evaluate this association in the male cohort due to a lack of exposed male cases, but it is unlikely that lead exposure was positively associated with meningioma risk in the male cohort. Non-significant associations with lead exposure were observed overall for cancers of the kidney and brain. The association with brain cancer appeared

to be limited to the female cohort. In addition, an elevated risk of lung and stomach cancer was observed with high lead exposure in the male cohort, but no association was observed in the female cohort. Our findings suggest that lead is associated with risk of several cancers, but are limited by small numbers of cases, particularly for kidney and brain cancer.

In 2006, when IARC classified lead as a probable carcinogen, the epidemiological evidence was the most consistent for stomach cancer, with elevated kidney and lung cancer risks observed in some but not all studies (IARC 2006). Evidence from human studies was considered limited; thus, several studies conducted since the IARC monograph have attempted to further evaluate the risks for these cancers. Some of these investigations have reported associations with meningioma and brain cancer that are consistent with our findings. A case-control study of death certificate data among U.S. women observed an association between jobs involving occupational lead exposure and risk of meningioma (n=161 cases) (Cocco et al. 1999). A more recent U.S. case-control study (n= 197 meningiomas) found evidence of an association between cumulative lead exposure and meningioma risk among males but not females, however, they observed a consistently increased risk of meningioma among a subset of lead-exposed individuals with susceptible genotypes of *ALAD2*, a gene influencing lead bioavailability (Bhatti et al. 2009; Rajaraman et al. 2006). Navas-Acien et al. reported an increased relative risk of meningioma with occupational lead exposure among Swedish males (n= 848 meningiomas) but had an insufficient number of exposed females in their cohort study to compute a risk estimate (Navas-Acien et al. 2002). In an occupational cohort of Finnish women, an increased relative risk of brain and central nervous system cancers (including 298 meningiomas) with high levels of occupational lead exposure was reported (Wesseling et al. 2002). Findings from several case-

control (Cocco et al. 1998) and cohort (Anttila et al. 1996; Ilychova and Zaridze 2012; van Wijngaarden and Dosemeci 2006; Wesseling et al. 2002) studies suggest an association between lead and brain cancer.

We also observed associations between lead exposure and an increased risk of kidney cancer, primarily in the male cohort, and increased risks of lung and stomach cancer in the male cohort only. Findings from other recent epidemiological studies, few of which included women, have been inconsistent for these cancers. A multicenter case-control study of both men and women conducted in Central and Eastern Europe observed an elevated risk of renal cell carcinoma (RCC) with high occupational lead exposure (Boffetta et al. 2011). A subsequent analysis from this study further reported that genetic variation in *ALAD* may modify the lead-RCC association (van Bommel et al. 2011). Southard et al. conducted a nested case-control study of Finnish male smokers and observed an increased relative risk of RCC (Odds Ratio = 2.0; 95% CI: 1.0, 3.9) with higher blood lead concentrations (Southard et al. 2012). A higher blood lead concentration was also associated with increased lung cancer risk in a separate Finnish cohort (Anttila et al. 1995). In contrast, a retrospective cohort study of male lead-exposed workers in Australia did not find increased risk of any cancers previously linked to lead exposure (Gwini et al. 2012). Rousseau et al. (Rousseau et al. 2007) evaluated the risk of 11 types of cancer and lead exposure in a case-control study of men in Canada and only found an association between organic lead and stomach cancer. In a subsequent study, Wynant et al. (Wynant et al. 2013) did not find an association between lead and lung cancer in a pooled analysis that included the previous study. We observed stronger associations with lung and stomach cancer among males with the lead fume metrics than with the lead dust metrics. Exposure to lead fumes was more common than

exposure to lead dust overall among study participants. Previous studies did not discern between the two lead measures, however, it is possible that the finer particulate produced by lead fumes is more readily absorbed through inhalation or ingestion than lead dust (NTP 2011).

The mechanisms through which lead may increase cancer risk remain unclear. Inhalation and oral ingestion are the two primary routes through which lead enters the body. Thus, both the lungs and stomach are organs that come into initial contact with lead exposure, but lead can enter the bloodstream and affect other organs. The brain and nervous system are especially sensitive to the potential toxic effects of lead due to its ability to pass through the blood-brain barrier (Inskip et al. 1995). The high reabsorptive activity of the renal proximal tubules also lends itself to the accumulation and uptake of lead in the kidney (IARC 2006). It has been suggested that lead may act through indirect mechanisms to facilitate carcinogenic effects of other DNA-damaging agents, as experimental studies have shown very little or no mutagenicity for the main forms of lead (IARC 2006; Winder and Bonin 1993). Therefore, it has been proposed that lead may play a role in carcinogenesis through mechanisms that involve oxidative damage, induction of apoptosis, altered cell signaling pathways, inhibition of DNA synthesis and repair of damage, and interaction with DNA-binding proteins (IARC 2006; Restrepo et al. 2000; Silbergeld 2003).

Differential associations by cohort/gender were observed between lead exposure and several cancer sites. There are several potential explanations for the observed differences. Gender-related differences in lead metabolism may have played a role in the differential associations observed (Bjorkman et al. 2000; Vahter et al. 2007). The different pattern of lead-exposed occupations reported between genders may have led to differences by cohort in workplace lead

exposure intensity and duration not captured by our exposure assessment. Although the most common occupation reported among males and females was the same, this occupation (installing or assembling electronic equipment) accounted for 42% of the exposure among females and only 14% among males. This would suggest that males were exposed to a wider range of occupational sources of lead than females. There is also a large difference in smoking prevalence between the two cohorts, although our findings are adjusted for pack-years smoked. We also note that the follow-up period for the male cohort was shorter than that of the female cohort (Table 2). Lastly, we cannot rule out the possibility that our gender-specific findings are due to chance given that we had small numbers of cases for some cancer sites; or due to residual confounding associated with gender.

Our study has many strengths, including the inclusion of two large population-based prospective cohorts with high response rates at recruitment and follow-up. Both cohorts captured extensive information on potential confounders, allowing us to adjust for potential confounding risk factors, such as smoking, that industry-based cohorts are rarely able to account for. Detailed lifetime occupational histories were also collected before cancer development, eliminating the potential for recall bias. Although we combined results from two large cohort studies, our findings were limited by small numbers of exposed cases for some cancer sites, most notably brain cancer and meningioma, and a relatively short follow-up period for the male cohort, thus limiting our statistical power to detect overall and sex-specific associations and can result in unstable risk estimates that should be interpreted with caution..

Another strength of our study is the ability to capture important time trends in lead exposure by using estimates based on a framework that calibrated an expert-based JEM using inspection measurements collected from Shanghai-area worksites. As is the case with all JEMs, the exposure assessment approach was unable to account for variation in exposure among workers who worked within a particular occupational-industrial grouping and time period (Koh et al. 2012). Another strength of the exposure assessment was that it distinguished between lead fume and lead dust, which partly captures differences in lead composition and bioavailability. This differentiation introduces some exposure misclassification because it is based on expert opinion to characterize the lead source based on observing the work activities while collecting the exposure measurements and, for the JEM, it is based on the expected source and work activities for a given occupation or industry. Further differentiation of lead exposure based on solubility or composition was not possible in this study given the limited occupational information and the wide variety but low prevalence of the lead-exposed occupations and industries represented in these cohorts. In addition, exposure estimates for other metals that may be correlated with occupational lead were not available; as a consequence, we are unable to rule out potential confounding by other metals. In regards to meningioma, there are currently no established associations between other metals and meningioma that would be expected to confound our observed association.

Conclusion

In conclusion, our findings, though limited by small numbers of cases, suggest that lead exposure is associated with increased risk of several cancer sites, in particular, meningioma, brain cancer and kidney cancer. The association between lead and meningioma and brain cancer that we

observed among women in our study underscores the importance of including women in future studies evaluating the carcinogenicity of lead.

References

- Abadin H, Ashizawa A, Stevens YW, Lladós F, Diamond G, Sage G, et al. 2007. Toxicological profile for lead. Atlanta (GA):U.S. Department of Health and Human Services, Public Health Service.
- Anttila A, Heikkilä P, Pukkala E, Nykyri E, Kauppinen T, Hernberg S, et al. 1995. Excess lung cancer among workers exposed to lead. *Scandinavian journal of work, environment & health* 21:460-469.
- Anttila A, Heikkilä P, Nykyri E, Kauppinen T, Pukkala E, Hernberg S, et al. 1996. Risk of nervous system cancer among workers exposed to lead. *Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine* 38:131-136.
- Bhatti P, Stewart PA, Hutchinson A, Rothman N, Linet MS, Inskip PD, et al. 2009. Lead exposure, polymorphisms in genes related to oxidative stress, and risk of adult brain tumors. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 18:1841-1848.
- Bjorkman L, Vahter M, Pedersen NL. 2000. Both the environment and genes are important for concentrations of cadmium and lead in blood. *Environmental health perspectives* 108:719-722.
- Boffetta P, Fontana L, Stewart P, Zaridze D, Szeszenia-Dabrowska N, Janout V, et al. 2011. Occupational exposure to arsenic, cadmium, chromium, lead and nickel, and renal cell carcinoma: A case-control study from central and eastern Europe. *Occupational and environmental medicine* 68:723-728.
- Cocco P, Dosemeci M, Heineman EF. 1998. Brain cancer and occupational exposure to lead. *Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine* 40:937-942.
- Cocco P, Heineman EF, Dosemeci M. 1999. Occupational risk factors for cancer of the central nervous system (cns) among us women. *American journal of industrial medicine* 36:70-74.

- Gottesfeld P, Pokhrel AK. 2011. Review: Lead exposure in battery manufacturing and recycling in developing countries and among children in nearby communities. *Journal of occupational and environmental hygiene* 8:520-532.
- Gwini S, MacFarlane E, Del Monaco A, McLean D, Pisaniello D, Benke GP, et al. 2012. Cancer incidence, mortality, and blood lead levels among workers exposed to inorganic lead. *Annals of epidemiology* 22:270-276.
- IARC. 2006. Inorganic and organic lead compounds. In: *Iarc monographs on the evaluation of carcinogenic risks to humans*. Lyon, France:International Agency for Research on Cancer.
- Ilychova SA, Zaridze DG. 2012. Cancer mortality among female and male workers occupationally exposed to inorganic lead in the printing industry. *Occupational and environmental medicine* 69:87-92.
- Inskip PD, Linet MS, Heineman EF. 1995. Etiology of brain tumors in adults. *Epidemiologic reviews* 17:382-414.
- Koh DH, Bhatti P, Coble JB, Stewart PA, Lu W, Shu XO, et al. 2012. Calibrating a population-based job-exposure matrix using inspection measurements to estimate historical occupational exposure to lead for a population-based cohort in shanghai, china. *Journal of exposure science & environmental epidemiology*.
- Kromhout H, Vermeulen R. 2001. Application of job-exposure matrices in studies of the general population: Some clues to their performance. *Eur Respir Rev* 11:80.
- Liang YX, Gang BQ, Gu XQ. 1995. The development of occupational exposure limits for chemical substances in china. *Regulatory toxicology and pharmacology* : RTP 22:162-171.
- Navas-Acien A, Pollan M, Gustavsson P, Plato N. 2002. Occupation, exposure to chemicals and risk of gliomas and meningiomas in sweden. *American journal of industrial medicine* 42:214-227.
- Needleman HL. 1992. *Human lead exposure*. Boca Raton:CRC Press.
- NTP. 2011. Lead and lead compounds. Report on carcinogens : carcinogen profiles / US Dept of Health and Human Services, Public Health Service, National Toxicology Program 12:251-256.

- Rajaraman P, Stewart PA, Samet JM, Schwartz BS, Linet MS, Zahm SH, et al. 2006. Lead, genetic susceptibility, and risk of adult brain tumors. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 15:2514-2520.
- Restrepo HG, Sicard D, Torres MM. 2000. DNA damage and repair in cells of lead exposed people. *American journal of industrial medicine* 38:330-334.
- Rousseau MC, Parent ME, Nadon L, Latreille B, Siemiatycki J. 2007. Occupational exposure to lead compounds and risk of cancer among men: A population-based case-control study. *American journal of epidemiology* 166:1005-1014.
- Shu XO, Li H, Yang G, Gao J, Cai H, Takata Y, et al. 2015. Cohort profile: The shanghai men's health study. *International journal of epidemiology*.
- Silbergeld EK. 2003. Facilitative mechanisms of lead as a carcinogen. *Mutation research* 533:121-133.
- Southard EB, Roff A, Fortugno T, Richie JP, Jr., Kaag M, Chinchilli VM, et al. 2012. Lead, calcium uptake, and related genetic variants in association with renal cell carcinoma risk in a cohort of male finnish smokers. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 21:191-201.
- Steenland K, Boffetta P. 2000. Lead and cancer in humans: Where are we now? *American journal of industrial medicine* 38:295-299.
- Vahter M, Akesson A, Liden C, Ceccatelli S, Berglund M. 2007. Gender differences in the disposition and toxicity of metals. *Environmental research* 104:85-95.
- van Bommel DM, Boffetta P, Liao LM, Berndt SI, Menashe I, Yeager M, et al. 2011. Comprehensive analysis of 5-aminolevulinic acid dehydrogenase (alad) variants and renal cell carcinoma risk among individuals exposed to lead. *PloS one* 6:e20432.
- van Wijngaarden E, Dosemeci M. 2006. Brain cancer mortality and potential occupational exposure to lead: Findings from the national longitudinal mortality study, 1979-1989. *International journal of cancer Journal international du cancer* 119:1136-1144.

- Ward EM, Schulte PA, Straif K, Hopf NB, Caldwell JC, Carreon T, et al. 2010. Research recommendations for selected iarc-classified agents. *Environmental health perspectives* 118:1355-1362.
- Wesseling C, Pukkala E, Neuvonen K, Kauppinen T, Boffetta P, Partanen T. 2002. Cancer of the brain and nervous system and occupational exposures in finnish women. *Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine* 44:663-668.
- Winder C, Bonin T. 1993. The genotoxicity of lead. *Mutation research* 285:117-124.
- Wynant W, Siemiatycki J, Parent ME, Rousseau MC. 2013. Occupational exposure to lead and lung cancer: Results from two case-control studies in montreal, canada. *Occupational and environmental medicine* 70:164-170.
- Zheng W, Chow WH, Yang G, Jin F, Rothman N, Blair A, et al. 2005. The shanghai women's health study: Rationale, study design, and baseline characteristics. *American journal of epidemiology* 162:1123-1131.

Table 1. Method for assigning categories of total lead exposure (never, low, high) from estimates of lead dust and lead fume exposure.

Combined Lead Metric		Lead Fume		
		0	≤ median ^a	> median ^a
Lead Dust	0	Never	Low	High
	≤ median ^a	Low	Low	High
	> median ^a	High	High	High

^aMedian values for cumulative lead fumes (0.33 mg/m³-years) and lead dust (1.32 mg/m³-years) were determined by the combined distribution of the exposed participants across both cohorts

Table 2. Selected characteristics of the Shanghai Women’s Health Study (SWHS) and the Shanghai Men’s Health Study (SMHS) cohorts.

Characteristics^a	SWHS (N=73,363)	SMHS (N=61,466)
Age at baseline (years); mean (range)	52.0 (40-70)	55.4 (40 -75)
Education		
Elementary school or less; N (%)	15,687 (21.4)	4,083 (6.7)
Middle school; N (%)	27,270 (37.2)	20,330 (33.5)
High school; N (%)	20,490 (27.9)	21,856 (36.1)
Professional education/college or higher; N (%)	9,903 (13.5)	14,334 (23.7)
Income ^b		
Low; N (%)	11,813 (16.1)	33,845 (55.2)
Lower middle; N (%)	28,063 (38.3)	21,539 (35.1)
Upper middle; N (%)	20,599 (28.1)	4,597 (7.5)
High; N (%)	12,872 (17.6)	1,358 (2.2)
Occupation ^c		
Professional, administrator; N (%)	21,026 (28.8)	16,308 (26.6)
Clerical worker; N (%)	15,198 (20.8)	13,469 (21.9)
Manual laborer; N (%)	36,862 (50.4)	31,619 (51.5)
Lifetime pack-years of cigarette use ^d		
Never; N (%)	71,320 (97.2)	18,669 (30.4)
Former – Low; N (%)	158 (0.2)	3,689 (6.0)
Former – High; N (%)	141 (0.2)	3,065 (5.0)
Current – Low; N (%)	875 (1.2)	18,012 (29.3)
Current – High; N (%)	868 (1.2)	18,024 (29.3)
Alcohol consumption		
Ever; N (%)	1,654 (2.3)	20,728 (33.7)
Grams per day; mean (range)	9.1 (0-150)	11.8 (0-608)

Characteristics ^a	SWHS (N=73,363)	SMHS (N=61,466)
Body Mass Index (kg/m ²); mean (range)	24.0 (13-49)	23.7 (12-40)
Menopausal status		
Pre-menopausal; N (%)	37,457 (51.1)	N/A
Post-menopausal; N (%)	35,891 (48.9)	N/A
Lead Dust		
Never; N (%)	70,378 (95.9)	57,241 (93.2)
Ever; N (%)	2,709 (3.7)	4,138 (6.7)
Year first exposed among exposed participants; median (range) ^c	1972 (1941-1999)	1976 (1943-2005)
Cumulative exposure (mg/m ³ *yr) among exposed participants; median (range) ^c	1.56 (0.01-11.2)	1.03 (0.006-7.8)
Annual exposure of exposed participants by time period (mg/m ³); median (range) ^c		
1930-1959	0.15 (0.04-0.52)	0.11 (0.02-0.30)
1960s	0.14 (0.03-0.52)	0.10 (0.02-0.30)
1970s	0.10 (0.03-0.37)	0.07 (0.01-0.21)
1980s	0.11 (0.03-0.42)	0.08 (0.01-0.24)
1990s	0.056 (0.010-0.32)	0.04 (0.003-0.18)
2000s	0.012 (0.008-0.030)	0.009 (0.002-0.025)
Lead Fume		
Never; N (%)	67,280 (91.7)	59,962 (97.6)
Ever; N (%)	5,807 (7.9)	1,417 (2.3)
Year first exposed among exposed participants; median (range) ^c	1972 (1945-1999)	1975 (1941-2005)
Cumulative exposure (mg/m ³ *yr) among exposed participants; median (range) ^c	0.29 (0.003-6.6)	0.46 (0.001-11.0)
Annual exposure of exposed participants by time period (mg/m ³); median (range) ^c		
1930-1959	0.028 (0.015-0.24)	0.063 (0.009-0.33)
1960s	0.025 (0.014-0.24)	0.049 (0.008-0.33)
1970s	0.020 (0.013-0.22)	0.038 (0.005-0.28)
1980s	0.020 (0.005-0.24)	0.30 (0.003-0.30)
1990s	0.005 (0.002-0.080)	0.008 (0.001-0.038)
2000s	0.003 (0.001-0.019)	0.004 (0.001-0.038)

Characteristics ^a	SWHS (N=73,363)	SMHS (N=61,466)
Lead Dust and Fume		
Never; N (%)	66,813 (91.1)	57,123 (93.0)
Ever; N (%)	6,274 (8.9)	4,256 (6.9)
Year first exposed; median (range) ^c	1972 (1941-1999)	1975 (1941-2005)
Years of Follow-Up; mean (range)	10.8 (0.1-13)	6.4 (0.1 - 9)

^aNumber of subjects with missing data noted for education (n=13 women, 863 men); income (n=16 women, 127 men); occupation (n=277 women, 70 men); smoking (n=1 woman, 7 men); alcohol drinking (n=0 women, 1 man); menopausal status (n=15 women); and lead dust and/or fume exposure data (n=276 women, 87 men).

^bIncome cutpoints were as follows: SWHS: <10,000 [low], 10,000-<20,000 [lower middle], 20,000-<30,000 [upper middle], and 30,000+ [high] yuan/year per household. SMHS: <1000 [low], 1000-<2000 [lower middle], 2000-<3000 [upper middle], and 3000+ [high] yuan/month.

^cLongest occupation reported during their occupational history

^dLifetime pack-years of cigarette use cutpoints were as follows: SWHS: never smoker, former - low smoker ≤ 7.4 pack-years, former - high smoker > 7.4 pack-years, current - low smoker ≤ 7.4 pack-years, current - high smoker > 7.4 pack-years; SMHS: never smoker, former -low smoker ≤ 22.2 pack-years, former – high smoker > 22.2 pack-years, current –low smoker ≤ 22.5 pack-years, current – high smoker > 22.5 pack-years.

^eResults presented are among exposed subjects only

Table 3. Meta-analysis summary estimates for the association between lead and cancer.

	Kidney		Lung		Stomach		Brain		Meningioma (SWHS only) ^a	
	Cases	RR (95% CI) ^b	Cases	RR (95% CI)	Cases	RR (95% CI)	Cases	RR (95% CI)	Cases	RR (95% CI)
Lead Dust^c										
Never	168	1.0	948	1.00	619	1.00	72	1.00	42	1.00
Ever	6	1.3 (0.6, 2.8)	26	0.9 (0.6, 1.4)	15	0.8 (0.4, 1.4)	5	2.3 (0.9, 5.8)	5	2.9 (1.1, 7.3)
Low	1	0.8 (0.1, 6.0)	13	0.9 (0.3, 3.1)	7	0.8 (0.4, 1.7)	2	2.0 (0.5, 8.3)	1	1.5 (0.2, 10.6)
High	5	2.3 (0.8, 6.7)	13	0.8 (0.5, 1.3)	8	0.8 (0.4, 1.7)	3	2.6 (0.8, 8.2)	4	3.8 (1.4, 10.7)
Lead Fumes^c										
Never	157	1.00	902	1.00	587	1.00	68	1.00	38	1.00
Ever	17	1.5 (0.9, 2.5)	72	0.9 (0.5, 1.8) ^d	47	1.0 (0.5, 2.0) ^d	9	1.8 (0.8, 4.1)	9	2.6 (1.2, 5.4)
Low	6	1.2 (0.4, 3.4)	22	0.8 (0.4, 1.3)	14	0.7 (0.4, 1.2)	6	2.9 (1.2, 6.7)	4	2.2 (0.8, 6.3)
High	11	1.8 (0.9, 3.7)	50	1.1 (0.6, 1.9)	33	1.2 (0.6, 2.4)	3	1.1 (0.3, 3.5)	5	3.0 (1.2, 7.6)
Combined Lead Dust and Fumes^e										
Never	157	1.0	900	1.0	585	1.0	67	1.0	38	1.0
Ever	17	1.4 (0.9, 2.3)	74	0.9 (0.5, 1.7) ^d	49	1.0 (0.5, 1.9) ^d	10	1.8 (0.7, 4.8)	9	2.4 (1.1, 5.0)
Low	5	1.0 (0.4, 2.5)	22	0.7 (0.4, 1.2)	14	0.7 (0.4, 1.2)	7	3.1 (1.0, 9.1)	3	1.7 (0.5, 5.4)
High	12	1.8 (1.0, 3.3)	52	1.1 (0.6, 1.9)	35	1.2 (0.6, 2.2)	3	1.0 (0.3, 3.2)	6	3.1 (1.3, 7.4)

Abbreviations: RR, relative hazard rate ratio; CI, confidence interval

^aMeningioma results are only from SWHS as there were no exposed meningioma cases (n=12) in SMHS

^bAdjusted for education, income level, lifetime pack-years of cigarette use, and menopause status (women only)

^cLevels of lead exposure: low: ≤ median; high: >median

^dTest for heterogeneity (Cochran's Q statistic) indicated heterogeneity at p< 0.05

^eLevels of combined lead exposure: low: exposed to lead dust or lead fumes, but not high exposure (≤ median); high: at least one high exposure (> median) to lead dust or lead fumes

Table 4. Cohort-specific associations between lead exposure and cancer risk.

	Kidney				Lung				Stomach				Brain			
	<u>SWHS</u>		<u>SMHS</u>		<u>SWHS</u>		<u>SMHS</u>		<u>SWHS</u>		<u>SMHS</u>		<u>SWHS</u>		<u>SMHS</u>	
	Cases	RR (95% CI) ^a	Cases	RR (95% CI)	Cases	RR (95% CI)	Cases	RR (95% CI)	Cases	RR (95% CI)	Cases	RR (95% CI)	Cases	RR (95% CI)	Cases	RR (95% CI)
Combined Lead Dust and Fumes^b																
Never	76	1.0	81	1.0	440	1.0	460	1.0	292	1.0	293	1.0	34	1.0	33	1.0
Ever	8	1.3 (0.6, 2.6)	9	1.6 (0.8, 3.1)	27	0.7 (0.5, 1.0)	47	1.2 (0.9, 1.7)	19	0.7 (0.4, 1.1)	30	1.4 (0.9, 2.0)	8	2.6 (1.2, 5.6)	2	0.9 (0.2, 3.8)
Low	4	1.3 (0.5, 3.5)	1	0.5 (0.1, 3.2)	10	0.6 (0.3, 1.1)	12	1.0 (0.5, 1.7)	7	0.6 (0.3, 1.2)	7	0.9 (0.4, 1.9)	6	4.2 (1.8, 10.1)	1	1.2 (0.2, 8.5)
High	4	1.2 (0.5, 3.4)	8	2.3 (1.1, 4.7)	17	0.8 (0.5, 1.3)	35	1.4 (0.98, 2.0)	12	0.8 (0.5, 1.5)	23	1.6 (1.03, 2.4)	2	1.2 (0.3, 5.0)	1	0.7 (0.1, 5.4)

Abbreviations: RR, relative hazard rate ratio; CI, confidence interval

^aAdjusted for education, income level, lifetime pack-years of cigarette use, and menopause status (women only).

^bLevels of lead exposure: low: exposed to lead dust or lead fumes, but not high exposure (\leq median); high: at least one high exposure ($>$ median) to lead dust or lead fumes.